

Complex Variables & Transforms (20A54302)

II - B.TECH & I- SEM

Prepared by:

Dr. B. NAGABHUSHANAM REDDY, Professor
Department of H &S



VEMU INSTITUTE OF TECHNOLOGY

(Approved By AICTE, New Delhi and Affiliated to JNTUA, Ananthapuramu)

Accredited By NBA(EEE, ECE & CSE) & ISO: 9001-2015 Certified Institution Near

Pakala, P.Kothakota, Chittoor- Tirupathi Highway

Chittoor, Andhra Pradesh-517 112

Web Site: www.vemu.org

Unit – 1

Complex - analysis

- **Function of Complex Variable/ Differentiation:**

If for each value of the complex variable $Z = X + iY$ in a given region 'R', we have one or more values of $w = f(z) = u + iv$, Then W is said to be a function of 'Z', and we have $w = f(z) = u + iv$.

Where u and v are real and imaginary parts of $f(z)$. $z = x + iy$
and

$f(z) = u(x, y) + iv(x, y)$ is a complex function.

- **Continuity of a Function:**

Let $f(z)$ is said to be continuous function at $z = z_0$ if

$$\lim_{z \rightarrow z_0} f(z) = f(z_0)$$

- **Differentiability of a Function:**

A function $f(z)$ is said to be differentiable at $z = z_0$ if

exists. It is denoted by $\lim_{\Delta z \rightarrow 0} \left(\frac{f(z + \Delta z) - f(z)}{\Delta z} \right)$ $f'(z_0)$

$$\text{i.e. } f'(z_0) = \lim_{\Delta z \rightarrow 0} \left(\frac{f(z + \Delta z) - f(z)}{\Delta z} \right)$$

- **Analytical**

Function:

The complex function $f(z)$ is said to be analytical function at $z=a$ if the function $f(z)$ has derivative at $z=a$ and neighbourhood of $z=a$.

Example:

1. Let $f(z) = z^2$ $f'(z) = 2z$

At $z=0$, $f'(z) = 2(0) = 0$ (finite) $f(z)$

has derivative at $z=0$

Finally $f(z)$ is called **analytical** function.

1

2. Let $f(z) =$

z

$-\frac{1}{z^2}$

$f'(z) =$ At $z=0$, $f'(z) =$

$\frac{-1}{(0)^2} = \infty$

$f(z)$ has no

derivative at $z=0$

Finally $f(z)$ is called **not analytical** function.

- **Singular Point:**

Let $z=a$ is said to be singular point if the function $f(z)$ is not analytical at $z=a$.

Example:

$f(z) = \frac{1}{z}$, $f'(z) = \infty$ $z = 0$ is called singular point.

- **Cauchy – Riemann Equations in Cartesian co-ordinates:**

- If $f(z)$ is continuous in some neighbourhood of z and differentiable at z then the first order partial derivatives satisfy the equations $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ and $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$ at the point z which are called the Cauchy-Riemann equations.

proof:

Let $f(z) = u+iv$ be an analytical function

By definition of analytical function, $f(z)$ has derivative.

$$\text{i.e. } f'(z) = \lim_{\Delta z \rightarrow 0} \left(\frac{f(z+\Delta z) - f(z)}{\Delta z} \right) \text{ exists (finite)}$$

$$1) \ z = x+iy \quad f(z) = u+iv \quad f(z) = u(x,y)+iv(x,y)$$

$$2) \ z = x+iy \quad \Delta z = \Delta x + i \Delta y \quad 3) \ f(z + \Delta z) = ?$$

$$z + \Delta z = x+iy + \Delta x + i \Delta y$$

$$z + \Delta z = (x + \Delta x) + i(y + \Delta y)$$

$$f(z + \Delta z) = u(x + \Delta x, y + \Delta y) + iv(x + \Delta x, y + \Delta y)$$

$$f'(z) = \lim_{\Delta x + i \Delta y \rightarrow 0} \left(\frac{[u(x + \Delta x, y + \Delta y) + iv(x + \Delta x, y + \Delta y)] - [u(x,y) + iv(x,y)]}{\Delta x + i \Delta y} \right) \rightarrow \textcircled{1}$$

$$\Delta x + i \Delta y \rightarrow 0$$

We know that $\Delta x + i \Delta y = 0 + i0$

$$x = 0, \Delta y = 0$$

Case (1) If $\Delta y = 0$, put $\Delta y = 0$ in ①.

$$f'(z) = \lim_{\Delta x \rightarrow 0} \left(\frac{[u(x+\Delta x, y)+iv(x+\Delta x, y)) - [u(x, y)+iv(x, y)]]}{\Delta x} \right) = \lim$$

$$f'(z) = \left(\lim_{\Delta x \rightarrow 0} \frac{[u(x+\Delta x, y)-u(x, y)]}{\Delta x} + \lim_{\Delta x \rightarrow 0} \frac{i[v(x+\Delta x, y)-v(x, y)]}{\Delta x} \right)$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \rightarrow \textcircled{2}$$

Case (2) If $\Delta x = 0$, put $\Delta x = 0$ in ①

$$f'(z) = \lim_{\Delta y \rightarrow 0} \left(\frac{[u(x, y+\Delta y)+iv(x, y+\Delta y)) - [u(x, y)+iv(x, y)]]}{i\Delta y} \right)$$

$$f'(z) = -i \lim_{\Delta y \rightarrow 0} \frac{[u(x, y+\Delta y)-u(x, y)]}{\Delta y} + \lim_{\Delta y \rightarrow 0} \frac{i[v(x, y+\Delta y)-v(x, y)]}{\Delta y}$$

$$f'(z) = -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \rightarrow \textcircled{3}$$

Equate ② & ③

Compare the real and imaginary parts

$$\begin{aligned} \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} &= -i \frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \\ \left\{ \begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} &= -\frac{\partial v}{\partial x} \end{aligned} \right\} \end{aligned} \quad (\text{If } ux = vy \text{ and } uy = -vx)$$

These are **Cauchy – Riemann** Equations in **Cartesian** co-ordinate System.

Cauchy – Riemann Equations in Polar co-ordinates:

Let $z=x+iy$

We know that $x=r\cos\theta$,

$$y=r\sin\theta$$

$$z = r\cos\theta + ir\sin\theta$$

$$z = r(\cos\theta + i\sin\theta) = re^{i\theta}$$

$$f(z)=u+iv \quad f(re^{i\theta}) = u(r, \theta)+iv(r, \theta) \rightarrow \textcircled{1}$$

Differentiate $\textcircled{1}$ w.r.t 'r',

$$f'(re^{i\theta}) e^{i\theta} = \frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \rightarrow \textcircled{2}$$

Differentiate $\textcircled{1}$ w.r.t 'θ',

$$f'(re^{i\theta}) (-ie^{i\theta}) = \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} \rightarrow \textcircled{3}$$

Substitute $\textcircled{2}$ in $\textcircled{3}$, We get

$$\left[\frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r} \right] = \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta} r e^{i\theta} \quad r i e^{i\theta} = \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta}$$

$$\frac{\partial u}{\partial r} - r \frac{\partial v}{\partial r} = \frac{\partial u}{\partial \theta} + i \frac{\partial v}{\partial \theta}$$

ir

ir

Lets compare real and imaginary parts

$$\frac{\partial u}{\partial \theta} = -r \frac{\partial v}{\partial r}$$

$$\frac{\partial v}{\partial \theta} = r \frac{\partial u}{\partial r}$$

These are **Cauchy – Riemann** Equations in **Polar** co-ordinate System. **Examples**

1) Show that $f(z) = xy + iy$ is not analytical

Solution : Given , $f(z) = xy + iy$

$$f(z) = u + iv \quad u = xy$$

$$v = y$$

$$\frac{\partial u}{\partial x} = y, \quad \frac{\partial v}{\partial x} = 0$$

$$\frac{\partial u}{\partial y} = x, \quad \frac{\partial v}{\partial y} = 1$$

$$\frac{\partial u}{\partial x} \neq \frac{\partial v}{\partial y}$$

$$\frac{\partial u}{\partial y} \neq -\frac{\partial v}{\partial x}$$

It doesn't **not satisfies C-R equations** and hence its **not an analytical function**.

2) Show that $f(z) = 2xy + i(x^2 - y^2)$ is not analytical function. Solution: Given $f(z) = 2xy + i(x^2 - y^2)$

$$f(z) = u + iv$$

$$u = 2xy \quad v = x^2 - y^2$$

$$\frac{\partial u}{\partial x} = 2y, \quad \frac{\partial v}{\partial x} = 2x$$

$$\frac{\partial u}{\partial y} = 2x, \quad \frac{\partial v}{\partial y} = -2y$$

$$\frac{\partial u}{\partial x} \neq \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} \neq -\frac{\partial v}{\partial x}$$

It doesn't **not** satisfies **C-R** equations and hence its **not an analytical** function.

3) Test the analyticity $f(z) = e^x(\cos y - i \sin y)$ and also find the $f'(z)$ Solution: Given $f(z) = e^x \cos y - i e^x \sin y$

$$f(z) = u + iv \quad u = e^x \cos y$$

$$v = -e^x \sin y$$

$$\frac{\partial u}{\partial x} = e^x \cos y, \quad \frac{\partial v}{\partial x} = -e^x \sin y$$

$$\frac{\partial u}{\partial y} = -e^x \sin y, \quad \frac{\partial v}{\partial y} = -e^x \cos y$$

$$\frac{\partial u}{\partial x} \neq \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} \neq -\frac{\partial v}{\partial x}$$

$f(z)$ is **not analytical** function and the $f'(z)$ does not exist.

4) Show that $f(z) = z \bar{z}^2$ is not analytical function

Solution : Given $f(z) = z \bar{z}^2$

$$f(z) = (x+iy)(x+iy) = (x+iy) [\sqrt{x^2 + y^2}]^2$$

$$f(z) = x(x^2 + y^2) + iy(x^2 + y^2) \quad f(z) =$$

$$u + iv$$

$$u = x(x^2 + y^2) = x^3 + xy^2 \quad v = y(x^2 + y^2) = x^2y + y^3$$

$$\frac{\partial u}{\partial x} = 3x^2 + y^2, \quad \frac{\partial v}{\partial x} = 2xy$$

$$\frac{\partial u}{\partial y} = 2xy, \quad \frac{\partial v}{\partial y} = x^2 + 3y^2$$

$$\frac{\partial u}{\partial x} \neq \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} \neq -\frac{\partial v}{\partial x}$$

$f(z)$ is **not analytical** function

5) Show that $w = \log z$ is an analytical function and also find $\frac{dw}{dz}$

Solution : Given $w = \log z$

$$\text{put } z = re^{i\theta}$$

$$\begin{aligned} i\theta &= \log r + \log e^{i\theta} \\ w &= \log re^{i\theta} \end{aligned}$$

$$= \log r + i\theta \log e$$

$$f(z) = w = \log r + i\theta = u + iv$$

$$u = \log r \quad v = \theta$$

$$\frac{\partial u}{\partial r} = \frac{1}{r}, \quad \frac{\partial v}{\partial r} = \theta$$

$$\frac{\partial u}{\partial \theta} = 0, \quad \frac{\partial v}{\partial \theta} = 1$$

$$r \frac{\partial u}{\partial r} = \frac{\partial v}{\partial \theta} \quad \& \quad \frac{\partial u}{\partial \theta} = -r \frac{\partial v}{\partial r}$$

$$r \left(\frac{1}{r} \right) = 1 \quad \& \quad 0 = 0 \quad \text{It is an **analytical** function } f(z) \\ = u+iv$$

$$f(re^{i\theta}) = u(r, \theta) + iv(r, \theta)$$

differentiate on both sides w.r.t 'r'

$$f'(re^{i\theta}) e^{i\theta} = \frac{\partial u}{\partial r} + i \frac{\partial v}{\partial r}$$

$$f'(z) e^{i\theta} = \frac{1}{r} + i \frac{\partial v}{\partial r}$$

$$f'(z) = \frac{1}{re^{i\theta}} = \frac{1}{z}$$

6) Show that $f(x) = \sin z$ is an analytical function everywhere in the complex plane

Solution : Given $f(x) = \sin z$

$$f(x) = \sin(x+iy) \quad f(x) = \sin x$$

$$\cos(iy) + \sin(iy) \cos x \quad f(X) = \sin x$$

$$\cosh y + i \sinh y \cos x \quad f(x) = u+iv$$

$$u = \sin x \cosh y \quad v = \sinh y \cos x$$

$$\frac{\partial u}{\partial x} = \cos x \cosh y, \quad \frac{\partial u}{\partial y} = \sin x \sinh y$$

$$= \sin x \sinh y, \quad = \cosh y \cos x \quad \& \quad \text{It is an **analytical** function}$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

7) Test the analyticity of the function $f(z) = e^x (\cos y + i \sin y)$ and find $f'(z)$. Solution : Given , $f(z) = e^x (\cos y + i \sin y) = u + iv$

$$u = e^x \cos y \quad v = e^x \sin y$$

$$\cos y, \quad \frac{\partial u}{\partial x} = e^x \quad \frac{\partial v}{\partial x} = e^x \sin y$$

$$\sin y \quad \frac{\partial u}{\partial y} = -e^x \quad \frac{\partial v}{\partial y} = e^x \cos y$$

$$\& \text{ It is an **analytical** function} \quad \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$f(z) = u + iv$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial y} = e^x \cos y + i e^x \sin y$$

$$f'(z) = e^x (\cos y + i \sin y)$$

$$f'(z) = e^x i e^y = e^{(x+iy)}$$

$$f'(z) = e^z$$

8) Determine P such that the function $f(z) = \frac{1}{2} \log (x^2 + y^2) + i \tan^{-1} \left(\frac{y}{x} \right)$ be an analytical function.
Solution :

$$\text{Given, } f(z) = \frac{1}{2} \log(x^2 + y^2) + i \tan^{-1}\left(\frac{px}{y}\right)$$

It is an analytical function, It

satisfies the C-R equation

$$u = \frac{1}{2} \log(x^2 + y^2) \quad \tan^{-1}\left(\frac{px}{y}\right)$$

$$\frac{\partial u}{\partial x} = \frac{1}{2} \frac{1}{x^2 + y^2} 2x, \quad \frac{\partial v}{\partial x} = \frac{1}{1 + \left(\frac{px}{y}\right)^2} \frac{p}{y}$$

$$\frac{\partial u}{\partial y} = \frac{1}{2} \frac{1}{x^2 + y^2} 2y$$

$$\frac{\partial v}{\partial y} = \frac{1}{1 + \left(\frac{px}{y}\right)^2} px \left(\frac{-1}{y^2}\right)$$

$$\frac{\partial v}{\partial y} = \frac{y^2}{y^2 + \left(\frac{px}{y}\right)^2} \left(\frac{-px}{y^2}\right)$$

$$\text{similarly: } \frac{\partial v}{\partial x} = \frac{py}{p^2x^2 + y^2} \quad \frac{\partial v}{\partial y} = \frac{-px}{y^2 + p^2x^2},$$

By given $f(z)$ is an analytical function, $f(z)$ satisfies C-R equations.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\frac{x}{x^2 + y^2} = \frac{-px}{y^2 + p^2x^2}$$

Comparing the equations we get:

$$p = -1$$

9) Prove that function $f(z)$ defined by $f(z) = -R$ equations are satisfied at the origin, yet $f'(0)$ does not exist.

$$\text{Solution: Given } f(z) = \frac{x^3(1+i) - y^3(1-i)}{x^2 + y^2}$$

i) To show that $f(z)$ is continuous at $z=0$

let $\lim_{z \rightarrow 0} f(z) = \lim_{x \rightarrow 0} \frac{x^3(1+i) - y^3(1-i)}{x^2+y^2}$ (given $f(0) = 0$) $\frac{x^3(1+i) - y^3(1-i)}{x^2+y^2}$, $z \neq 0$ and $f(0)$ is continuous and C

$$\lim_{x \rightarrow 0} \frac{x(1+i)}{x^2}$$

$$f(z) = f(z) =$$

$$\lim_{x \rightarrow 0} x(1+i) = 0 = f(0)$$

$f(z)$ is

continuous

ii) To show that C-R equations are satisfied at origin

$$f(z) = \frac{x^3 + x^3 i - y^3 + i y^3}{x^2 + y^2} = \frac{x^3 - y^3}{x^2 + y^2} + \frac{i(x^3 + y^3)}{x^2 + y^2} f(z)$$

$$= u + iv$$

$$u = \frac{x^3 - y^3}{x^2 + y^2}, \quad v = \frac{x^3 + y^3}{x^2 + y^2}$$

$$v =$$

$$\frac{\partial u}{\partial x} = \lim_{x \rightarrow 0} \frac{u(x,0) - u(0,0)}{x}$$

$$\frac{\partial u}{\partial x} = \lim_{x \rightarrow 0} \frac{x-0}{x} \Rightarrow \lim_{x \rightarrow 0} 1 = 1$$

$$\frac{\partial u}{\partial x} = 1$$

$$\frac{\partial u}{\partial y} = \lim_{y \rightarrow 0} \frac{u(0,y) - u(0,0)}{y}$$

$$\frac{\partial u}{\partial y} = \lim_{y \rightarrow 0} \frac{-y-0}{y} \Rightarrow \lim_{y \rightarrow 0} -1 = -1$$

$$\frac{\partial u}{\partial y} = -1$$

$$\frac{\partial v}{\partial x} = \lim_{x \rightarrow 0} \frac{v(x,0) - v(0,0)}{x}$$

$$\frac{\partial v}{\partial x} = \lim_{x \rightarrow 0} \frac{x-0}{x} = \lim_{x \rightarrow 0} 1 = 1$$

$$\frac{\partial v}{\partial x} = 1$$

$$\frac{\partial v}{\partial y} = \lim_{y \rightarrow 0} \frac{v(0,y) - v(0,0)}{y}$$

$$\frac{\partial v}{\partial y} = \lim_{y \rightarrow 0} \frac{y-0}{y} = \lim_{y \rightarrow 0} 1 = 1$$

$$\frac{\partial v}{\partial y} = 1$$

$$\text{C} - \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \lim_{x \rightarrow 0} \frac{\partial u}{\partial y} = - \lim_{y \rightarrow 0} \frac{\partial v}{\partial x}$$

R Equations are satisfied at origin **iii)** To

show that $f'(z)$ does not exist at origin

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z}$$

$$f'(z) =$$

$$y \rightarrow 0 \quad z$$

$$\frac{(\quad)}{(\quad)}$$

$$x^3(1+i-y^2) + 3y(1-i)^2$$

—

$$\frac{3(1+i)-0}{2+y^2}$$

$$\frac{(\quad)}{(\quad)}$$

$$f'(z) = \lim_{x \rightarrow 0} \frac{f(x+iy) - f(x)}{iy}$$

$$\lim_{x \rightarrow 0} \frac{x}{x}$$

$$f'(z) =$$

$$= \lim_{x \rightarrow 0} \frac{1+i-3}{1+i-3} = 1+i \quad (\text{Finite})$$

$f'(z)$ Exists

At $y = mx$

$$f'(z) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z} = \lim_{x \rightarrow 0} \frac{\frac{x^3(1+i) - m^3x^3(1-i)}{x^2 + x^2m^2}}{x+imx}$$

$$f'(z) = \lim_{x \rightarrow 0} \frac{x^3[(1+i) - m^3(1-i)]}{x^2(1+m^2)x(1+im)}$$

$$f'(z) = \lim_{x \rightarrow 0} \frac{[(1+i) - m^3(1-i)]}{(1+m^2)(1+im)} = \frac{[(1+i) - m^3(1-i)]}{(1+m^2)(1+im)}$$

$f'(z) =$ **(Infinite) $f'(z)$ depends upon the 'm' value, so that the $f'(z)$ does not exist at origin**

Part – B

Laplace Equations

the equation of the form $\frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 0$ or $\frac{\partial^2 \varphi}{\partial x^2} + \frac{\partial^2 \varphi}{\partial y^2} = 0$

Harmonic Function

The function u and v are said to be harmonic, if it satisfies Laplace Equations
i.e

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

or

$$\frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2} = 0$$

Milne – Thomson Method

When u is given find f(z) :

1) To find $\frac{\partial u}{\partial x}$ and $\frac{\partial u}{\partial y}$

2) To find $f'(z) = u + iv$

Differentiate w.r.t 'x' we get

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} - i \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial y} \right) \quad (\text{From C-R equation})$$

$$f'(z) =$$

$$\frac{\partial u}{\partial x} = \phi_1(z_1, 0) \quad \frac{\partial u}{\partial y} = \phi_2(z_2, 0) \quad f'(z) =$$

$$\phi_1(z_1, 0) - i \phi_2(z_2, 0)$$

$$\text{Integrate w.r.t 'z'} \quad f(z) = \phi_1(z_1, 0) dz - i \phi_2(z_2, 0) dz$$

+ c When v is given find f(z):

$$1) \text{ To find } \frac{\partial v}{\partial y} \text{ and } \frac{\partial v}{\partial x}$$

$$2) \text{ To find } f(z) = u + iv$$

Differentiate w.r.t 'x', we get

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$f'(z) = \frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

$$\frac{\partial v}{\partial y} = \phi_1(z_1, 0)$$

$$\frac{\partial v}{\partial x} = \phi_2(z_2, 0) \quad (\text{From C-R equation})$$

$$f'(z) = \phi_1(z_1, 0) + i \phi_2(z_2, 0)$$

$$\text{Integrate w.r.t 'z' } f(z) = \int [\phi_1(z_1, 0) + i \phi_2(z_2, 0)] dz + c$$

- 1) Construct an analytical function $f(z)$ when $u = x^3 - 3xy^2 + 3x + 1$ is given

$$\frac{\partial u}{\partial x} = 3x^2 - 3y^2 + 3$$

$$\frac{\partial u}{\partial y} = -6xy$$

Solution:

By Milne Thomson Method

$$f(z) = u + iv$$

$$\frac{\partial u}{\partial x} = \phi_1(z, 0) = 3z^2 + 3$$

$$\frac{\partial u}{\partial y} = \phi_2(z, 0) = -6z$$

$$\frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = \phi_1(z, 0) - i \phi_2(z, 0) = - \left(\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \right) \quad \text{6(z) (0) = 0}$$

$$f'(z) = \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} = - \frac{\partial v}{\partial x}$$

$$f'(z) = \phi_1(z, 0)$$

$$\text{Integrate w.r.t 'z' } f(z) = \int \phi_1(z, 0) dz + c$$

$$= \int (3z^2 + 3 - 0) dz + c$$

$$= \int (3z^2 + 3) dz + c$$

$$= \frac{3z^3}{3} + 3z + c$$

$$f(z) = z^3 + 3z + c$$

2) Construct an analytical function $f(z)$ when $u = \sin x \cosh y$ is given

Solution: $= \cos x \sinh y$

$$= \sin x \sinh y$$

By Milne Thomson

Method

$f(z) = u + iv$

$$\frac{\partial u}{\partial x} = \phi_1(z, 0) = \cos z(1) = \cos z$$

$$\frac{\partial u}{\partial y} = \phi_2(z, 0) = \sin z(0)$$

$$\frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} = \phi_1(z, 0) - i \phi_2(z, 0)$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} = 0$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \quad \& \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$f'(z) = \cos z$

Integrate w.r.t 'z' $f(z) =$

$$\int [\phi_1(z, 0) - i \phi_2(z, 0)] dz + c$$

$$\int \cos z dz + c \quad \mathbf{f(z) = \sin z + c}$$

3) Find the analytical function $f(z) = u + iv$ if $u + v = \frac{\sin 2x}{\cosh 2y - \cos 2x}$

Solution:

$$u + v = \frac{\sin 2x}{\cosh 2y - \cos 2x}$$

$$f(z) = u + iv$$

$$if(z) = u - v$$

$$(1 + i)f(z) = (u - v) + i(u + v)$$

$$f(z) = u + iv$$

$$\frac{\partial v}{\partial x} = \frac{[\cosh 2y - \cos 2x] 2 \cos 2x - \sin 2x [0 + 2 \sin 2x]}{[\cosh 2y - \cos 2x]^2}$$

$$\frac{\partial v}{\partial x} = \frac{2 \cos 2x \cosh y - 2 \cos^2 2x - 2 \sin^2 2x}{[\cosh 2y - \cos 2x]^2}$$

$$\frac{\partial v}{\partial x} = \frac{2 \cos 2x \cosh y - 2}{[\cosh 2y - \cos 2x]^2}$$

$$\frac{\partial v}{\partial x} = \phi_2(z, 0)$$

$$\frac{\partial v}{\partial x} = \frac{2 \cos 2z \cosh 0 - 2}{[\cosh 0 - \cos 2z]^2} = \frac{2[\cos 2z - 1]}{[1 - \cos 2z]^2} = \frac{-2[1 - \cos 2z]}{[1 - \cos 2z]^2}$$

$$\frac{\partial v}{\partial x} = \frac{-2}{2 \sin^2 z}$$

$$\text{Where } F(z) = (1+i)f(z)$$

$$u+v = V$$

$$\overline{\partial_x} = \phi_2(z, 0) = -\operatorname{cosec} 2z$$

$$\frac{\partial v}{\partial y} = \phi_1(z, 0) = \frac{[\cosh 2y - \cos 2x] 0 - \sin 2x [\sinh 2y(2)]}{[\cosh 2y - \cos 2x]^2}$$

$$\phi_1(z, 0) = \frac{\partial v}{\partial y} = \frac{-2 \sin 2x \sinh y}{[\cosh 2y - \cos 2x]^2}$$

$$\frac{\partial v}{\partial y} = \frac{-0 \sin 2z}{[\cosh 2y - \cos 2z]^2} = 0$$

$$f(z) = u + iv$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\frac{\partial v}{\partial y} + i \frac{\partial v}{\partial x}$$

$$f'(z) =$$

$$f(z) = \int [\phi_1(z,0) + i \phi_2(z,0)] dz + c$$

$$f(z) = \int -\operatorname{cosec}^2 z (i) dz + c$$

$$f(z) = -i(-\cot z) + c = i \cot z + c$$

$$f(z) = i \cot z + c$$

$$(1+i) f(z) = \frac{i}{1+i} + \frac{c}{1+i} = i \cot z + c$$

$$f(z) = \frac{i(1-i)}{2}$$

$$\cot z f(z) =$$

$$\cot z + c_1$$

$$i+1$$

$$f(z) = \frac{1}{2} \cot z + c_1$$

$$\frac{\partial u}{\partial x} = e^x x^2 \cos y + 2x e^x \cos y - e^x y$$

$$\phi_1(z,0) = \frac{\partial u}{\partial x} = e^z z^2$$

$$\phi_1(z,0) = \frac{\partial u}{\partial x} = e^z z^2 + 2z e^z$$

$$\frac{\partial u}{\partial y} = -e^x x^2 \sin y - 2xy e^x \sin y$$

$$\phi_2(z,0) = \frac{\partial u}{\partial y} = 0 + 0 - 0 - 0 = 0$$

4) Find the
 $e^x [(x^2 -$

Solution: $u = e^x x^2$

analytical function, whose real part is $u = y^2)(\cos y - 2xy \sin y)]$

$\cos y - e^x y^2 \cos y - 2xy e^x \sin y$

$x^2 \cos y - 2y e^x \sin y - 2xy e^x \sin y$

$$\cos(0) + 2z e^z \cos(0) - 0 - 0 - 0$$

$$\sin y + e^x \sin y - 2y e^x \cos y - 2x e^x \sin y - 2xy e^x \cos y$$

$$f(z) = u + iv$$

$$f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x} \\ = \frac{\partial u}{\partial x} - i \frac{\partial v}{\partial y}$$

$$f'(z) =$$

$$f(z) = \int_0^z (e^z z^2 + 2z e^z - 0) dz + c$$

$$f(z) = \int (e^z z^2 + 2z e^z - 0) dz + c$$

$$= \int e^z (z^2 + 2z) dz + c = \int e^z z^2$$

$$dz + 2 \int z e^z dz$$

$$u = z^2 \quad dv = e^z dz \quad du = 2z dz \quad v = e^z \quad f(z) = e^z$$

$$z^2 - 2 \int z dz e^z dz + 2 \int z e^z dz + c \quad \mathbf{f(z) = e^z}$$

$$\mathbf{z^2 + c}$$

- 5) The analytical function whose imaginary part is $v(x,y) = 2xy$ **Solution:**

$$\begin{aligned}
 v &= 2xy \\
 \frac{\partial v}{\partial x} &= 2y = \phi_2(z,0) = 2(0) = 0 \\
 \frac{\partial v}{\partial y} &= 2x = \phi_1(z,0) = 2(z) = 2z \quad f(z) \\
 &= \phi(z,0) \int dz = \int 2z dz + c \\
 &= z^2 + c \\
 f(z) &= z^2 + c \\
 \mathbf{f(z) = z^2 + c}
 \end{aligned}$$

- 6) Find harmonic conjugate at $u = e^{x^2-y^2} \cos 2xy$ and also find $f(z)$

Solution :

$$u = e^{x^2-y^2} \cos 2xy$$

$$\frac{\partial u}{\partial x} = e^{x^2-y^2} \cos 2xy (2x) - e^{x^2-y^2} \sin 2xy (2y)$$

$$\phi_1(z,0) = e^{z^2-0} \cos 0 (2z) - e^{x^2-y^2} (0)$$

$$\begin{aligned}
 \phi_1(z,0) &= e^{z^2} 2z \frac{\partial u}{\partial y} = e^{x^2-y^2} \cos 2xy (-2y) - \\
 &e^{x^2-y^2} \sin 2xy (2x)
 \end{aligned}$$

$$\phi_2(z,0) = 0 - 0$$

$$\phi_2(z,0) = 0 \quad f(z)$$

$$= u + iv \quad f'(z) =$$

$$\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} f'(z) =$$

$$f'(z) = \phi_1(z,0) - i \phi_2(z,0)$$

$$f(z) = \int [\phi_1(z,0) - i \phi_2(z,0)] dz + c \quad f(z) = \int e^{z^2} 2z$$

$$dz + c \quad (\text{put } z^2 = t \Rightarrow 2z dz = dt) \quad f(z) = \int e^t dt +$$

$$c = e^t + c$$

$$f(z) = e^{z^2} + c \quad f(z) = e^{(x+iy)^2} f(z) =$$

$$e^{x^2-y^2+2xyi} + c \quad f(z) = e^{x^2-y^2} e^{2xyi} + c \quad u+iv =$$

$$e^{x^2-y^2} [\cos 2xy + i \sin 2xy] + c \quad u+iv = e^{x^2-y^2}$$

$$\cos 2xy + i e^{x^2-y^2} (\sin 2xy) + c$$

$$v = e^{x^2-y^2} \sin 2xy + c$$

7) Find the analytical function $f(z)$ such that $\text{Re}[f'(z)] = 3x^2 - 4y - 3y^2$ and $f(1+i) = 0$.

Solution :

$$\operatorname{Re}[f'(z)] = 3x^2 - 4y - 3y^2$$

$$f(z) = u + iv$$

Integrate w.r.t 'y' we get

$$x^2y - \frac{4y^2}{2} - \frac{3y^3}{3} + f(x)$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$f'(z) =$$

$$\frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$\frac{\partial u}{\partial x}$$

$$\operatorname{Re}[f'(z)] =$$

$$\frac{\partial u}{\partial x} = 3x^2 - 4y - 3y^2$$

$$\frac{\partial v}{\partial y} = 3x^2 - 4y - 3y^2$$

Integrate w.r.t 'x' we get & $u = \frac{3x^3}{3} - 4xy - 3y^2x + f(y)$ $v = 3$

$$u = x^3 - 4xy - 3y^2x + f(y)$$

$$v = 3x^2y - y^3 - 2y^2 + f(x)$$

Differentiate w.r.t 'y' we get

$$\frac{\partial u}{\partial y} = -4x - 6xy + f'(y)$$

Differentiate w.r.t 'x' we get

$$\frac{\partial v}{\partial x} = 6xy + f'(x)$$

From C-R equations

$$\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$-4x - 6xy + f'(y) = -6xy - f'(x)$$

$$-4x + f'(y) = -f'(x)$$

Compare equation on both sides

$$\text{i.e } f'(x) = 4x, f'(y) = 0$$

$$f(x) = 4 \int x dx$$

$$f(y) = c$$

$$= \frac{4x^2}{2} + c$$

$$f(x) = 2x^2 + c \quad f(y) = c$$

$$f(z) = u+iv \quad f(z) = [x^3 - 4xy - 3y^2x] + i[3x^2y - y^3 - 2y^2] + 2x^2 + c$$

$$\text{given } f(1+i) = 0 \quad f(z) = u+iv$$

$$z = x+iy = (1+i)$$

$$\text{put } x = 1, y = 1 \quad f(z) = [1-4-3] + i[3-2-1]$$

$$+2 + c \quad f(1+i) = 0 = -6 + 2i + c$$

$$= 6 - 2i$$

$$f(z) = [x^3 - 4xy - 3y^2x] + i[3x^2y - y^3 - 2y^2] + 2x^2 + 6 - 2i$$

8) Find the analytic function $f(z) = u+iv$ if $u-v = e^x(\cos y - \sin y)$ **Solution:**

$$f(z) = u+iv \quad i f(z) = iu-v$$

$$(1+i) f(z) = (u-v) + i(u+v)$$

$$f(z) = u+iv \quad u = u-v = e^x$$

$$(\cos y - \sin y)$$

$$\begin{aligned}
 F(z) &= (1+i) f(z) \quad \cos y - e^x \sin y = \\
 \frac{\partial u}{\partial x} &= e^x \\
 \frac{\partial u}{\partial y} &= -e^x \\
 f'(z) &= \frac{\partial u}{\partial x} - i \frac{\partial v}{\partial y} \\
 f(z) &= \int [\phi_1(z,0) - i \phi_2(z,0)] dz + c \\
 f(z) &= \int (e^z + i e^z) dz + c \\
 \mathbf{f(z) = (e^z + i e^z) + c (1+i)} \\
 f(z) &= e^z + i e^z + c \\
 f(z) &= \frac{e^z(1+i)}{(1+i)} + \frac{c}{1+i} \mathbf{f(z)} \\
 &= \mathbf{e^z + c}
 \end{aligned}$$

Harmonic Conjugate

1) Show that function $u = 2xy + 3y$ is harmonic and find harmonic conjugate.

Solution:

$$u = 2xy + 3y$$

$$\frac{\partial u}{\partial x} = 2y$$

$$\frac{\partial^2 u}{\partial x^2} = 0$$

2

$$\frac{\partial u}{\partial y} = 2x + 3$$

$$\frac{\partial^2 u}{\partial y^2} = 0$$

2

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \quad u \text{ satisfies Laplace}$$

equation

'u' is a **Harmonic** function

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

$$dv = -(2x+3) dx + 2y dy$$

$$= \int -(2x+3) dx + \int 2y dy$$

$$v = -\left(\frac{2x^2}{2} + 3x\right) + \frac{2y^2}{2} + c$$

$$v = -x^2 + y^2 - 3x + c$$

2) Show that $u = 2\log(x^2 + y^2)$ is harmonic and find its harmonic conjugate.

Solution:

$$u = 2\log(x^2 + y^2)$$

$$\frac{\partial u}{\partial x} = 2 \frac{1}{x^2+y^2} 2x$$

$$\frac{\partial u}{\partial y} = 2 \frac{1}{x^2+y^2} 2y$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{(x^2+y^2)(4) - 4x(2x)}{(x^2+y^2)^2}$$

$$\frac{\partial^2 u}{\partial y^2} = \frac{(x^2+y^2)(4) - 4y(2y)}{(x^2+y^2)^2}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = \frac{4x^2+4y^2 - 8x^2+4x^2+4y^2 - 8y^2}{(x^2+y^2)^2}$$

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

$$dv = - \frac{\frac{\partial u}{\partial y}}{\frac{\partial u}{\partial x}} dx + \frac{\frac{\partial u}{\partial x}}{\frac{\partial u}{\partial y}} dy \quad \partial u$$

$$dv = \frac{-4y}{x^2+y^2} dx + \frac{4x}{x^2+y^2} dy$$

$$dv = \frac{-4}{x^2+y^2} (y dx - x dy)$$

$$v = - 4 \int \left[\frac{x dy - y dx}{x^2+y^2} \right]_v$$

$$= - 4 \int d \tan^{-1}\left(\frac{y}{x}\right)$$

$$v = - 4 \tan^{-1}\left(\frac{y}{x}\right) + c$$

$$d \tan^{-1}\left(\frac{y}{x}\right) = \frac{1}{1+(\frac{y}{x})^2} \left[\frac{x dy - y dx}{x^2} \right]_x$$

3) Find f(z) if the imaginary part is $r^2 \cos 2\theta + r \sin \theta$ **Solution:**

$$V = r^2 \cos 2\theta + r \sin \theta$$

Integrate w.r.t. θ we get $u = \frac{1}{2} \frac{x^2 + y^2}{r^2} + f(\theta)$

$$\frac{\partial u}{\partial \theta} = -r^2 \sin \theta + r \cos \theta + f'(\theta)$$

Differentiate (1) w.r.t. θ

$$\frac{\partial u}{\partial \theta} = -r^2 \sin \theta + r \cos \theta + f'(\theta) \rightarrow (3)$$

Compare the equations (2) & (3)

$$f'(\theta) = 0 \rightarrow (1)$$

$$f(\theta) = c$$

$$u = \frac{1}{2} \frac{x^2 + y^2}{r^2} + c$$

$$\rightarrow (2) \quad 4) \text{ Show that } \left[\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right] = -2 \frac{1}{r^2} \cos 2\theta - r \sin \theta$$

Solution:

$$f(z) = u + iv$$

$$\text{real } f(z) = u$$

$$[\text{real } f(z)]^2 = u^2$$

$$\frac{\partial(u^2)}{\partial x} = 2u \frac{\partial u}{\partial x}$$

$$\frac{\partial^2(u^2)}{\partial x^2} = 2u \frac{\partial^2 u}{\partial x^2} + 2 \frac{\partial u}{\partial x} \frac{\partial u}{\partial x} \rightarrow (1)$$

Similarly,

$$\rightarrow (2)$$

Add
equation

$$\left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right] u^2 = 2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] \quad (1) \quad \frac{\partial^2(u^2)}{\partial y^2} = 2u \frac{\partial^2 u}{\partial y^2} + 2 \frac{\partial u}{\partial y} \frac{\partial u}{\partial y} \quad \&$$

$$(2) \quad \left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right] u^2 = 2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] + 2u \left[\right]$$

$$\{ f(z) = u+iv \Rightarrow f'(z) = \frac{\partial u}{\partial x} - i \frac{\partial u}{\partial y} \}$$

$$|f'(z)|^2 = \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2$$

$$\left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right] u^2 = |f'(z)|^2$$

5) If $f(z)$ is analytical function with constant modulus ,then show that $f(z)$ is constant.

Solution:

let $f(z)$ is constant modulus

$$f(z) = u+iv$$

$$|f(z)| = \sqrt{u^2 + v^2} = \text{constant}$$

$$\sqrt{u^2 + v^2} = c$$

$$u^2 + v^2 = c^2 = c_1$$

Differentiate w.r.t 'x'

$$2u \frac{\partial u}{\partial x} + 2v \frac{\partial v}{\partial x} = 0 \rightarrow (1)$$

$$= 0 \rightarrow \textcircled{2} \quad 2u \frac{\partial u}{\partial y} + 2v \frac{\partial v}{\partial y} \quad \text{Differentiate w.r.t 'y'}$$

$$\text{equations} \quad 2u \frac{\partial v}{\partial y} - 2v \frac{\partial u}{\partial y} \quad \text{By C-R}$$

$$\textcircled{1} \ominus \quad 2u \frac{\partial u}{\partial y} + 2v \frac{\partial v}{\partial y} = 0 \rightarrow \textcircled{3}$$

$$= 0 \rightarrow \textcircled{4} \quad \frac{\partial v}{\partial y} - v^2 \frac{\partial u}{\partial y}$$

$$\text{Multiply } \textcircled{3} * v \ominus \quad uv \quad u^2 \frac{\partial u}{\partial y} + uv \frac{\partial v}{\partial y} = 0$$

$$\textcircled{4} * u \ominus \quad = 0$$

$$\text{Subtract} \quad \text{then} \quad uv$$

$$u = c$$

Similarly

$$u = \frac{\partial v}{\partial y} - v^2 \frac{\partial u}{\partial y} - u^2 \frac{\partial u}{\partial y} - uv \frac{\partial v}{\partial y} = 0$$

$$- \frac{\partial u}{\partial y} (u^2 + v^2) = 0$$

$$u^2 + v^2 \neq 0$$

$$\frac{\partial u}{\partial y} = 0$$

$$\int \frac{\partial u}{\partial y} = c$$

$$v = c f(z) \text{ is}$$

constant

Conformal Mapping :

A transformation $w = f(z)$ is said to be conformal if it preserves angle between oriented curves in magnitude as well as in orientation.

Bilinear Transformation :

The transformation $w = f(z) = \frac{az+b}{cz+d}$ is called the bilinear transformation or mobius transformation. Where a, b, c, d are complex constants.

The method to find the bilinear transformation if three points and their images are given as follows:

We know that we need four equations to find 4 unknowns. To find a bilinear transformation we need three points and their images.

in cross ratio, three are four points $(w, w_1, w_2, w_3) = (z, z_1, z_2, z_3)$

$$\frac{(w-w_1)(w_2-w_3)}{(w_1-w_2)(w_3-w)} = \frac{(z-z_1)(z_2-z_3)}{(z_1-z_2)(z_3-z)}$$

Since we have to get $w = \frac{-az+b}{cz+d}$, we take one point as 'z' and its image as 'w'

Problems about bilinear transformation:

1) Find the bilinear transformation on which maps the points (-1, 0, 1) into the points (0,i,3i) in w-plane

Solution :

In z-plane, $z_1 = -1, z_2 = 0, z_3 = 1$

In w-plane, $w_1 = 0, w_2 = i, w_3 = 3i$

In cross ratio,

$$(w, 0, i, 3i) = (z, -1, 0, 1)$$

$$\begin{aligned} \frac{(w-w_1)(w_2-w_3)}{(w_1-w_2)(w_3-w)} &= \frac{(z-z_1)(z_2-z_3)}{(z_1-z_2)(z_3-z)} \\ \frac{(w-0)(i-3i)}{(0-i)(3i-w)} &= \frac{(z+1)(0-1)}{(-1-0)(1-z)} \\ \frac{(w)(-2i)}{(-i)(3i-w)} &= \frac{-(z+1)}{-(1-z)} \end{aligned}$$

$$-2wi(1-z) = (z+1) [-i(3i-w)]$$

$$-2wi + 2wiz = -[-3-wi](z+1)$$

$$-2wi + 2wiz = 3z + wiz + 3 + wi$$

$$\frac{(w-0)(0-1)}{(0-i)(1-0)} = \frac{(0-1)(i-0)}{(1-0)(0-z)}$$

$$\frac{-w}{-i} = \frac{-i}{-z} \quad w = \frac{i^2}{z} = \frac{-1}{z} \quad \mathbf{w} = \frac{-1}{z}$$

=

$$-3wi + wiz = (3z + 3)$$

$$= \quad z_1 = \alpha = \frac{1}{z_1} = \frac{1}{0} \quad [z_1' = 0], \quad z_2 = i, \quad z_3 = 0$$

$$w[i(3-z)] = z(z+1) \quad \mathbf{w} = \frac{-3(z+1)}{-i(3-z)}$$

2) Find the bilinear transformation which maps the points $(\alpha, i, 0)$ in the z-plane into $(0, i, \alpha)$ in the w-plane.

$$w_1 = 0, \quad w_2 = i, \quad w_3 = \frac{1}{0} = \alpha = \frac{1}{w_3} \quad [w_3' = 0]$$

maps the points $(\alpha, i, 0)$ in

Solution: In z-plane, z

$$\frac{(w-w_1)(w_2-w_3)}{(w-w_3)(w_2-w_1)} = \frac{(z-z_1)(z_2-z_3)}{(z-z_3)(z_2-z_1)}$$

In w-plane,

$$(w-w_1)$$

$$(w-w_3)(w_2-w_1) = (z-z_3)(z_2-z_1)$$

$$z_1' z_3 - ((w-w_1-w_2)(w-w_3)(-z_1 z_2)(z_3-z) w_3' z_1')$$

3) Find the bilinear transformation that maps the points $(0, i, \alpha)$ respectively into $(0, 1, \alpha)$.

Solution:

$$\text{In } z\text{-plane, } z_1 = 0, z_2 = i, z_3 = \frac{1}{0} = \frac{1}{\alpha} = \frac{1}{z_1} [z_3' = 0]$$

$$w_1 = 0, w_2 = 1, w_3 = \frac{1}{w_1} = \frac{1}{0} = \alpha [w_3' = 0]$$

In w-plane,

$$\frac{(w-w_1)(w_2-w_3)}{(w_1-w_2)(w_3-w_1)} = \frac{(z-z_1)(z_2-z_3)}{(z_1-z_2)(z_3-z_1)}$$

$$\frac{(w-0)(0-1)}{(0-1)(1-0)} = \frac{(z-0)(i(0)-0)}{(0-i)(1-0)}$$

$$\frac{-w}{-1} = \frac{-z}{-i}$$

$$w = -iz$$

Fixed point :

$$\text{The transformation } w = \frac{az+b}{cz+d}$$

The roots of this transformation are called fixed points or invariant points.

$$z = \frac{az+b}{cz+d} \text{ (we know that } w = f(z) \text{) } z(cz+d) =$$

$$az+b \quad cz^2+dz = az+b \quad cz^2+(d-a)z - b = 0$$

Problems:

1) Find the fixed points of the transformation $w =$

Solution: The roots of above transformation are called fixed points

i fixed points $\pm i$

2) The fixed points of the transformation $w =$

Solution:

$$\begin{aligned} \text{put } w &= z \\ z &= \frac{z-1+i}{z+2} \\ z(z+2) &= (z-1+i) \quad (a=1, b=1, c=1-i) \\ z^2+2z &= z-1+i \\ z^2+z+i-1 &= 0 \\ z &= \frac{-b \pm \sqrt{b^2-4ac}}{2a} = \frac{-1 \pm \sqrt{1+4(1-i)}}{2} \\ -1 \pm \frac{1+\sqrt{4-4i}}{2} &= \frac{-1 \pm \sqrt{3-4i}}{2} \\ \frac{-1 + \sqrt{3-4i}}{2} & \quad \frac{-1 - \sqrt{3-4i}}{2} \end{aligned}$$

3) Determine the bilinear transformation whose fixed points are 1,-1 Solution:

Given fixed points are $z = 1, -1$

The roots of the transformation is $w = \frac{az+b}{cz+d}$ are called fixed points **put $w = z$**

$$z = \frac{az+b}{cz+d}$$

$$cz^2 + (d-a)z - b = 0 \quad (z+1)(z-1) = 0$$

$$z^2 - 1 = 0 \quad (c=1, d=0, a=0, b=1)$$

$$w = \frac{0z+1}{1z+0} = \frac{1}{z}$$

Problems on images:

1) Write the image of the triangle with vertices $(i, 1+i, 1)$ in the z -plane under the transformation $w = 3z+4-2i$

Solution:

y

$$(x, y) = (1, 0)$$

In w -plane:

in z-plane Transformation $z = i$ \odot y
 $x+iy = 0+i$ $w = 3z+4-2i$ $(x,y) = (0,1)$ $w =$
 $3(x+iy)+4-2i$ $z = 1+i$ \odot $x+iy = 1+i$ $u+iv = w$

$(x,y) = (1,1)$ $u = 3x+4, v = 3y-2$

x

z- plane


$(1,0)$
 $z = 1$ \odot $x+iy = 1$

$$\text{i) } (x,y) = (0,1) \quad (u,v) = (4,1) \quad \text{ii) } (x,y) =$$

$$(1,1) \quad (u,v) = (7,1) \quad \text{iii) } (x,y) = (1,0) \quad (u,v) = (7,-2)$$

Conclusion:

The image of the triangle whose vertices $(i,1+i,1)$ is mapped as triangle whose vertices $(4,1), (7,1), (7,-2)$ in w -plane under the transformation $w=3z+4-2i$



2) Find the image of the infinite strip $0 < y < \frac{1}{2}$ under the transformation $w = \frac{1}{z}$


Solution:

In z -plane

the infinite strip between the lines $y=0, y = \frac{1}{2}$.

Transformation:

$$w = \frac{1}{z} \quad \frac{1}{z} = \frac{u-iv}{u^2+v^2} \quad x+iy = \frac{u}{u^2+v^2} + i \frac{-v}{u^2+v^2}$$



In w-plane

$$i) y = 0 \Rightarrow 0 = \frac{-v}{u^2+v^2}$$

z-plane

$$ii) y = \frac{1}{2} \Rightarrow \frac{1}{2} = \frac{-v}{u^2+v^2}$$

$$0 = -v \quad u^2 + v^2 = -2v \quad v = 0 \quad \text{Conclusion: } 1$$

The image of infinite strip $0 < y < \frac{1}{2}$ is transferred as straight line ($v=0$) or circle under the transformation $w = \frac{z-i}{z+i}$

3) Find the image of the region in the z-plane between the lines $y = 0$ and $y = \frac{\pi}{2}$ under the transformation $w = e^z$

Solution:

In z-plane

The lines are $y = 0, y = \frac{\pi}{2}$

Transformation

$$w = e^z$$

$$u+iv = e^{x+iy} = e^x e^{iy} \quad y = 0 \quad u+iv = e^x$$


$$[\cos y + i \sin y] \quad u = e^x \cos y \quad v = e^x \sin y$$

In w-plane

$$i) y = 0 \quad u = e^x, \quad v = 0$$

$$ii) y = \frac{\pi}{2} \quad u = 0, \quad v = e^x$$

$$v = 0$$



Conclusion:

The image of the region lines $y = 0$ & $y = \frac{\pi}{2}$ are transferred as first quadrant in the w -plane under the transformation $w = e^z$

- 4) Show that transformation $w = z + \frac{1}{z}$ maps the circle $|z| = c$ into the ellipse $u = (c + \frac{1}{c}) \cos \theta$, $v = (c - \frac{1}{c}) \sin \theta$. Also discuss the case when $c = 1$ in detail.

Solution:

Z-plane

Transformation

1

The circle $|z| = c$

$|z|$

w

$$\sqrt{\quad} = z$$

+

z

$$x + iy = c$$

$$w = r e^{i\theta} + \frac{1}{r e^{i\theta}}$$

$$x^2 + y^2 = c^2 \quad u + iv = r(\cos \theta + i \sin \theta) + \frac{1}{r}(\cos \theta - i \sin \theta) \quad + y^2 = c^2 \quad u + iv =$$

$$\left(r + \frac{1}{r}\right) \cos \theta + i \left(r - \frac{1}{r}\right) \sin \theta \quad u = \left(r + \frac{1}{r}\right) \cos \theta \quad v = \left(r - \frac{1}{r}\right) \sin \theta$$

w-plane

$$|z| = c$$

y

$$|z| = r \quad (r = c)$$

we know that

$$\sin^2 \theta = 1 \quad \frac{u^2}{\left(c + \frac{1}{c}\right)^2} + \frac{v^2}{\left(c - \frac{1}{c}\right)^2} = 1$$


Case:

When $c = 1$

$$|z| = 1, \quad \frac{u^2}{a^2} + \frac{v^2}{b^2} = 1$$

$$r = 1$$

$$u = 2 \cos \theta, \quad v = 0$$




$$\cos^2 \theta +$$

$$u+iv = 2 \cos \theta + i (0) 2 \sin \theta \quad u = 2 \quad v = 0$$

Conclusion:

The image of circle $z = c$ is transferred as ellipse $\frac{u^2}{a^2} + \frac{v^2}{b^2} = 1$ in w -plane and also the image of circle $z = 1$ when $c = 1$ is transferred as straight lines $u = 2$ & $v = 0$ in w -plane under the transformation $w = z + \frac{1}{z}$.



5) Discuss the transformation of $w = \sin z$ using example.

$+ b$


Solution:

Transformation $w = \sin z$

$$w = \sin(x+iy) \quad w = \sin x \cosh y + i \cos x \sinh y$$

$$u+iv = \sin x \cosh y + i \cos x \sinh y$$

$$u = \sin x \cosh y \quad v = \cos x \sinh y$$



Example: In z -plane

$x = c$

$$\cosh y = \frac{u}{\sin c}, \quad \sinh y = \frac{v}{\cos c}$$

$$x^2 + y^2 = 1$$

In w -plane

$$u^2 + v^2 = 1$$

$$\sin x$$

$$\cos x$$

Conclusion:


put $x = c$

$$\frac{u^2}{\sin^2 c} - \frac{v^2}{\cos^2 c} = 1$$

The image line x

$= c$ is transferred as hyperbola

$\frac{u^2}{a^2} - \frac{v^2}{b^2} = 1$ in w -plane under the transformation $w = \sin z$.




6) Discuss the transformation of $w = \cos z$

Solution: Transformation on $w = \cos z$

$$w = \cos(x+iy) \quad w = \cos x \cosh y - \sin x \sinh y \quad u+iv =$$

$$\cos x \cosh y - i \sin x \sinh y \quad u = \cos x \cosh y$$

$$v = \frac{u}{\cosh y},$$



$\sin x \sinh y$ In

z- $\frac{v}{\sinh y}$

plane In w -plane $y = c \quad \cos x = \sin x = -$

$$\cos^2 x + \sin^2 x = 1$$


$$\frac{u^2}{\cosh^2 y} + \frac{v^2}{\sinh^2 y} = 1$$

put $y = c$

y

$$\frac{u^2}{\cosh^2 c} + \frac{v^2}{\sinh^2 c} = 1$$

$$\frac{u^2}{a^2} + \frac{v^2}{b^2} = 1$$



Conclusion:

The image of line $y = c$ is transferred as ellipse $\frac{u^2}{a^2} + \frac{v^2}{b^2} = 1$ under the transformation $w = \cos z$.

Unit – 2

Complex Integration


Line Integral:

suppose $f(z)$ is a complex function in the region R , and C is a smooth curve in R . Consider an interval

$x_1 < x_2 \dots < x_n < b$ are points in (a, b) .
 (a, b) and $a <$

$\Delta x_r = x_r - x_{r-1}$ are chord vectors, then

$$\lim_{n \rightarrow \infty} \sum_{r=1}^n \Delta x_r = \int_a^b f(z) dz$$



Where the summation tends to a limit and independent of the points choice. The limit exists if $f(z)$ is continuous along the path.

Evaluation of the integrals: $f(z) dz = (u + iv)(dx + idy) = (udx - vdy + i(udy + vdx))$ where u and v are functions of x .

()

Problems:

1) Evaluate $\int_C x^2 + ixy dz$ from A(1, 1) to B(2, 8) along $x = t$ and $y = t^3$.

Solution: Along $x = t, y = t^3$, $dx = dt, dy = 3t^2 dt$, The limits for t are 1 and 2

$$\int_C x^2 + ixy (dx + idy) = \int_1^2 (x^2 dx - xy dy) + i \int_1^2 (xy dx + x^2 dy)$$

$$= \int_1^2 (t^2 dt - 3t^6 dt) + i \int_1^2 (t^4 dt + t^5 dt)$$

and upper limit)

$$= -\frac{1094}{2} + \frac{124i}{5}$$

$\int_0^{1+i} z^2 dz$ along $y = x^2$

2) Evaluate $\int_0^z z^2 dz$

$\int_0^{1+i} z^2 dz$ along $y = x^2$, $dy = 2x dx$

Solution: $\int_0^z z^2 dz$

$$\int_0^{1+i} (x^2 - y^2 + 2ixy)(dx + idy)$$

$$= \int_0^1 (x^2 - x^4) dx - 2 \int_0^1 x^3 \cdot 2x dx + i \int_0^1 (x^2 - x^4) 2x dx + 2i \int_0^1 x^3 dx$$

$$= \left[\frac{x^3}{3} - \frac{x^5}{5} \right]_0^1 - 4 \left[\frac{x^5}{5} \right]_0^1 + i \left[\frac{x^3}{3} - \frac{x^5}{5} \right]_0^1 + 2i \left[\frac{x^4}{4} \right]_0^1$$

$$= \frac{0}{2} \left(\frac{x}{2} \right)$$

$$= -\frac{1}{3} + \frac{i}{3}$$

$$2+i$$

3) Evaluate $\int_{1-i}^{2+i} (2x + 1 + iy) dz$ along $(1-i)$ to $(2+i)$.

Solution: Along $(1-i)$ to $(2+i)$ is the straight line AB joining $(1,-1)$ to $(2,1)$.

$$\text{The equation of AB is } y-1 = -\frac{(-1-1)}{(1-2)}(x-2) \quad y-2x =$$

$$-3, y = 2x-3, dy = 2dx$$

X varies from 1 to 2

$$2+i2$$

$$\int_{1-i}^{2+i} (2x+1+iy) dz = \int_1^2 [2x+1 dx - (2x-3)2dx + i[2x-3]dx + (2x+1)2dx]$$

$$= \int_1^2 (-2x+7) dx + i \int_1^2 (6x-1) dx$$

$$= -\frac{2x^2}{2} + 7x + i \left(\frac{6x^2}{2} - x \right) \Big|_1^2 \text{ (apply the lower$$


and upper limit)

$$2+i$$

$$\int_{1-i}^{2+i} (2x+1+iy) dz = 4+8i$$

$$\int_{(1,1)}^{(0,0)} [2+5y+i(x^2-y^2)] dz \text{ along } y^2 = x.$$

4) Evaluate $\int_{(0,0)}^{(1,1)} [3x$



Solution: Along $y^2 = x$, $2ydy = dx$, y varies from 0 to 1.

$$\int_{(0,0)}^{(1,1)} [3x^2 + 5y + i(x^2 - y^2)] [dx + i dy] = \int_0^1 3y^4 2y dy + 5y 2y - (y^4 - y^2) dy + i[(3y^4 + 5y) dy + (y^4 - y^2) 2y dy]$$

$$= 5 \frac{y^6}{6} - \frac{y^3}{3} + 11 \frac{y^5}{5} + i(2 + 3 \frac{y^4}{4} - 2 + 5 \frac{y^2}{2}) \text{ (apply the lower}$$

and upper limit)

$$= \frac{129}{30} + \frac{44i}{15}$$

$\int_{(0,0)}^{(1,1)} (x^2 y dx + (x^2 - y^2) dy)$ along a) $y = 3x^2$ b) $y = 3x$.

5) Evaluate $\int_{(0,0)}^X$

Solution: a) $y = 3x^2$, $dy = 6x dx$, x varies from 0 to 1.

$$\int_{(0,0)}^{(1,1)} x^2 y dx + (x^2 - y^2) dy = \int_0^1 3x^4 dx + (x^2 - 9x^4) 6x dx$$

(1,3)

5

4

$$\int_{(0,0)}^{(1,3)} x^2 y dx + (x^2 - y^2) dy = 3 \underline{\hspace{1cm}}$$

$$x^5 + 6x^4 - 54x^6$$

69

= - $\underline{\hspace{1cm}}$

10

(1,3)

b) $y = 3x$, $dy = 3dx$, x varies from 0 to 1.

$$\int_{(0,0)}^{(1,3)} x^2 y dx + (x^2 - y^2) dy =$$

$$\int_0^1 3x^3 dx + (x^2 - 9x^2) 3dx$$

x^4

x^3

= 3

$\underline{\hspace{1cm}}$ - 24

$\underline{\hspace{1cm}}$ (apply the lower

4

3

and upper limit)

29

= -

$\underline{\hspace{1cm}}$

4

6) Evaluate $\oint_C (3z + 1) dz$ where C is the boundary of the square with vertices at the points $z = 0$, $z = 1$, $z = 1+i$, $z = i$ and the orientation of C is anti-clockwise. **Solution: C is the square OABC**

$$+ c^2(3z + 1)dz + c^3($$

Along $C_1 = OA$
()

$$dy=0 \quad C(0,1)$$


$$A(1,0)$$

Along $C_2 = AB$

X varies from 0 to 1
 $c^1 3z+1 dz = \int_0^1 (3x+1)dx = 3 \frac{x^2}{2} + x$ (apply the lower and upper limit)

$$\frac{5}{2} =$$

y



$$Z=0 \quad 0 \quad Z=1$$

$$x = 1, dx = 0 \quad y$$

varies from 0 to 1

$$\int_0^1 \int_1^3 c_2(3z + 1) dz = i \int_0^1 [3(1+iy)+1] dy = 4i - 2$$

Along $c_3 = BC$ $y = 1, dy = 0 \quad x$

varies from 1 to 0

$$\int_1^0 \int_0^3 c_3(3z + 1) dz = - \int_1^0 [3(x + i) + 1] dx = -\frac{1}{2} - 3i - 1$$

Along $c_4 = CO$ $x = 0, dx = 0 \quad y$

varies from 1 to 0

$$\int_1^0 \int_0^3 c_4(3z + 1) dz = - \int_1^0 [3iy + 1] i dy = -\frac{1}{2} - i$$

$$\int_0^1 \int_0^3 c_4(3z + 1) dz = \frac{1}{2} + 4i - \frac{1}{2} - 2 = 4i - 2$$

$$-3i - i + \frac{1}{2} = 0$$

$$\oint c(3z+1) dz = 0$$

Evaluate $\int_{(0,0)}^{(1,1)} [3x^2 + 4xy + ix^2] dz$ along $y = x^2$ 7)
Solution: $y = x^2, dy = 2x dx,$

$$\int_{(0,0)}^{(1,1)} [3x^2 + 4xy + ix^2] dz = \int_0^1 (3x^2 + 4x^3 + ix^2)(dx + i2x dx)$$

$$= \int_0^1 (3x^2 + 4x^3 - 2x^3) dx + i \int_0^1 (6x^3 + 8x^4 + x^2) dx$$

$$= \left[3x^3 + \frac{4}{2}x^4 - \frac{2}{2}x^4 \right]_0^1 + i \left[\frac{6}{4}x^4 + \frac{8}{5}x^5 + \frac{1}{3}x^3 \right]_0^1$$

$$= \left[\frac{3}{2}x^4 + \frac{2}{5}x^5 + \frac{1}{3}x^3 \right]_0^1 \quad (\text{apply the lower and upper limit})$$

$$= \frac{3}{2} + \frac{2}{5} + \frac{1}{3}$$

8) Evaluate $\int_C (y^2 + 2xy) dx + (x^2 - 2xy) dy$, where is the boundary of the region by $y = x^2$ and $x = y^2$

Solution:

C_1 : Along OA, $y = x^2, dy = 2x dx$ X varies from 0 to 1 $\int_{C_1} (y^2 + 2xy) dx + (x^2 - 2xy) dy = \int_0^1 (x^4 + 2x^3) dx + (x^2 - 2x^3) 2x dx$

$$= \int_0^1 (x^4 + 2x^3 + 2x^3 - 4x^4) dx$$

$$= \int_0^1 (-2x^4 + 4x^3) dx$$

$$= \left[-\frac{2}{5}x^5 + x^4 \right]_0^1$$

$$= -\frac{2}{5} + 1 = \frac{3}{5}$$

C_2 : Along ABO, $x = y^2, dx = 2y dy$ y varies from 1 to 0 -

$$\int_{C_2} (y^2 + 2xy) dx + (x^2 - 2xy) dy = \int_1^0 (y^4 + 2y^3) 2y dy + (y^4 - 2y^3) dy$$

$$\int_C (y^2 + 2xy)dx + (x^2 - 2xy)dy = -1$$

$$\int_C (y^2 + 2xy)dx + (x^2 - 2xy)dy = -1$$

Cauchy's theorem

If $f(z)$ is analytical and $f'(z)$ is continuous inside and on a closed curve C , then $\int_C f(z)dz = 0$.

Proof: Suppose R is the region bounded by C

$$f(z) = u + iv \quad z = x + iy$$

$$x + iy$$

Where C

$$\int_C f(z)dz = \int_C (u + iv)(dx + idy) = \int_C (u dx - v dy) + i \int_C (v dx + u dy)$$


$$+ v dx)$$

$$\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y}$$


Since $f'(z)$ is continuous, $\frac{\partial u}{\partial x}$, $\frac{\partial u}{\partial y}$, $\frac{\partial v}{\partial x}$, $\frac{\partial v}{\partial y}$ exist and are continuous in R .

According to Green's theorem

$$\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y}$$



simple



$$\oint_C u dx + v dy = \iint_R (\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y}) dx dy$$

$$(\quad) \quad \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \quad \frac{\partial V}{\partial x} - \frac{\partial U}{\partial y}$$

$$\oint_C f(z) dz = \iint_R (-\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y}) dx dy + i \iint_R (\frac{\partial v}{\partial y} - \frac{\partial u}{\partial x}) dx dy$$

$$(\quad) \quad \frac{\partial U}{\partial x} - \frac{\partial u}{\partial y} \quad \frac{\partial V}{\partial x}$$

Since f(z) is analytic

$$\oint_C f(z) dz = \iint_R (\frac{\partial v}{\partial y} - \frac{\partial u}{\partial x}) dx dy + i \iint_R (\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y}) dx dy$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$\frac{\partial u}{\partial x} = 0$$

$$\oint_C f(z) dz = 0$$

Cauchy's Integral Formula

If f(z) is analytical within and on a simple closed curve and a is any point inside C, then

$$\frac{1}{2\pi i} \oint_C \frac{f(z) dz}{z-a}$$

$$f(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{z-a}$$

proof: C is a closed curve and a is any point inside C, Enclose a within a circle C whose radius is r and the centre is at a. Now C is inside C.

f(z) is not analytical

inside C.

$$(z-a)$$

By Cauchy's theorem for multiple connected region $\oint_C g(z) dz = \oint_{C'} g(z) dz$

$$\oint_C f(z) dz = \oint_{C'} f(z) dz$$

$$\oint_C g(z) dz = \oint_{C'} g(z) dz$$

Where

$$C' \text{ is } |z-a| = r$$

$$|z-a| = r$$

$$z-a = re^{i\theta}, z = a + re^{i\theta}$$

$$dz = rie^{i\theta} d\theta$$

θ varies from 0 to 2π in C'

$$\oint_C f(z-a) dz = \oint_{C'} f(z-a) dz = \int_0^{2\pi} f(a + re^{i\theta}) rie^{i\theta} d\theta = i \int_0^{2\pi} f(a + re^{i\theta}) d\theta$$

As $r \rightarrow 0, C' \rightarrow 0$


$$\oint_C f(z) dz = \oint_{C'} f(z) dz$$

$$\oint_C f(z-a) dz = i \int_0^{2\pi} f(a) d\theta = f(a) 2\pi i$$

$$f(a) = \frac{1}{2\pi i} \oint_C f(z-a) dz$$

Cauchy's integral formula for the derivatives

$$\frac{f^{(n)}(a)}{n!} = \frac{1}{2\pi i} \oint_C f(z) dz$$



$$f(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)}$$

Differentiating with respect to a successively

$$f'(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^2}$$

$$f''(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^3}$$

$$f'''(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^4}$$

$$f^{(iv)}(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^5}$$

.

.

.

$$f^{(n)}(a) = \frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^{n+1}}$$

We can evaluate easily the integrals of complex functions using this formula.

Problems:

$$\oint_C z e^z dz$$

1) Evaluate $\oint_C \frac{1}{(z+2)^3} dz$ where C is $|z| = 3$. **Solution:**

$z = -2$ lies inside $|z| = 3$

According to Cauchy's integral formula

$$\frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^{n+1}} = \frac{f^{(n)}(a)}{n!}$$

$$f(z) = z e^z$$

$$n=2$$

$$f'(z) = z e^z + e^z$$

$$f''(z) = z e^z + 2e^z$$

$$f''(-2) = -2e^{-2} + 2e^{-2} = 0$$

$$\oint_C \frac{z e^z dz}{(z+2)^3} = 0$$

$$\oint_C \frac{1}{(z+2)^3} dz = 0$$

$$dz$$

2) Evaluate $\oint_C \frac{1}{z^3(z+4)} dz$ where C is $|z| = 2$ using Cauchy's integral formula.

Solution: $z = 0$ lies inside C and $z = -4$ lies outside.

According to Cauchy's integral formula

1

$$f''(a) = \frac{2!}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^{n+1}} \quad [a=0] \quad \text{and } f(z) = \frac{1}{(z+4)}$$

$$f'(z) = -\frac{1}{(z+4)^2} \quad f''(z) = \frac{2}{(z+4)^3} \quad \text{and } f''(0) = \frac{2}{4^3} = \frac{1}{32}$$

$$\oint_C \frac{1}{z^3(z+4)} dz = \frac{1}{32}$$

3) Evaluate $\oint_C \frac{(z^3 - \sin 3z) dz}{(z - \frac{\pi}{2})^3}$ where C is $|z| = 2$ using Cauchy's integral formula.

Solution: According to Cauchy's integral formula

$$\frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^3} = \frac{f''(a)}{2!} \quad [a = \frac{\pi}{2} \text{ and } f(z) = z^3 - \sin 3z]$$

$\frac{\pi}{2} < 2$, $z = \frac{\pi}{2}$ lies inside C: $|z| = 2$

$$f'(z) = 3z^2 - 3\cos 3z$$

$$f''(z) = 6z - 9\sin 3z$$

$$f''(\frac{\pi}{2}) = 3\pi - 9$$

$$\oint_C \frac{(z^3 - \sin 3z) dz}{(z - \frac{\pi}{2})^3} = \pi i (3\pi - 9)$$

4) Evaluate $\oint_C e^{-z} dz$ where C is $|z| = 2$ using Cauchy's integral formula.

$$\oint_C e^{-z} dz$$

Solution: $\oint_C e^{-z} dz = \oint_C \frac{1}{(z-1)^3} dz$

$z = 1$ lies inside C i.e. $|z| = 2$

$$f(z) = e^{-z}$$

According to Cauchy's integral formula

$$\frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^3} = \frac{f''(a)}{2!}$$

$f(a)$, $[a = 1]$

$$\frac{1}{2\pi i} \oint_C f(z) dz$$

$$f^{(l)}(a) = \frac{l!}{(l-1)!} \frac{1}{c} \oint_C \frac{f(z) dz}{(z-a)^3}$$

$$2\pi i$$

$$f'(z) = -e^{-z}, f'(z) = e^{-z}, f'(1) = e^{-1}$$

$$\frac{\oint_C e^{-z} dz}{c(z-1)^3} = \frac{i\pi}{e}$$

5) Using Cauchy's integral formula evaluate $\oint_C \frac{z^4 dz}{(z+1)(z-i)^2}$ where C is ellipse and $9x^2 + 4y^2 = c$

36.

Solution:

$$\oint_C \frac{z^4 dz}{(z+1)(z-i)^2}$$

$$= \frac{1}{2\pi i} \oint_C \frac{z^4 dz}{(z+1)(z-i)^2} = \frac{1}{2\pi i} \oint_C \frac{z^4 dz}{(z+1)(z-i)^2} = \frac{1}{2\pi i} \oint_C \frac{z^4 dz}{(z+1)(z-i)^2}$$

Splitting into partial fractions $z = -1$ and $z = i$ lie inside $9x^2 + 4y^2 = 36$

$$\frac{1}{2\pi i} \oint_C \frac{f(z) dz}{(z-a)^2}$$

$$f(a) =$$

$$\frac{f'(z)}{(z-a)^2} = f'(a)$$

$$2\pi i$$

$$f(z) = z^4, a = -1, f(-1) = 1, a = i, f(i) = 1$$


$$f'(z) = 4z^3 \text{ and } f'(i) = -4i$$

$$\frac{1}{(1+i)^2} = 1, \frac{1}{(1+i)}$$

$$\begin{aligned} \oint_C \frac{z^4 dz}{(z+1)(z-i)^2} &= \frac{z^4}{(1+i)^2} \left(2\pi i - 2\pi i + 2\pi i (-4i) \right) \\ &= 4\pi(1-i) \frac{8\pi}{(1+i)} \end{aligned}$$

log z dz

6) Evaluate $\oint_C \frac{1}{(z-1)^3} dz$ where C is $|z-1|=2$ using Cauchy's integral formula
Solution:



According to Cauchy's integral formula

$$\oint_C f(z) dz = \frac{f^{(n)}(a)}{n!} \cdot \text{Area of } C$$

$$\oint_C \frac{1}{(z-a)^3} dz = 2\pi i$$

$$a=1$$

$|z-1|=2$ is a circle whose centre is (1,0)

radius is , a=1 lies inside C

1 and

$$\frac{1}{z^2} \quad \frac{f(z)}{z^2} dz$$

$$f(z) =$$

$$\log z, f'(z) = \frac{1}{z}, f''(z) = -\frac{1}{z^2}, f''(1) = -1$$

$$\frac{z}{1}$$

$$f''(a) = \pi i \oint_C \frac{f(z)}{(z-a)^3} dz$$

$$= -\pi i \oint_C \frac{1}{(z-1)^3} dz$$

$$\oint_C \frac{(z^2 - z - 1) dz}{z(z-i)^2}$$

7) Evaluate $\oint_C \frac{1}{z(z-i)^2}$ where C is $|z-2|=1$
Solution:

According to Cauchy's integral formula

$$\oint_C \frac{f(z)}{(z-a)^2} dz = 2\pi i f'(a)$$


z=0 inside C and z=i is outside C

$$f(z) = \frac{1}{z^2}, [a=0, f'(0) = -1]$$

2

$$\oint_C \frac{1}{z(z-i)^2} dz = 2\pi i$$

$$\oint_C (3z^2 + 7z + 1) dz$$



9) If $F(a) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-a)} dz$ using Cauchy's integral formula where C is $|z|=2$, $F(1)$, $F(3)$, $F'(1-i)$.

$$(3z^2+7z+1)dz$$

Solution: Suppose $F(a) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-a)} dz$

$$F(1) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-1)} dz, \text{ [z=1 lies inside C]}$$

$$f(z)dz$$

$$\oint_C \frac{f(z)}{(z-a)} dz = 2\pi i f(a)$$

$$[f(z) = 3z^2 + 7z + 1, f(1) = 3 + 7 + 1 = 11]$$

$$(3z^2 + 7z + 1)$$

$$2$$

$$C$$

$$=$$

$$2\pi i \cdot 11 = 22\pi i = F(1)$$

$$(z-3)$$

$$2$$

$$(3z^2 + 7z + 1)$$

$$F(z) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-3)} dz,$$

$$[z=3 \text{ is outside } C]$$

$$(z-3)$$

$$(3z^2 + 7z + 1)$$

$$2$$

$$C$$

$$\oint_C \frac{f(z)}{(z-3)} dz = 0 = F(3)$$

$$(z-3)$$

$$a = 1-i \text{ is inside } C$$

$$F(a) = 2\pi i (3a^2 + 7a + 1)$$

$$F'(a) = 2\pi i (6a + 7)$$

$$F''(a) = 12\pi i$$

$$F''(1-i) = 12\pi i$$

Complex Power Series

Taylor's Theorem:

If $f(z)$ is analytic inside and on a simple closed circle C with centre at a , then for z inside C

$$f(z) = f(a) +$$

$$f'(a)(z-a) + \frac{f''(a)}{2!}(z-a)^2 + \frac{f'''(a)}{3!}(z-a)^3 + \dots$$

2!

3!

Proof: Let z be any point inside C , then enclose z with a circle c' , with centre at a , let w be a point on c' , then

converges $= \frac{1}{w-a} \left(1 - \frac{z-a}{w-a} \right) = \frac{1}{w-a} \left(1 - \frac{z-a}{w-a} + \frac{(z-a)^2}{(w-a)^2} - \frac{(z-a)^3}{(w-a)^3} + \dots \right)$

uniformly $= \frac{1}{w-a} \left[1 + \frac{z-a}{w-a} + \frac{(z-a)^2}{(w-a)^2} + \frac{(z-a)^3}{(w-a)^3} + \dots + \frac{(z-a)^n}{(w-a)^n} + \dots \right]$


$$|z-a| < |w-a|$$

multiplying $\left| \frac{z-a}{w-a} \right| < 1$

both sides by $f(w)$ and integrating with respect to w on c' $\int_{c'} f(w) \frac{1}{w-a} dw = \int_{c'} f(w) \left(1 - \frac{z-a}{w-a} + \frac{(z-a)^2}{(w-a)^2} - \frac{(z-a)^3}{(w-a)^3} + \dots \right) dw$

$$= \int_{c'} f(w) dw - (z-a) \int_{c'} \frac{f(w)}{w-a} dw + \frac{(z-a)^2}{2!} \int_{c'} \frac{f(w)}{(w-a)^2} dw - \frac{(z-a)^3}{3!} \int_{c'} \frac{f(w)}{(w-a)^3} dw + \dots$$

$f(w)$ is analytic on c'



Therefore, this series

$$f(z) = \frac{1}{2\pi i} \oint_C \frac{f(w)dw}{w-z}$$

and $n! = 2\pi i c^l (w-a)_{n+1}$ Dividing by $2\pi i$

$$\frac{1}{(w-a)^{n+1}} = \frac{1}{(z-a)^{n+1}} + 2\pi i \frac{(z-a)^{-n}}{(z-a)^2} + 2\pi i \frac{(z-a)^{-n-1}}{(z-a)} + \dots + 2\pi i \frac{(z-a)^{-n-n}}{(z-a)}$$

$$f(z) = f(a) + (z-a)f'(a) + \frac{(z-a)^2}{2!}f''(a) + \dots + \frac{(z-a)^n}{n!}f^{(n)}(a) + \dots$$

This is Taylor's series of $f(z)$

if $z-a = h$

$$f(a+h) = f(a) + h f'(a) + \frac{h^2}{2!} f''(a) + \dots + \frac{h^n}{n!} f^{(n)}(a) + \dots \quad \text{if } |h| \text{ is small}$$

$$a=0, h=z$$

$$f(z) = f(0) + z f'(0) + \frac{z^2}{2!} f''(0) + \dots + \frac{z^n}{n!} f^{(n)}(0) + \dots$$

$f(z) = a_0 + a_1(z-a) + a_2(z-a)^2 + \dots$

This is a Maclaurin's series of $f(z)$

Laurent series

If $f(z)$ is analytic in a ring R bounded by two concentric circles C_1 and C_2 of radii r_1 and r_2 , ($r_1 > r_2$) with centre at a then for all z in R

$$f(z) = a_0 + a_1(z-a) + a_2(z-a)^2 + \dots + \frac{b_1}{z-a} + \frac{b_2}{(z-a)^2} + \dots$$

$$\text{Where } a_n = \frac{1}{2\pi i} \oint_{C_1} \frac{f(w) dw}{(w-a)^{n+1}}$$

$$b_n = \frac{1}{2\pi i} \oint_{C_2} \frac{f(w) dw}{(w-a)^{n+1}}$$

$$\text{and } b_n = \frac{1}{2\pi i} \oint_{C_2} \frac{f(w) dw}{(w-a)^{n+1}}$$


Where C' is any curve in R encircling C_2

Proof: Consider cross cut PQ and $f(z)$ is analytic in the region R' bounded by PQ , z is any point in R' .

$$f(z) = \frac{1}{2\pi i} \int_{PQ} \frac{f(w) dw}{(w-z)} - \frac{1}{2\pi i} \int_{C_2} \frac{f(w) dw}{(w-z)} + \frac{1}{2\pi i} \int_{C_1} \frac{f(w) dw}{(w-z)}$$

$$f(z) = \frac{1}{2\pi i} \left[\int_{C_1} \frac{f(w) dw}{(w-z)} - \int_{C_2} \frac{f(w) dw}{(w-z)} \right] \quad \text{Equation 1}$$

∞



$$\frac{1}{(z-a)} + \frac{1}{(z-a)^2} + \dots$$

Where C_1 and C_2 are described anticlockwise

Consider

$$\frac{1}{(w-a)^2} \frac{f(w)dw}{2\pi i \oint_{C_1} \frac{f(w)dw}{(w-z)}} = \frac{1}{2\pi i \oint_{C_1} \frac{f(w)dw}{(w-a)}} + \frac{1}{2\pi i \oint_{C_1} \frac{f(w)dw}{(w-a)^2}} + \dots + \frac{1}{2\pi i \oint_{C_1} \frac{f(w)dw}{(w-a)^{n+1}}} + \dots$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{f^{(n)}(a)}{(w-a)^{n+1}}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n!} \frac{f^{(n)}(a)}{(z-a)^{n+1}} \quad \text{Equation 2}$$

Where

$$a_n = \frac{1}{n!} \frac{f^{(n)}(a)}{(w-a)^{n+1}}$$

Consider

$$\frac{1}{2\pi i \oint_{C_2} \frac{f(w)dw}{(w-z)}}$$

For C_2 , $w-a < z-a$

$$f'(z) = \frac{1}{z}, f'(1) = 1,$$

$$f''(z) = -\frac{1}{z^2}, f''(1) = -1,$$

$$f'''(z) = \frac{2}{z^3}, f'''(1) = 2, f^{(4)}(z) =$$

$$-\frac{3!}{z^4}, f^{(4)}(1) = -3!$$

$$\log z = (z-1) - \frac{1}{2}(z-1)^2 + \frac{1}{3}(z-1)^3 - \frac{1}{4}(z-1)^4 + \dots + \frac{(-1)^{n-1}}{n}(z-1)^n + \dots$$

2) Obtain all the Laurent series of the function $\frac{1}{(z+1)z(z-2)}$ about $z = -1$

Solution:

$$f(z) = \frac{1}{(z+1)z(z-2)}$$

put $z+1 = u, z = u-1$

$$2 = u-3$$

$$\frac{1}{(z+1)z(z-2)} = \frac{1}{u(u-1)(u-3)} = \frac{A}{u} + \frac{B}{u-1} + \frac{C}{u-3}$$

$$\lim_{u \rightarrow 0} \frac{u - (1 - (u - 3) \sqrt{7u - 9})}{u}$$

$$u \rightarrow 1 \text{ u } (u-3) \\ 7u-9 \text{ C}$$

()

$$= -3 - (1+u+u^2+u^3+\dots) - (1+u+u^2+\dots)u^3 \quad \frac{2}{3} \quad - \quad -$$

3) Expand $\frac{1}{(z^2 - 3z + 2)}$ is the region

Solution:

(i) $|z - 1| < 1$ $\frac{1}{(z^2 - 3z + 2)} = \frac{1}{(z - 2)} - \frac{1}{(z - 1)} =$

$$\frac{1}{(z-2)} - \frac{1}{(z-1)} = \frac{1}{(z-1-1)} - \frac{1}{(z-1)}$$

$$= - \frac{1}{[1-(z-1)]} - \frac{1}{(z-1)} = \left(1 - (z-1)\right)^{-1} - \frac{1}{(z-1)}$$

$$= - (1+(z-1) + (z-1)^2 + (z-1)^3 + \dots) - \frac{1}{(z-1)}$$

(ii)

$$\left| \frac{z}{|z|} \right| < \frac{z}{|z|}, z < 2, |z| < 1, |z| < 1$$

$$\frac{1}{(z-2)} = \frac{1}{2} \left(1 - \frac{z}{2}\right)^{-1} - \frac{1}{z} \left(1 - \frac{1}{z}\right)^{-1}$$

$$= \frac{1}{2} (1 + \frac{z}{2} + \frac{z^2}{4} + \frac{z^3}{8} + \dots) - \frac{1}{z} (1 + \frac{1}{z} + \frac{1}{z^2} + \dots)$$

$$= \frac{1}{2} (1 + \frac{z}{2} + \frac{z^2}{4} + \frac{z^3}{8} + \dots) - \frac{1}{z} (1 + \frac{1}{z} + \frac{1}{z^2} + \dots)$$

(iii)

$$\overline{|z|} > 2, 2 < |z|, |z| < 1, z$$

$$\frac{1}{z(1-z^2)} - \frac{1}{z(1-\frac{1}{z})} = \frac{1}{z(1-z^2)} - \frac{1}{z(1-\frac{1}{z})}$$

$$= \frac{1}{z(1-z^2)} - \frac{1}{z(1-\frac{1}{z})}$$

$$= \frac{1}{z(1-z^2)} - \frac{1}{z(1-\frac{1}{z})}$$

$$(1 + \frac{1}{z} + \dots) - \frac{1}{z(1-z^2)}$$

$$n-1$$

$$1 - \frac{1}{n-1}$$

$$= \frac{1}{z}$$

$$(2 = \sum_{n=1}^{\infty} \frac{1}{n})$$

$$z$$

$$= \sum_{n=1}^{\infty} \frac{2}{z^n} - \sum_{n=1}^{\infty} \frac{1}{z^n}$$

$$\frac{2}{z^n} - \frac{1}{z^n}$$

$$\frac{2}{z} + \frac{2^2}{z^2} + \dots$$

$$\frac{1}{z^n}$$

$$\frac{1}{z} (1 + \frac{1}{z} + \frac{1}{z^2} + \dots)$$

$$(z^2-1)$$

Laurent series expansion of the function _____ if $2 < z < 3$.

$$(z+2)(z+3)$$

Solution:

$$f(z) = \frac{(z^2-1)}{(z+2)(z+3)} = 1 - \frac{(5z+7)}{(z^2+5z+6)}$$

$$\frac{3}{8} = 1 +$$

-

$$(z+2) (z+3)$$

$$= 1 +$$

$$\frac{3}{z(1+\frac{2}{z})}$$

$$\frac{8}{3(1+\frac{z}{3})}$$

$$\frac{1}{z}$$

$$= 1 + \frac{3}{z} \left(1 + \frac{2}{z}\right)^{-1} - \frac{8}{3} \left(1 + \frac{z}{3}\right)^{-1}$$

$$= \frac{3}{z} \left(1 - \frac{2}{z} + \frac{2^2}{z^2} - \frac{2^3}{z^3} + \dots\right) - \frac{8}{3} \left(1 - \frac{z}{3} + \frac{z^2}{3^2} - \frac{z^3}{3^3} + \dots\right)$$

$$= 1 + \dots$$

$$\begin{aligned}
 &= 1 + 3 \sum_{n=1}^{\infty} \frac{(-1)^{n-1} n^{-1} z^{n-1}}{z^n} + 8 \sum_{n=1}^{\infty} \frac{(-1)^n}{3} \frac{8z^{n-1}}{z^n} \\
 &= 1 + \sum_{n=1}^{\infty} (-1)^n \left(-\frac{2^{n-1}}{z^n} + \frac{8z^{n-1}}{3z^n} \right)
 \end{aligned}$$

e^{2z}

5) Expand $f(z) = \frac{e^{2z}}{(z-1)^3}$ about $z=1$ as Laurent series. Also indicate the region of convergence of the series.

Solution:

$$\begin{aligned}
 f(z) &= \frac{e^{2z}}{(z-1)^3} \\
 \text{put } z-1 &= u, z = 1+u \\
 \frac{e^{2z}}{(z-1)^3} &= \frac{e^{2(1+u)}}{u^3} = \frac{e^2 e^{2u}}{u^3} = \frac{e^2}{u^3} \left(1 + 2u + \frac{(2u)^2}{2!} + \dots \right) \\
 &= \frac{e^2}{(z-1)^3} \left(1 + 2(z-1) + \frac{(2(z-1))^2}{2!} + \dots \right) \\
 &= e^2 \left(\frac{1}{(z-1)^3} + \frac{1}{(z-1)^2} + \frac{2}{z-1} + \dots \right)
 \end{aligned}$$

z

6) Express $f(z) = \frac{z}{(z-1)(z-3)}$ in a series of positive and negative powers of $z-1$.

Solution:

$$f(z) = \frac{z}{(z-1)(z-3)}$$

z A B

$$\frac{1}{z} = \frac{1}{(z-1)(z-3)} = \frac{1}{(z-1)} - \frac{1}{(z-3)}$$

$$A = \lim_{z \rightarrow 1} \frac{1}{z-3} = -\frac{1}{2}$$

$$B = \lim_{z \rightarrow 3} \frac{1}{z-1} = \frac{1}{2}$$

$$\begin{aligned} f(z) &= \frac{3}{2(z-3)} - \frac{1}{2(z-1)} = \frac{3}{2(z-1-2)} - \frac{1}{2(z-1)} \\ &= \frac{3}{-4(1-\frac{z-1}{2})} - \frac{1}{2(z-1)} \\ &= -\frac{3}{4} \left(1 - \frac{z-1}{2}\right)^{-1} - \frac{1}{2(z-1)} \\ &= -\frac{3}{4} \sum_{n=0}^{\infty} \left(\frac{z-1}{2}\right)^n - \frac{1}{2(z-1)} \end{aligned}$$

Contour Integration

Singular points

Singular point: A point at which $f(z)$ ceases to be analytic is called a singular point.

Isolated singular point: Suppose $z=a$ is a singular point of a function $f(z)$ and no other singular point of $f(z)$ exists in a circle with centre at a , then $z=a$ is said to be an isolated singular point.

In such a case $f(z)$ can be expanded by Laurent series around $z=a$

Pole: If the principal part of $f(z)$ consists of a finite number of terms b_1, b_2, \dots, b_n $b_n \neq 0$

then $(z-a)$ is said to be a pole of order n .

if $n=1$, $z=a$ is said to be a simple pole. (note: if $f(z)$ has a pole at $z=a$, then $\lim_{z \rightarrow a} f(z) = \infty$)

Removable singularity: If a single valued function $f(z)$ is not defined at $z=a$ $\lim_{z \rightarrow a} f(z)$ exists and $f(z)$ exists, then $z=a$ is said to be a removable singularity $f(z) = \lim_{z \rightarrow a} f(z)$, $z=0$ is a removable singularity.

Essential singularity: If the principal part of $f(z)$ consists of an infinite number of terms, then $z=a$ is said to be an essential singularity

$e_z = 1 + \frac{1}{z} + \frac{1}{2!z^2} + \frac{1}{3!z^3} + \dots$ $z=0$ is an essential singularity.

Singularity at infinity: Suppose we substitute $z = \frac{1}{w}$, $f(\frac{1}{w}) = F(w)$ (say), then the singularity at $w=0$ of $F(w)$ is called the w singularity at infinity. e^z has an essential singularity at $z = \infty$, since e_z has an essential singularity at $z=0$.

Entire function: A function which is analytic everywhere in the finite plane is called an entire function or integral function.

Examples: e^z , $\sin z$, $\cos z$ are entire functions.

Note: An entire function can be represented by a Taylor series which has an infinite radius of convergence. Conversely, if a power series has an infinite radius of convergence, it represents an entire function.

Liouville's theorem: If $f(z)$ is analytic and bounded, i.e. $|f(z)| < m$ for some constant m in the entire complex plane, then $f(z)$ is a constant.

Residue: We know that $\oint_C \frac{f(z)}{(z-a)^{n+1}} dz = 2\pi i b_n$ where C is $|z-a|=R$ and $\oint_C \frac{f(z)}{(z-a)^{n+1}} dz = 0$, if $n \neq -1$.

$\oint_C f(z) dz = 2\pi i b_{-1}$ where C is the circle with centre at a and $f(z)$ is expanded in Laurent series. b_{-1} is said to be the residue of $f(z)$ at $z=a$ [the coefficient of $\frac{1}{(z-a)}$ in the principal part of the Laurent series of $f(z)$].

Cauchy's Residue Theorem:

Statement: If $f(z)$ is an analytic function inside and on a closed curve 'C' except at a finite number of points, inside C, then

$$\oint_C f(z) dz = 2\pi i \left(\sum \text{residues at the points where } f(z) \text{ is not analytic and which lie inside } C \right).$$

If the poles are of order one and n then the residues are

$$\lim_{z \rightarrow a} \frac{d}{dz} [(z-a)^n f(z)], \lim_{z \rightarrow a} \frac{1}{(n-1)!} \frac{d^{n-1}}{dz^{n-1}} [(z-a)^n f(z)]$$

1) Find the poles of the function and the corresponding residues at each pole, $f(z) = \frac{1}{e^z(z^2+1)}$

Solution: The given function is $f(z) = \frac{1}{e^z(z^2+1)}$, $f(z)$ is not analytic at $z = i$ and $z = -i$

Therefore, the poles of $f(z)$ are i and $-i$, both are simple poles. If $z=a$ is a simple pole, then the residue at $z=a$ is $\lim_{z \rightarrow a} (z-a)f(z)$

$$\text{Res } z=i = \lim_{z \rightarrow i} (z-i)f(z) = \lim_{z \rightarrow i} \frac{1}{e^z(z+i)} = \frac{1}{e^i(2i)} = -\frac{1}{2e^i}$$

$$\text{Res } z = -i = \lim_{z \rightarrow -i} (z+i) f(z) = \lim_{z \rightarrow -i} (z+i) \frac{e^{iz}}{(z-i)(z+i)^2} = \lim_{z \rightarrow -i} \frac{e^{iz}}{(z-i)} = \frac{e^{-i}}{-2i} = \frac{i}{2}$$

2) Find the poles of the function and the corresponding residues at each pole, $f(z) = \frac{\sin^2 z}{(z-\pi)^2}$

Solution:

The given function is $f(z) = \frac{\sin^2 z}{(z-\pi)^2}$, $z = \pi$ is a double pole

$$\begin{aligned} \text{Res at } z = \pi &= \lim_{z \rightarrow \pi} \frac{d}{dz} \left[(z-\pi)^2 f(z) \right] \\ &= \lim_{z \rightarrow \pi} 2 \sin z \cos z = 2 \sin \pi \cos \pi = 0 \end{aligned}$$

$z \sin z$

3) Find the residue of $\frac{z \sin z}{(z-\pi)^3}$ at $z = \pi$.

Solution: The given function is $f(z) = \frac{z \sin z}{(z-\pi)^3}$, $z = \pi$ is a pole of order 3

If $z = a$ is a pole of order 3, then residue at $z = a$ is

$$\lim_{z \rightarrow a} \frac{1}{(z-a)^{n-1}} \frac{d^{n-1}}{dz^{n-1}} f(z) \quad [(z-a)^{n-1} f(z)] \quad (a = \pi)$$

$$\text{Res at } z = \pi = \lim_{z \rightarrow \pi} \frac{d}{dz} (z \sin z)$$

$$= \lim_{z \rightarrow \pi} \frac{1}{2} (z^2 \cos z + \sin z) = \lim_{z \rightarrow \pi} \frac{1}{2} (z^2 \cos z + \sin z) = -1.$$

4) Evaluate $\oint_C \frac{(\cos \pi z^2 + \sin \pi z^2) dz}{(z-1)^2(z-2)}$ where C is $|z| = 3$.

Solution: The given function and $z = 2$ is a simple pole,

$$\text{Res at } z = 1 = \lim_{z \rightarrow 1} \frac{d}{dz} \left(\frac{(\cos \pi z^2 + \sin \pi z^2)}{(z-2)} \right)$$

$$= \lim_{z \rightarrow 1} \frac{(z-2)(-2z \sin \pi z^2 + 2z \cos \pi z^2) - \cos \pi z^2 - \sin \pi z^2}{(z-2)^2} = 1$$

is $f(z) = \frac{1}{z-1}$ is a double pole both lie inside C.

$$\oint_C [z-1]^{-2} f(z) dz = \lim_{z \rightarrow 1} \frac{d}{dz} (\cos \pi z^2 + \sin \pi z^2)$$

$$\text{Res at } z = 2 = \lim_{z \rightarrow 2} (z-2) f(z) = \lim_{z \rightarrow 2} \frac{(\cos \pi z^2 + \sin \pi z^2) dz}{2(z-2)}$$

According to residue theorem

$$-\left(\frac{(\cos \pi z^2 + \sin \pi z^2) dz}{2(z-2)} \right) = -2\pi i (\text{sum of the residues}) = 2\pi i (3+1) = 8\pi i$$

$$\int_C z \sec z \, dz \quad x^2 + 9y^2 = 9$$

5) Evaluate $\int_C \frac{1}{1-z^2} dz$ where C is $4x^2 + 9y^2 = 9$

Solution:

The given function is $f(z) = \frac{1}{1-z^2}$. $z=1$ and -1 are simple poles and $4x^2 + 9y^2 = 9$ is an ellipse whose semi major and minor axes are 1 and $\frac{3}{2}$. 1 and -1 both

lie inside C.

$$\text{Res at } z=1 = \lim_{z \rightarrow 1} (z-1)f(z) = \lim_{z \rightarrow 1} \frac{1}{z+1} = \frac{1}{2}$$


$$\text{Res at } z=-1 = \lim_{z \rightarrow -1} (z+1)f(z) = \lim_{z \rightarrow -1} \frac{1}{z-1} = -\frac{1}{2}$$

$$\int_C z \sec z \, dz = \int_C \frac{1}{1-z^2} dz = 2\pi i (\text{sum of the residues, by residue}$$

theorem)

$$= 2\pi i (-\frac{1}{2} - \frac{1}{2}) = -2\pi i$$

$$e^z dz$$



6) Evaluate $\oint_C \frac{e^z dz}{(z+2)(z-1)}$ Where C is the circle $|z-1|=1$.

Solution: The given function is $f(z) = \frac{e^z}{(z+2)(z-1)}$, $z = -2$ and 1 are simple poles, $z=1$ lies inside C and $z = -2$ lies outside C.

$$(z-1)f(z) = \lim_{z \rightarrow 1} \frac{e^z}{z+2} = \frac{e}{3} \quad \text{Res at } z=1 = \lim_{z \rightarrow 1} \frac{e^z}{z+2}$$

$$\oint_C f(z) dz = 2\pi i$$

(sum of residues at the poles which lie inside C)

$$\oint_C \frac{e^z dz}{(z+2)(z-1)} = 2\pi i \cdot \frac{e}{3}$$

$$\oint_C \frac{e^z dz}{(z+2)(z-1)} = \frac{2\pi i e}{3}$$


Evaluation of real integrals in unit circle

We can evaluate the integrals of the type $\int_0^{2\pi} f(\cos \theta, \sin \theta) d\theta$ where $f(\cos \theta, \sin \theta)$ is a rational function, using residue theorem.

$e^{i\theta}$, we can write $\cos \theta = \frac{z + \frac{1}{z}}{2}$

$e^{i\theta} + e^{-i\theta}$ we know that if $z = e^{i\theta}$

$$\cos \theta = \frac{1}{2} \left(z + \frac{1}{z} \right) \text{ and } \sin \theta = \frac{1}{2i} \left(z - \frac{1}{z} \right)$$



$$\sin \theta = \frac{(z - \frac{1}{z})}{2i}$$

$$\begin{aligned} e^{i\theta} &= \frac{dz}{iz} \\ \text{and} \\ d\theta &= \frac{dz}{iz} \end{aligned}$$

By this substitution we can change the integral into a function of z .

We know that $\oint_C f(z) dz = 2\pi i$ (sum of the integrals)

We

||

take C is $z=1$, then θ varies from 0 to 2π

$$\int_0^{2\pi} f(\cos\theta, \sin\theta) d\theta = \oint_C g(z) dz \quad \text{where } C \text{ is } z=1$$

$$g(z) = f\left[\frac{1}{2}\left(z + \frac{1}{z}\right), \frac{1}{2i}\left(z - \frac{1}{z}\right)\right] \frac{dz}{iz}$$

We can evaluate using residue theorem

Problems

$$\int_0^{2\pi} d\theta \quad \int_0^{2\pi}$$

1) Show that $\int_0^{2\pi} (a + b \sin \theta) d\theta = \sqrt{a^2 - b^2}$, $a > b > 0$ using residue theorem.

Solution:

Consider $C = |z|=1$, $z = e^{i\theta}$

$$\cos \theta = \frac{1}{2} \left(z + \frac{1}{z} \right), \quad \sin \theta = \frac{1}{2i} \left(z - \frac{1}{z} \right)$$

$$\int_0^{2\pi} (a + b \sin \theta) d\theta = \int_C \left[a + 2bi \left(z - \frac{1}{z} \right) \right] \frac{dz}{iz}$$

$$= \int_C \left[\frac{a}{z} + 2b - \frac{2b}{z} \right] dz$$

$$f(z) = \frac{a - 2b}{z} + 2b$$

$$\int_C f(z) dz$$

$$= \int_C \left[\frac{a - 2b}{z} + 2b \right] dz$$

$$bz^2 + 2aiz - b = b(z - \alpha)(z - \beta)$$

$$\text{where } (\alpha + \beta) = -\frac{2ai}{b}, \quad \alpha\beta = -\frac{b}{b}$$

$$\alpha = \frac{-ai + i\sqrt{a^2 - b^2}}{b} \quad \beta = \frac{-ai - i\sqrt{a^2 - b^2}}{b}$$

$$\alpha = \frac{-ai + i\sqrt{a^2 - b^2}}{b} \quad \beta = \frac{-ai - i\sqrt{a^2 - b^2}}{b}$$

$$\alpha < 1 \text{ and } \beta > 1 \quad \alpha \text{ lies in } C \quad \int_C f(z) dz = 2\pi i \operatorname{Res} f(z) \text{ at } \alpha$$

$$\operatorname{Res} f(z) \text{ at } \alpha = \lim_{z \rightarrow \alpha} (z - \alpha) f(z) = \lim_{z \rightarrow \alpha} \frac{a - 2b}{z} = \frac{a - 2b}{\alpha}$$

$$b(\alpha - \beta)$$

$$= \frac{2}{b \left[\frac{-ai + i\sqrt{a^2 - b^2}}{b} + \frac{ai + i\sqrt{a^2 - b^2}}{b} \right]}$$

$$= \frac{1}{i\sqrt{a^2 - b^2}}$$

$$\oint_C f(z) dz = \frac{1}{i} \frac{2 \pi i}{2\pi i \sqrt{a^2 - b^2}} = \frac{2\pi i}{i\sqrt{a^2 - b^2}}$$

$$\int_0^{2\pi} \frac{d\theta}{a + b \sin \theta} = \frac{2\pi}{\sqrt{a^2 - b^2}}$$

2) Evaluate $\int_0^{2\pi} \frac{d\theta}{(6 - 3 \cos \theta)^2}$ using residue theorem

Solution: $\int_0^{2\pi} \frac{d\theta}{(6 - 3 \cos \theta)^2}$

Substitute $z = e^{i\theta}$

$$\cos \theta = \frac{1}{2} \left(z + \frac{1}{z} \right), \quad \sin \theta = \frac{1}{2i} \left(z - \frac{1}{z} \right)$$

$$dz = i e^{i\theta} d\theta \quad \text{and} \quad d\theta = \frac{dz}{iz}$$

$$\int_0^{2\pi} \frac{d\theta}{(6 - 3 \cos \theta)^2} = \oint_C \frac{dz}{4z^2 (z^2 - 4z + 3)^2}$$

$$\oint_C \frac{dz}{(z^2 - 4z + 1)^2} = \frac{1}{2\pi i} \oint_C \frac{dz}{(z - \alpha)^2 (z - \beta)^2}$$

The poles are α and β where $\alpha = 2 - \sqrt{3}$ and $\beta = 2 + \sqrt{3}$ and both are double poles, among which α lies inside C .

$$\begin{aligned} \text{Res at } z = \alpha &= \lim_{z \rightarrow \alpha} \frac{d}{dz} [(z - \alpha)^2 f(z)] \\ &= \lim_{z \rightarrow \alpha} \frac{d}{dz} \left[\frac{1}{(z - \beta)^2} \right] = \frac{-2}{(\alpha - \beta)^3} \end{aligned}$$

$$\begin{aligned} (\alpha + \beta) &= 4, \quad \alpha - \beta = -2\sqrt{3} \\ \text{Res at } z = \alpha &= \frac{1}{2\pi i} \oint_C \frac{dz}{(z - \alpha)^2 (z - \beta)^2} \\ &= \frac{1}{2\pi i} \left[\frac{4}{24\sqrt{3}} - \frac{1}{6\sqrt{3}} \right] = \frac{1}{24\sqrt{3}} \end{aligned}$$

3) Evaluate $\oint_C \frac{dz}{(a + b \cos \theta)^2}$, $a > b > 0$ using residue theorem

Solution:

$$\begin{aligned} \oint_C \frac{dz}{(a + b \cos \theta)^2} &= \oint_C \frac{dz}{(a + \frac{b}{2} (z + \frac{1}{z}))^2} \\ \text{put } z &= e^{i\theta}, \quad \frac{1}{2} \oint_C \frac{dz}{(a + \frac{b}{2} (z + \frac{1}{z}))^2} \\ dz &= e^{i\theta} d\theta \\ \cos \theta &= \frac{z + \frac{1}{z}}{2} \\ \oint_C \frac{dz}{(a + \frac{b}{2} (z + \frac{1}{z}))^2} &= \oint_C \frac{4z dz}{(2a + b(z + \frac{1}{z}))^2} \end{aligned}$$

$$\oint_C \frac{dz}{(a + b \cos \theta)^2} = \frac{1}{2\pi i} \oint_C \frac{4z dz}{(2a + b(z + \frac{1}{z}))^2}$$

The poles are α and β , both are double poles

$$\alpha = \frac{-a + \sqrt{a^2 - b^2}}{b}, \quad \beta = \frac{-a - \sqrt{a^2 - b^2}}{b}$$

Where $\alpha =$ and β

$$= b$$

a lies inside C

$$\frac{d}{dz}$$

$$\text{Residue at } z = \alpha = \lim_{z \rightarrow \alpha} (z - \alpha) \frac{d}{dz} \left[\frac{1}{(z - \beta)^2} \right]$$

$$\frac{1}{(z - \beta)^2}$$

$$= - \left(\frac{2}{(z - \beta)^3} \right)$$

$$b(\alpha - \beta)$$

$$\frac{1 - 2ab^3}{2}$$

$$a$$

$$= -b \left(\frac{1 - 2ab^3}{b^8(a^2 - b^2)^{3/2}} \right) = \frac{a}{4(a^2 - b^2)^{3/2}}$$

$$2\pi \int_0^{2\pi} d\theta$$

$$0 \quad \frac{1}{(a + b \cos \theta)^2} = 2\pi i \text{ (Res } z = \alpha \text{ by residue theorem)}$$

$$\frac{2\pi i a^4}{2}$$

$$\frac{2\pi a}{2}$$

$$2$$

$$=$$

$$\frac{3}{2} = \frac{3}{2}$$

$$4i(a^2 - b^2)(a^2 - b^2)^{3/2}$$

Contour integration when the poles lie on imaginary axis

$$f(x)$$

We can evaluate integrals of the type

$$\int_{-\infty}^{\infty} \frac{h(x)}{g(x)} dx = h(x), \text{ using residue theorem.}$$

Consider $\oint_C h(z) dz$ when the poles of $h(z)$ lie on imaginary axis. We take positive imaginary axis. Integration is taken over the semicircle and the line $-R$ to R . The poles lie on upper half plane. If the poles lie on real axis

$$\int_{-R}^R h(x) dx = -\oint_C h(z) dz$$


$$\int_{-R}^R h(x) dx + \int_r h(z) dz$$

We know that by residue theorem $\oint_C h(z) dz = 2\pi i$ (sum of the residues of $h(z)$ at its poles which lie on upper half plane)

$$\int_{-R}^R h(x) dx + \int_r h(z) dz = 2\pi i (\text{sum of the residues})$$

In the limiting case $R \rightarrow \infty$ we get

$$\int_{-\infty}^{\infty} h(x) dx \quad (\text{if } \int_r h(z) dz = 0)$$



Problems:

Evaluate by contour integration $\int_{-\infty}^{\infty} \frac{dx}{1+x^2}$ **1)**


Solution: Consider $\oint_C \frac{1}{1+z^2} dz$ where C is the contour consisting of semicircle r and the line (diameter) from $-R$ to R .

dz dz dz

$$\frac{dz}{1+z^2} = \frac{R}{1+z^2} + \frac{r}{1+z^2}$$

$$\frac{dz}{1+z^2} = 0$$

$$-\infty < x < \infty$$



The poles of $f(z)$ are $\pm i$, i lie on upper half plane.

$$\text{Res at } z=i = \lim_{z \rightarrow i} (z-i) f(z) = \lim_{z \rightarrow i} \frac{z}{1+z^2} = \frac{i}{2i} = \frac{1}{2}$$

(residue at $z=i$)

$$= 2\pi i \cdot \frac{1}{2} = \pi$$

$$2 \int_0^{\infty} \frac{dx}{1+x^2} = \int_{-\infty}^{\infty} \frac{dx}{1+x^2} \quad [f(x) \text{ is even}]$$

$$= \frac{1}{2} \int_{-\infty}^{\infty} \frac{dz}{1+z^2} = \frac{1}{2} \cdot \pi$$

Solution:

2) Evaluate $\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)(4+x^2)} dx$ using residue theorem.

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-R}^R f(z) dz + \int_{\Gamma} f(z) dz \quad [\int_{\Gamma} f(z) dz = 0]$$

$$= \oint_C f(z) dz = \oint_C f(z) dz$$

The poles of $f(z) = \frac{1}{z^2(1+z^2)(4+z^2)}$ are $i, -i, 2i, -2i$.

$$\frac{1}{z^2(1+z^2)(4+z^2)}$$

All are simple poles i and $-i$ lie on upper half plane.

Res at $z=i = \lim_{z \rightarrow i} (z-i)f(z)$

$$= \lim_{z \rightarrow i} \frac{1}{(1+z)(4+z^2)} = -\frac{1}{6i}$$

Res at

$$z=2i = \lim_{z \rightarrow 2i} (z-2i)f(z)$$

$$= \lim_{z \rightarrow 2i} \frac{1}{(z+2i)(1+z^2)} = -\frac{1}{4i(-3)} = \frac{3}{4i}$$

According to residue theorem

$$\oint_C f(z) dz$$

$2\pi i$ (sum of residues)

$$= 2\pi i \left(\frac{1}{-6i} + \frac{3}{4i} \right) = 3$$

$$\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)(4+x^2)} dx = 3$$

$$\int_{-\infty}^{\infty} \frac{x^2}{(1+x^2)(4+x^2)} dx$$

3) Evaluate $\int_{-\infty}^{\infty} \frac{x^5}{1+x^6} dx$ using residue theorem.

Solution:

$$-\int_{-\infty}^{\infty} f(x) dx$$

$$= \int_{-R}^R f(z) dz + \int_{\Gamma} f(z) dz \quad \left[\int_{\Gamma} f(z) dz = 0 \right]$$

$$= \int_{-R}^R f(z) dz$$

$$-\int_{-R}^R f(x) dx = \int_{-R}^R f(z) dz$$

The poles are $e^{(2n+1)\pi i/6}$ where $n=0,1,2,3,4,5$

$$[-1 = \cos \pi + i \sin \pi = e^{-\pi i} = \cos(2n+1)\pi + i \sin(2n+1)\pi]$$

$$(-1)^{\frac{1}{6}} = \frac{\cos(2n+1)\pi}{6} + i \frac{\sin(2n+1)\pi}{6} = e^{2n+1\pi i/6}$$

When $n = 0, 1, 2$ i.e., $e^{\pi i/6}, e^{\pi i/2}, e^{5\pi i/6}$ lie on upper half plane.

$$\text{Res at } z \rightarrow e^{\pi i/6} = \lim_{z \rightarrow e^{\pi i/6}} (z - e^{\pi i/6}) f(z)$$


form $\frac{0}{0}$

$$= \lim_{z \rightarrow e^{\pi i/6}} \frac{z^2(z - e^{\pi i/6})}{(1+z^6)}$$

$$= \lim_{z \rightarrow e^{\pi i/6}} \frac{(3z^2 - 2z e^{-\pi i/6})}{6z^5}$$

πi

$$z \rightarrow e^{\pi i/6}$$



$$\frac{(3z-2e^{-z/6})}{6z^4} = \lim$$

$$\pi i \quad z \rightarrow e^{\pi i/6}$$

$$-3\pi i$$

$$\frac{e^{-z/6}}{6z^4} = \frac{1}{6} e^{-\frac{1}{2} \frac{\pi}{2}} = e^{-\frac{\pi}{2}} = (\cos \frac{\pi}{2} - i \sin \frac{\pi}{2}) = -i$$

$$6e$$

Res at $z \rightarrow e^{3\pi i/6} = \lim_{z \rightarrow e^{\pi i}} = (z - e^{\pi i})^0 f(z)$ form

$$\frac{3\pi i}{6}$$

$$z \rightarrow e^{3\pi i/6}$$

$$= \lim_{z \rightarrow e^{\pi i}} \frac{z^2(z - e^{\pi i})}{(1+z^6)}$$

$$= \lim_{z \rightarrow e^{2\pi i}} \frac{(3z^2 - 2ze^{-z/2})}{6z^5} \quad \pi i$$

$$\pi i$$

$$= \lim_{z \rightarrow e^{2\pi i}} \frac{(3z - 2e^{-z/2})}{6z^4}$$

$$\pi i$$

$$\frac{1}{6} e^{-\frac{3\pi}{2}} = (\cos \frac{3\pi}{2} - i \sin \frac{3\pi}{2}) = i$$

$$5\pi i \quad 5\pi i$$

$$\begin{aligned}
 \text{Res at } z \rightarrow e^{5\pi i/6} &= \lim_{z \rightarrow e^{5\pi i/6}} (z - e^{5\pi i/6}) f(z) \\
 &= \lim_{z \rightarrow e^{5\pi i/6}} \frac{z^2 (z - e^{5\pi i/6})}{(1 + z^6)} \\
 &= \lim_{z \rightarrow e^{5\pi i/6}} \frac{z^2 (3z^5 - e^{5\pi i/6} z^5)}{6z^5} \\
 &= \lim_{z \rightarrow e^{5\pi i/6}} \frac{1}{6} e^{-15\pi i/6} = \frac{1}{6} e^{-15\pi i/6} = \frac{1}{6} e^{-5\pi i/2} = \frac{1}{6} (\cos 5\pi/2 + i \sin 5\pi/2) \\
 &= \frac{1}{6} (0 + i) = \frac{i}{6}
 \end{aligned}$$

According to residue theorem

$$\begin{aligned}
 2\pi i (\text{sum of residues}) &= \oint_C f(z) dz \\
 &= 2 \left(\frac{1}{6} + \frac{1}{6} + \frac{1}{6} \right) = \frac{1}{3} \oint_C \frac{z^2}{1+z^6} dz \\
 &= \frac{1}{3} \left(\int_0^{2\pi} \frac{z^2}{1+z^6} dz \right)
 \end{aligned}$$

$$\pi \qquad \pi i \qquad i \qquad i$$

$$- \qquad -$$

$$\infty \, dx$$

$$\infty = 3$$

$$\infty \, dx$$

$$=$$

$$6$$


$$-i \sin) = -$$

$$\frac{6}{6}$$

4) Evaluate $\int_{-\infty}^{\infty} \frac{dx}{(x^2+1)^3}$ using residue theorem.

Solution:

$$\begin{aligned} &= \int_{-R}^R f(z) dz + \int_{\Gamma} f(z) dz \quad [\int_{\Gamma} f(z) dz = 0] \\ &= \int_{-R}^R f(z) dz \\ &\int_{-R}^R \frac{1}{(z^2+1)^3} dz = \int_{-R}^R f(z) dz \end{aligned}$$



The function is $f(z) = \frac{1}{(z^2+1)^3}$. The poles are i and $-i$ of order 3, $z=i$ lies on upper half plane and inside the semicircle

$$\begin{aligned} \text{Res at } z=i &= \lim_{z \rightarrow i} \frac{1}{2!} \frac{d^2}{dz^2} [(z-i)^3 f(z)] \\ &= \lim_{z \rightarrow i} \frac{1}{2!} \frac{d^2}{dz^2} \left(\frac{1}{(z+i)^3} \right) \\ &= \lim_{z \rightarrow i} \frac{3}{(z+i)^5} \\ &= \frac{3}{(i+i)^5} = \frac{3}{(2i)^5} = \frac{3}{32i} = -\frac{3i}{32} \end{aligned}$$

$$(2i)^5 = 16$$

According to residue theorem

$$\oint_C f(z) dz = 2\pi i$$

(residue at $z = i$) $\left(\frac{1}{z-i} \right) \pi i$

$$= 2\pi i \left(\frac{3}{16i} \right) = \frac{3\pi}{8}$$

$$\int_{-\infty}^{\infty} \frac{dx}{(x^2+1)^3} = \frac{3\pi}{8}$$

Evaluation of the integrals of the type

$$\int_{-\infty}^{\infty} f(x) dx$$

∞ e **Jordan's**

Lemma

If $f(z)$ is a function of z satisfying the following properties:

- (i) $f(z)$ is analytic in upper half plane except at a finite number of poles
- (ii) $f(z) \rightarrow 0$ uniformly as $|z| \rightarrow \infty$ with $0 \leq \arg z \leq \pi$
- (iii) a is a positive integer, then

$$\lim_{r \rightarrow \infty} \oint_C f(z) e^{iaz} dz = 0$$

Where C is a semicircle with radius r and centre at the origin

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(z) dz = 2\pi i$$


(sum of the residues which lie on upper half plane)

Problems

$$\int_{-\infty}^{\infty} \frac{\cos x dx}{(x^2+16)(x^2+9)}$$

1) Evaluate $\int_{-\infty}^{\infty} \frac{\cos x dx}{(x^2+16)(x^2+9)}$ using residue theorem.

Solution: $\int_{-\infty}^{\infty} f(z) e^{imz} dz = \lim_{R \rightarrow \infty} \int_{-R}^R f(z) e^{imz} dz$



$$\int_{-R}^R f(z) e^{imz} dz$$

$\Rightarrow \int_{-R}^R f(z) e^{imz} dz = 0$ (Jordan's Lemma)

$$\int_{-\infty}^{\infty} f(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(z) dz = 2\pi i$$

-R

R

(sum of the residues which lie on upper half plane)

$$e^{iz} dz$$

$\frac{1}{(z^2+16)(z^2+9)}$ $z=3i, -3i, 4i$ and $-4i$ are simple poles. $3i$ and $4i$ lie on upper half

plane.

Res at z =

$$3i = \lim_{z \rightarrow 3i} (z -$$

$$3i)f(z)$$

$$\frac{e^{iz}}{(z^2+16)(z+3i)}$$

$$= \lim_{z \rightarrow 3i} \frac{z}{(z^2+16)(z+3i)}$$

$$\frac{e^{-3} - ie^{-3}}{(-9+16)(6i)} = \frac{-ie^{-3}}{7(6i)} = \frac{e^{-3}}{42}$$

Res at z =

$$4i = \lim_{z \rightarrow 4i} (z -$$

$$4i)f(z)$$

$$\frac{e^{iz}}{(z+4i)(z^2+9)}$$

$$= \lim_{z \rightarrow 4i} \frac{z}{(z+4i)(z^2+9)}$$

$$\frac{e^{-4} - ie^{-4}}{(9-16)(8i)} = \frac{ie^{-4}}{56}$$

$$\oint_C \frac{e^{iz} dz}{(z^2+16)(z^2+9)} = 2\pi i \left(\frac{-i}{42e^3} + \frac{i}{56e^4} \right) = \frac{\pi(4e^{-3} - 3e^{-4})}{42e^3 + 56e^4} = 84$$


$$\text{R.P} \oint_C \frac{e^{iz} dz}{(z^2+16)(z^2+9)} = \oint_C \frac{\cos z \, dz}{(z^2+16)(z^2+9)}$$

$$\int_{-\infty}^{\infty} \frac{\cos x \, dx}{(x^2+16)(x^2+9)} = \frac{\pi(4e^{-3} - 3e^{-4})}{84}$$

$$\int_0^{\infty} \frac{x \sin x \, dx}{(a^2 + x^2)}$$

2) Evaluate $\int_{-\infty}^{\infty} \frac{e^{imx}}{(a^2 + x^2)} dx$

Solution: $\int_{-\infty}^{\infty} f(z) e^{imz} dz = \lim_{R \rightarrow \infty} \int_{-R}^R f(z) e^{imz} dz + \int_{\text{arc}}$



$$\Rightarrow \int_{-\infty}^{\infty} e^{imx} f(z) dz = 0$$

$$f(z) = \frac{1}{(a^2 + z^2)}$$

$-R$ R

$z = ai$ and $-ai$ are simple poles.

Res at $z = ai = \lim_{z \rightarrow ai} (z - ai) f(z)$

$$\lim_{z \rightarrow ai}$$

$$ze^{iz}$$

$$=$$

$$\frac{z - ai}{(z + ai)^2}$$

$$= \frac{e^{-a}}{2}$$

$$= e^{-a}$$

$$= e^{-a}$$

$$= e^{-a}$$

$$= e^{-a}$$

$$= e^{-a}$$

$$\frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{e^{-ax}}{a^2 + x^2} dx = \frac{1}{2} \int_{-\infty}^{\infty} \frac{e^{-ax}}{a^2 + x^2} dx = \frac{\pi}{2} e^{-a}$$

$$= \frac{\pi}{2} e^{-a}$$

$$= \frac{\pi}{2} e^{-a}$$

Unit -3

LAPLACE TRANSFORMS

LAPLACE TRANSFORM

Definition:

Let $f(t)$ be a function of t , defined $\forall t \geq 0$. If the integral $\int_0^{\infty} e^{-st} f(t) dt$ exists, then it is called the Laplace Transform of

$f(t)$ and it is denoted by $L\{f(t)\}$ or $f(s)$.

Here s is parameter, real or complex. L is called Laplace Transform operator.

$$L\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt$$

Def: Piece-wise Continuous Function:

A function is said to be piece-wise continuous (or) Sectionally Continuous) over the closed interval $[a,b]$ if it is defined on that interval and is such that the interval can be divided into a finite number of sub intervals, in each of which $f(t)$ is continuous and both right and left hand limits at every end point of the sub intervals.

Def: Functions of Exponential Order:

A function $f(t)$ is said to be of exponential order as $t \rightarrow \infty$ if

$$\lim_{t \rightarrow \infty} (e)^{-at} f(t) = \text{finite quantity}$$

(or)

If for a given positive integer T , \exists a positive number M
 Such that $|f(t)| < Me^{at} \quad \forall t \geq T$,

Sufficient Conditions for existence of Laplace Transform are 1)

f(t) is Piece-wise Continuous Function in [a, b] where a>0, 2)

f(t) is of Exponential Order function.

Linear Property:

Theorem: If c_1, c_2 are constants and f_1, f_2 are functions of t , then

$$L[c_1 f_1(t) + c_2 f_2(t)] = c_1 L[f_1(t)] + c_2 L[f_2(t)]$$

Proof: The definition of Laplace Transform is

$$L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt \text{ -----(1)}$$

By definition

$$\begin{aligned} L[c_1 f_1(t) + c_2 f_2(t)] &= \int_0^{\infty} e^{-st} [c_1 f_1(t) + c_2 f_2(t)] dt \\ &= \int_0^{\infty} e^{-st} c_1 f_1(t) dt + \int_0^{\infty} e^{-st} c_2 f_2(t) dt \\ &= c_1 \int_0^{\infty} e^{-st} f_1(t) dt + c_2 \int_0^{\infty} e^{-st} f_2(t) dt \end{aligned}$$

$$= c_1 L[f_1(t)] + c_2 L[f(t)]$$

Laplace Transform (L.T) of some Standard Functions:

1

1) Show that $L\{1\} = \frac{1}{s}$

Solution: By definition of L.T $L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt$ -----(1)

Put $f(t)=1$ o.b.s $L[1] = \int_0^{\infty} e^{-st} \cdot 1 \cdot dt$

$$= \left[\frac{e^{-st}}{-s} \right]_0^{\infty} = \frac{1}{s}$$

2) $L[c] = L[c \cdot 1] = c \cdot L[1] = c \cdot (1/s) = c/s$

3) Show that $L[e^{at}] = \frac{1}{s-a}$

Solution: By definition of L.T,
 $L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt$ -----(1)

$$\begin{aligned} L[e^{at}] &= \int_0^{\infty} e^{-st} e^{at} dt \\ &= \int_0^{\infty} e^{-(s-a)t} dt \\ &= \left[\frac{e^{-(s-a)t}}{-(s-a)} \right]_0^{\infty} \\ &= \frac{1}{s-a} \quad (e^{-\infty} = 0) \end{aligned}$$

Put $f(t) = e^{at}$ o.b.s in (1) $L[$

Note: $L[e^{-at}] = \frac{1}{s+a}$

4) Show that $L[\cos at] = \frac{s}{s^2+a^2}$ and $L[\sin at] = \frac{a}{s^2+a^2}$

Solution: W.k.t $e^{i\theta} = \cos \theta + i \sin \theta$

$$e^{iat} = \cos at + i \sin at$$

$$L[e^{iat}] = L[\cos at + i \sin at]$$

$$L[\cos at + i \sin at] = L[e^{iat}]$$

$$\begin{aligned} &= \frac{1}{s-ia} \quad (L[e^{at}] = \frac{1}{s-a}) \\ &= \frac{s+ia}{(s-ia)(s+ia)} \\ &= \frac{s+ia}{s^2+a^2} \\ &= \frac{s}{s^2+a^2} + i \frac{a}{s^2+a^2} \end{aligned}$$

Equating real and imaginary parts we get

$$L[\cos at] = \frac{s}{s^2+a^2} \quad \text{and} \quad L[\sin at] = \frac{a}{s^2+a^2}$$

5) Find $L[\sin at]$

$$\frac{e^{at} - e^{-at}}{2}$$

$$\text{Solution: } L[\sin at] = L\left[\frac{1}{2}\left[\frac{1}{s-a} - \frac{1}{s+a}\right]\right] = \frac{1}{2}[L\{e^{at}\} - L\{e^{-at}\}]$$

$$= \frac{1}{2}\left[\frac{s+a-s+a}{s^2-a^2}\right]$$

$$= \frac{a}{s^2-a^2}$$

6) Find $L[\cos at]$

$$\frac{e^{at} + e^{-at}}{2}$$

$$\text{Solution: } L[\cos at] = L\left[\frac{1}{2}(e^{at} + e^{-at})\right] = \frac{1}{2}[L\{e^{at}\} + L\{e^{-at}\}]$$

$$= \frac{1}{2}\left[\frac{1}{s-a} + \frac{1}{s+a}\right]$$

$$= \frac{1}{2}\left[\frac{s+a+s-a}{s^2-a^2}\right] = \frac{s}{s^2-a^2}$$

s

7) Show that (i) $L[t^n] = \frac{n!}{s^{n+1}}, \quad n > -1$

(ii) $L[t^n] = \frac{n!}{s^{n+1}}, \quad n \text{ is +ve integer}$

Solution: : By definition of L.T

$$L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt \text{-----(1)}$$

$$L[t^n] = \int_0^{\infty} e^{-st} t^n dt \quad \text{put } st = x \text{ i.e } t = x/s$$

$$= \int_0^{\infty} e^{-x} \left(\frac{x}{s}\right)^n \frac{dx}{s} \quad dt = \frac{dx}{s}$$

$$= \frac{1}{s^{n+1}} \int_0^{\infty} e^{-x} x^n dx$$

$$= \frac{1}{s^{n+1}} \rho(n+1), \quad \text{for } (n+1) > 0$$

$$L[t^n] = \rho(n+1)/s^{n+1}, \quad n > -1$$

$$L[t^n] = n!/s^{n+1}, \quad n \text{ is +ve integer}$$

FORMULAE

1

$$1) \quad L\{1\} =$$

$\frac{s}{c}$

$$2) \quad L\{c\} =$$

$\frac{1}{s}$

$$3) \quad L[e^{at}] = \frac{1}{s-a} \quad , \quad L[e^{-at}] = \frac{1}{s+a}$$

s

$$4) \quad L[\cos at] = \frac{a}{s^2 + a^2}$$

$$5) \quad L[\sin at] = \frac{a}{s^2 + a^2}$$

$$6) \quad L[\sin hat] = \frac{s}{s^2 - a^2}$$

$$7) \quad L[\cos hat] = \frac{s}{s^2 - a^2}$$

$$8) \quad L(t^n) = \frac{n!}{s^{n+1}}, \quad n > -1$$

$$9) \quad L(t^n) = \frac{n!}{s^{n+1}}, \quad n \text{ is +ve integer}$$

PROBLEMS

1. Find the Laplace Transformation (L.T) of $t^2 + 2t + 3$

$$\text{Solution: } L[t^2 + 2t + 3] = L[t^2] + 2L[t] + L[3]$$

$$= \frac{2!}{s^3} + 2 \cdot \frac{1!}{s^2} + \frac{3}{s}$$

$$t^{\frac{5}{2}} + 4] \quad \text{2. Find } L[$$

$$\text{Solution: } L[t^{\frac{5}{2}} + 4] = L[t^{\frac{5}{2}}] + L[4]$$

$$= \frac{\Gamma(\frac{7}{2})}{s^{\frac{7}{2}}} + \frac{4}{s}$$

$$\text{3. Find } L[$$

$$\text{Solution: } L[e^{3t} + 3e^{-2t}] = L[e^{3t}] +$$

$$3L[e^{-2t}]$$

$$= \frac{1}{s-3} + 3 \frac{1}{s+2}$$

$$\text{4. Find } L[\sin 3t + \cos^2 2t]$$

$$\text{Solution: } L[\sin 3t + \cos^2 2t] = L[\sin 3t] + L[\cos^2 2t]$$

$$= \frac{3}{s^2+9} + L\left[\frac{1+\cos 4t}{2}\right]$$

$$= \frac{3}{s^2+9} + \frac{1}{2} \{ L[1] + L[\cos 4t] \}$$

$$= \frac{3}{s^2+9} + \frac{1}{2} \left[\frac{1}{s} + \frac{s}{s^2+16} \right]$$

$$\text{5. Find } L[f(t)] \text{ if } f(t) = 0, \quad 0 < t < 2$$

$$= 3, \quad t > 2$$

Solution: By definition of L.T

$$\begin{aligned}
 & \int_0^{\infty} e^{-st} f(t) dt \\
 \mathcal{L}[f(t)] &= \int_0^{\infty} e^{-st} f(t) dt \\
 &= \int_0^2 e^{-st} f(t) dt + \int_2^{\infty} e^{-st} f(t) dt \\
 &= 0 + \int_2^{\infty} e^{-st} \cdot 3 dt \\
 &= 3 \left[\frac{e^{-st}}{-s} \right]_2^{\infty} \\
 &= 3 \left(\lim_{t \rightarrow \infty} \frac{e^{-st}}{-s} - \frac{e^{-2s}}{-s} \right) \\
 &= \frac{3}{s} (0 - (-e^{-2s})) \\
 &= \frac{3}{s} e^{-2s}
 \end{aligned}$$

$$e_{-\infty} = 0$$

First shifting Theorem (F.S.T):

If $\mathcal{L}[f(t)] = f(s)$ then $\mathcal{L}[e^{at} f(t)] = f(s-a)$

Proof : By definition of L.T

$$L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt = f(s) \text{-----(1)}$$

$$\begin{aligned} L[e^{at}f(t)] &= \int_0^{\infty} e^{-st} e^{at} f(t) dt \\ &= \int_0^{\infty} e^{-(s-a)t} f(t) dt \text{ Put } s-a=p = \int_0^{\infty} e^{-pt} f(t) dt \\ &= f(p) = f(s-a) \end{aligned}$$

$$\text{Note: } L[e^{-at}f(t)] = f(s+a)$$

Problems:

1) Find $L[t^3 e^{-3t}]$

Solution : let $f(t) = t^3$

$$L[f(t)] = L[t^3] = \frac{3!}{s^{3+1}} = \frac{6}{s^4} = f(s)$$

By F.S.T, $L[e^{-at}f(t)] = f(s+a)$

$$f(t) = f(s+3)$$

$$a=3 \quad L[e^{-3t}]$$

$$L[e^{-3t}t^3] = \frac{6}{(s+3)^4}$$

2) Find $L [e^{-t}(3 \sin 2t - 5 \cosh 2t)]$

Solution : Let $f(t) = (3 \sin 2t - 5 \cosh 2t)$ L

$$[f(t)] = L[(3 \sin 2t - 5 \cosh 2t)]$$

$$= 3 \frac{2}{s^2 + 4} - 5 \frac{s}{s^2 - 4} = f(s)$$

By F.S.T , $L[e^{-at} f(t)] = f(s+a)$ $a=1$

$$L[e^{-1t} f(t)] = f(s+1)$$

$$= \frac{6}{(s+1)^2 + 4} - \frac{5(s+1)}{(s+1)^2 - 4}$$

$$L [e^{-t}(3 \sin 2t - 5 \cosh 2t)] = \frac{6}{s^2 + 2s + 5} - \frac{5s + 5}{s^2 + 2s - 3}$$

Second Shifting Theorem (S.S.T)

STATEMENT:- If $L[f(t)] = f(s)$ and $g(t) = f(t-a)$, $t > a$

$= 0$, $t < a$ then $L\{g(t)\} = e^{-as} f(s)$

PROOF:- By definition of L.T

$$\begin{aligned}
 L[f(t)] &= \int_0^{\infty} e^{-st} f(t) dt = f(s) \text{-----(1)} \\
 L[g(t)] &= \int_0^{\infty} e^{-st} g(t) dt \\
 &= 0 + \int_a^{\infty} e^{-st} f(t-a) dt \quad \text{put } t-a=x \quad t=a+x \\
 &= e^{-as} \int_0^{\infty} e^{-sx} f(x) dx \quad dt=dx, (x=0 \text{ to } \infty)
 \end{aligned}$$

$$= e^{-as} f(s)$$

Example :

Find Laplace Transform of $g(t) = \begin{cases} \cos\left(t - \frac{2\pi}{3}\right), & \text{if } t > \frac{2\pi}{3} \\ 0, & \text{if } t < \frac{2\pi}{3} \end{cases}$

Solution: Let $f(t) = \cos t$, $a = \frac{2\pi}{3}$

$$f(t-a) = \cos\left(t - \frac{2\pi}{3}\right) = \cos\left(t - \frac{2\pi}{3}\right) f(t)$$

$$L[f(t)] = L\left[\cos t\right] = \frac{s}{s^2+1} = f(s)$$

[

By S.S.T $L[g(t)] = e^{-as} f(s)$

$$= \left(e^{-\frac{2\pi}{3}s}\right) \frac{s}{s^2+1}$$

Change of scale property:

If $L[f(t)] = f(s)$ then $L[f(at)] = \frac{1}{a} f\left(\frac{s}{a}\right)$

NOTE: $L\left[f\left(\frac{t}{a}\right)\right] = a f(as)$

Example: If $L[f(t)] = \frac{9s^2 - 12s + 15}{(s-1)^3}$ then find $L[f(3t)]$

Solution: Given

$$\frac{9s^2 - 12s + 15}{(s-1)^3} = f(s)$$

$L[f(t)] =$ by Change of

scale property, $L[f(at)]$

$=$

$$\frac{1}{a} f\left(\frac{s}{a}\right)$$

$$L[f(3t)] = \frac{1}{3} f\left(\frac{s}{3}\right)$$

$$= \frac{1}{3} \left[\frac{9\left(\frac{s}{3}\right)^2 - 12\left(\frac{s}{3}\right) + 15}{\left(\frac{s}{3} - 1\right)^3} \right]$$

$$= \frac{1}{3} \left[\frac{s^2 - 4s + 15}{(s-3)^3 / 27} \right]$$

$$= \frac{9(s^2 - 4s + 15)}{(s-3)^3}$$

Laplace transform of the derivative of $f(t)$

□ If $f(t)$ is continuous for all $t \geq 0$ and $f(t)$ is piecewise continuous, then

$L\{f(t)\}$ exists, provided $\lim_{t \rightarrow \infty} e^{-st}f(t) = 0$ and □□

$$L\{f'(t)\} = sL\{f(t)\} - f(0)$$

$$L\{f^{(n)}(t)\} = s^n L\{f(t)\} - s^{n-1}f(0) - s^{n-2}f'(0) - \dots - f^{(n-1)}(0)$$

Example Derive Laplace transform of $\sin at$

Let $f(t) = \sin at$ then $f'(t) = a \cos at$ and $f''(t) = -a^2 \sin at$

Also $f(0) = 0$, $f'(0) = a$ from this also $f''(0) = 0$, also from this

By derivative formula,

$$L[f''(t)] = s^2 L[f(t)] - s f(0) - f'(0) \text{-----(1)}$$

$$L\{-a^2 \sin at\} = s^2 L(\sin at) - a$$

$$(-a^2) L(\sin at) + a = s^2 L(\sin at) \quad a =$$

$$(s^2 + a^2) L(\sin at)$$

$$L(\sin at) = \frac{a}{s^2 + a^2}$$

Laplace transform of the integration of $f(t)$

$$\text{If } L[f(t)] = f(s) \text{ then } L\left[\int_0^t f(t) dt\right] = \frac{f(s)}{s}$$

Example:

Find L.T. of $\int_0^t \sin at \, dt$ Solution:

Let

$$L[f(t)] = L[\sin at] = \frac{a}{s^2 + a^2} \quad f(t) = \sin at$$

$$\int_0^t f(t) dt = \frac{f(s)}{s}$$

$$= f(s)$$

$$L\left[\int_0^t \sin at \, dt\right] = \frac{1}{s} \left(\frac{a}{s^2 + a^2}\right)$$

Multiplication by t :

$$\text{If } L[f(t)] = f(s) \text{ then } L[t f(t)] = -\frac{d}{ds} [f(s)]$$

$$L[t^2 f(t)] = (-1)^2 \frac{d^2}{ds^2} [f(s)]$$

$$= (-1)^n \frac{d^n}{ds^n} [f(s)]$$

$$L[t^n f(t)] =$$

Example : Find $L[t \sin^2 t]$

Solution: Let

$$f(t) = \sin^2 t$$

$$\begin{aligned} \text{let } \sin^2 t &= L\left[\frac{1 - \cos 2t}{2}\right] \\ \frac{1}{2} (L[1] - L[\cos 2t]) &= \frac{1}{2} \left(\frac{1}{s} - \frac{s}{s^2 + 4}\right) = \frac{2}{s(s^2 + 4)} = f(s) \\ &= -\frac{d}{ds} [f(s)] \\ &= -\frac{d}{ds} \left[\frac{2}{s(s^2 + 4)}\right] \\ &= -2 \left[\frac{-1}{\{s(s^2 + 4)\}^2}\right] \frac{d}{ds}(s(s^2 + 4)) \\ &= \left[\frac{2}{\{s(s^2 + 4)\}^2}\right] \frac{d}{ds}(s^3 + 4s) \end{aligned}$$

$$L[f(t)] = L[$$

By theorem $L[t f(t)]$

$$\begin{aligned} &= \left[\frac{2}{\{s(s^2 + 4)\}^2}\right] \\ &= \frac{6s^2 + 8}{s^2(s^2 + 4)^2} \quad] (3s^2 + 4) \text{ Division } \end{aligned}$$

by t:

If $L[f(t)] = f(s)$ then $L\left[\frac{f(t)}{t}\right] = \int_s^\infty f(s) ds$, provided $\lim_{t \rightarrow 0} \frac{f(t)}{t}$ exists.

Problems: (1) Find

$L\left[\frac{e^{-3t} - e^{-4t}}{t}\right]$

Solution: Let $f(t) = e^{-3t} - e^{-4t}$

$$L[f(t)] = L[e^{-3t} - e^{-4t}] = \frac{1}{s+3} - \frac{1}{s+4} = f(s) \text{ w.k.t}$$

$$, L\left[\frac{f(t)}{t}\right] = \int_s^\infty f(s) ds$$

$$\left[\frac{e^{-3t} - e^{-4t}}{t}\right] = \int_s^\infty \left(\frac{1}{s+3} - \frac{1}{s+4}\right) ds$$

$L\left[\frac{e^{-3t} - e^{-4t}}{t}\right]$

$$= \log(s+3) - \log(s+4)$$

$$= \log\left(\frac{s+3}{s+4}\right) = \log\left[s\left(1 + \frac{3}{s}\right) - s\left(1 + \frac{4}{s}\right)\right]$$

$$= \log 1 - \log\left(\frac{s+4}{s+3}\right)$$

$$= 0 - \log\left(\frac{s+4}{s+3}\right) = \log\left(\frac{s+3}{s+4}\right)$$

(2). Find L.T of $\frac{\cos at - \cos bt}{t}$

Solution: Let $f(t) = \cos at - \cos bt$

$$L[f(t)] = L[\cos at - \cos bt]$$

$$f(s) = \frac{s}{s^2+a^2} - \frac{s}{s^2+b^2}$$

w.k.t, $L\left[\frac{f(t)}{t}\right] = \int_s^\infty f(s) ds$

$$\left[\frac{\cos at - \cos bt}{t}\right] = \int_s^\infty \left(\frac{s}{s^2+a^2} - \frac{s}{s^2+b^2}\right) ds \quad \Bigg|_s^\infty$$

$$= \left[\log(s^2 + a^2) - \log(s^2 + b^2) \right]_s^\infty$$

$$= \left(\log(s^2 + a^2) - \log(s^2 + b^2) \right) \Bigg|_s^\infty$$

$$= \frac{1}{2} \log \left(\frac{s^2 + b^2}{s^2 + a^2} \right)$$

Evaluation of

integrals by Laplace transforms:

(1). Using L.T. Evaluate $\int_0^\infty \left[\frac{e^{-t} - e^{-2t}}{t} \right] dt$

Solution: First we will find $L\left[\frac{e^{-t} - e^{-2t}}{t}\right]$ let

$$f(t) = e^{-t} - e^{-2t}$$

$$L[f(t)] = L[e^{-t} - e^{-2t}]$$

$$= \frac{1}{s+1} - \frac{1}{s+2} = f(s)$$

w.k.t, $L\left[\frac{f(t)}{t}\right] = \int_s^\infty f(s) ds$,

$$L\left[\frac{e^{-t} - e^{-2t}}{t}\right] = \int_s^\infty \left(\frac{1}{s+1} - \frac{1}{s+2} \right) ds$$

$$= \left[\log(s+1) - \log(s+2) \right]_{s=\infty}^{\infty} = \log \left(\frac{s+1}{s+2} \right) \Big|_{s=\infty}^{\infty}$$

$$= \log \left(\frac{\infty+1}{\infty+2} \right) - \log \left(\frac{\infty+1}{\infty+2} \right)$$

$$= \log \left(\frac{\infty+1}{\infty+2} \right) - \log \left(\frac{\infty+1}{\infty+2} \right)$$

$$\left. \begin{aligned} \frac{s(1+\frac{1}{s})}{s(1+\frac{2}{s})} &= \log 1 - \log \left(\frac{s+1}{s+2} \right) \\ &= 0 - \log \left(\frac{s+1}{s+2} \right) = \log \left(\frac{s+2}{s+1} \right) \end{aligned} \right\}$$

$$\frac{e^{-t} - e^{-2t}}{t} \Big] = \log \left(\frac{s+2}{s+1} \right)$$

therefore, $L[$

The definition of Laplace Transform is

$$L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt$$

$$L\left[\frac{e^{-t} - e^{-2t}}{t}\right] = \int_0^{\infty} e^{-st} \left[\frac{e^{-t} - e^{-2t}}{t}\right] dt = \log \left(\frac{s+2}{s+1} \right)$$

Put $s=0$ on both sides

$$\int_0^{\infty} 1 \left[\frac{e^{-t} - e^{-2t}}{t}\right] dt = \log \left(\frac{2}{1} \right) = \log 2$$

$$\int_0^{\infty} \left(\frac{\cos at - \cos bt}{t} \right) dt$$

$$\frac{\cos at - \cos bt}{t} \quad L[\quad]$$

2. Using LT find

Solution: First we find

: Let $f(t) = \cos at - \cos bt$

$$L[f(t)] = L[\cos at - \cos bt] f(s)$$

$$= \frac{s}{s^2 + a^2} - \frac{s}{s^2 + b^2}$$

w.k.t, $\frac{f(t)}{t} = \int_s^\infty f(s) ds$

$$L\left[\frac{\cos at - \cos bt}{t}\right] = \int_s^\infty \left(\frac{s}{s^2+a^2} - \frac{s}{s^2+b^2}\right) ds$$

$$= \frac{1}{2} \left[\log(s^2+a^2) - \log(s^2+b^2) \right]_s^\infty$$

$$= \frac{1}{2} \left[\lim_{s \rightarrow \infty} (\log(s^2+a^2) - \log(s^2+b^2)) - (\log(s^2+a^2) - \log(s^2+b^2)) \right]$$

$$= \frac{1}{2} \log\left(\frac{s^2+b^2}{s^2+a^2}\right)$$

$$= \frac{1}{2} \log\left(\frac{s^2+b^2}{s^2+a^2}\right)$$

By definition of LT,

$$\int_0^\infty e^{-st} \left(\frac{\cos at - \cos bt}{t}\right) dt = \frac{1}{2} \log\left(\frac{s^2+b^2}{s^2+a^2}\right)$$

Put s=0 o.b.s

$$\int_0^\infty \left(\frac{\cos at - \cos bt}{t}\right) dt = \frac{1}{2} \log\left(\frac{b^2}{a^2}\right)$$

$$= \log\sqrt{\left(\frac{b^2}{a^2}\right)} = \log(b/a)$$

3. ST $\int_0^\infty \left(\frac{\cos 5t - \cos 3t}{t}\right) dt = \log(3/5)$ Note: put a=5, b=3 in above problem

Laplace Transform of Periodic Function:

Definition : A function f(t) is said to be periodic with period T , if

$\forall t, f(t+T) = f(t)$ where T is positive constant.

The least value of $T > 0$ is called the periodic function of f(t).

Example: $\sin t = \sin(2\pi + t) = \sin(4\pi + t) = \dots$ — Here $\sin t$ is periodic function with period 2π .

Formula :- If $f(t)$ is periodic function with period $T \forall t$ then

$$L[f(t)] = \frac{1}{1-e^{-sT}} \int_0^T e^{-st} f(t) dt$$

Problem : Find the L. T of the function $f(t) = e^t$, $0 < t < 5$ and $f(t) = f(t+5)$

$$\begin{aligned} & \frac{1}{1-e^{-s5}} \int_0^5 e^{-st} f(t) dt \\ &= \frac{1}{1-e^{-s5}} \int_0^5 e^{-st} e^t dt \end{aligned}$$

Solution : Here $T=5$ $L[f(t)] = \frac{1}{1-e^{-5s}} \left[\frac{e^{(1-s)t}}{1-s} \right]_0^5 = \frac{1}{1-e^{-5s}} \left[\frac{e^{5(1-s)}}{1-s} \right]$

The unit step function or Heaviside's unit function :

It is denoted by $u(t-a)$ or $H(t-a)$ and is defined as $H(t-a) = 0, t < a$
 $= 1, t > a$ L.T.

of unit step function:

Prove that $L[H(t-a)] = \frac{e^{-as}}{s}$

Solution : $L[H(t-a)] = \int_0^{\infty} e^{-st} H(t-a) dt$

$$= \int_0^a e^{-st} H(t-a) dt + \int_a^{\infty} e^{-st} H(t-a) dt$$

$$= \int_0^a 0 + \int_a^{\infty} e^{-st} \cdot 1 dt$$

$$= \left(\frac{e^{-st}}{-s} \right)$$

$$= \left(\frac{e^{-sa}}{s} \right) \cdot dt$$

Inverse Laplace Transform :

Definition : If $f(s)$ is the Laplace Transform of $f(t)$ then $f(t)$ is called the inverse Laplace Transform of $f(s)$ and is denoted by $L^{-1} f(s)$. i.e., $f(t) = [L^{-1} f(s)]$

$$L^{-1} f(s) = [()]$$

L^{-1} is called inverse Laplace Transform operator, but not reciprocal.

Example : If $L[e^{at}] = \frac{1}{s-a}$ then $e^{at} = L^{-1}[\frac{1}{s-a}]$

Linear Property :

If $f_1(s)$ and $f_2(s)$ are L.T. of $f_1(t)$ and $f_2(t)$ respectively then

$L^{-1}[c_1 f_1(s) + c_2 f_2(s)] = c_1 L^{-1}[f_1(s)] + c_2 L^{-1}[f_2(s)]$ where c_1 , c_2 constants.

Standard Formulae :

$$\frac{1}{s} \Rightarrow L^{-1}\left[\frac{1}{s}\right] = 1$$

$$(2) \quad L[e^{at}] = \frac{1}{s-a} \quad \Rightarrow L^{-1}\left[\frac{1}{s-a}\right] = e^{at}$$

$$(3) \quad L[e^{-at}] = \frac{1}{s+a} \quad \Rightarrow L^{-1}\left[\frac{1}{s+a}\right] = e^{-at}$$

$$(4) \quad L[\sin at] = \frac{a}{s^2+a^2} \Rightarrow L^{-1}\left[\frac{1}{s^2+a^2}\right] = \frac{1}{a} \sin at$$

$$(5) \quad L\left[\cos \frac{s}{s^2+a^2} at\right] \Rightarrow L^{-1}\left[\frac{s}{s^2+a^2}\right] = \cos at$$

$$5) \quad L\left[\sin \frac{a}{s^2-a^2} at\right] \Rightarrow L^{-1}\left[\frac{1}{s^2-a^2}\right] = \frac{1}{a} \sinh at$$

$$6) \quad L\left[\cos \frac{s}{s^2-a^2} at\right] \Rightarrow L^{-1}\left[\frac{s}{s^2-a^2}\right] = \cosh at$$

$$7) \quad L(t^n) = \rho(n+1)/s^{n+1}, \quad n > -1 \Rightarrow L^{-1}\left[\frac{1}{s^{n+1}}\right] = \frac{t^n}{\rho(n+1)}$$

$$8) \quad L(t^n) = n!/s^{n+1}, \quad n \text{ is +ve integer} \Rightarrow L^{-1}\left[\frac{1}{s^{n+1}}\right] = \frac{t^n}{n!} \text{ Problems:}$$

$$(1) \quad L^{-1}\left[\frac{1}{s^2} + \frac{1}{s+4} + \frac{1}{s^2+4} + \frac{s}{s^2-9}\right] \quad \text{Find}$$

solution :

$$L^{-1}\left[\frac{1}{s^2}\right] + L^{-1}\left[\frac{1}{s+4}\right] + L^{-1}\left[\frac{1}{s^2+4}\right] + L^{-1}\left[\frac{s}{s^2-9}\right]$$

$$= t + e^{-4t} + \frac{1}{2} \sin 2t + \cosh 3t.$$

$$L^{-1}\left[\frac{1}{s^2+25}\right]$$

$$L^{-1}\left[\frac{1}{s^2+25}\right] = L^{-1}\left[\frac{1}{s^2+5^2}\right] = \frac{1}{5} \sin 5t$$

$$L^{-1}\left[\frac{1}{2s-5}\right]$$

(2) Find solution

:

(3) Find

solution : $L^{-1} \left[\frac{1}{2s-5} \right] = \frac{1}{2} L^{-1} \left[\frac{1}{s-5/2} \right] = \frac{1}{2} e^{5/2 t}$ solution :

(4) Find $L^{-1} \left[\frac{2s+1}{s(s+1)} \right]$

$$L^{-1} \left[\frac{2s+1}{s(s+1)} \right] = L^{-1} \left[\frac{s+s+1}{s(s+1)} \right] = L^{-1} \left[\frac{1}{s+1} + \frac{1}{s} \right] = e^{-t} + 1$$

(5) Find $L^{-1} \left[\frac{3s-8}{4s^2+25} \right]$] }

solution: $L^{-1} \left[\frac{3s-8}{4s^2+25} \right] = \frac{1}{4} L^{-1} \left[\frac{3s-8}{s^2+(5/2)^2} \right]$
 $= \frac{1}{4} \left\{ 3 L^{-1} \left[\frac{s}{s^2+(5/2)^2} \right] - 8 L^{-1} \left[\frac{1}{s^2+(5/2)^2} \right] \right\}$
 $= \frac{3}{4} \cos \frac{5}{2} t - 8 \times \frac{1}{25} \sin \frac{5}{2} t$
 $= \frac{3}{4} \cos \frac{5}{2} t - \frac{8}{25} \sin \frac{5}{2} t$

Sin $\frac{5}{2} t$

$$= \frac{3}{4} \cos \frac{5}{2} t - \frac{8}{25} \sin \frac{5}{2} t$$

FIRST SHIFTING THEOREM OF INVERSE L.T:

If $L^{-1} [f(s)] = f(t)$ then $L^{-1} [f(s-a)] = e^{at} f(t)$
 $= e^{at} L^{-1} [f(s)]$

PROOF:

By definition of L.T

$$\int_0^{\infty} e^{-st} f(t) dt = f(s) \text{-----(1)}$$

$$L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt$$

$$= \int_0^{\infty} e^{-(s-a)t} f(t) dt$$

$$L[e^{at}$$

$$f(t) dt \text{ Put } s-a=p = \int_0^{\infty} e^{-pt} f(t)$$

dt

$$= f(p) = f(s-a)$$

$$L[e^{at}f(t)] = f(s-a)$$

$$\Rightarrow L^{-1}[f(s-a)] = e^{at} f(t) \quad (\text{or}) \quad L^{-1}[f(s-a)] = e^{at} L^{-1}[f(s)]$$

$$\text{Note: } L^{-1}[f(s+a)] = e^{-at} L^{-1}[f(s)]$$

PROBLEMS

$$L^{-1}\left[\frac{s+3}{(s+3)^2+8^2}\right]$$

1) Find

$$L^{-1}\left[\frac{s+3}{(s+3)^2+8^2}\right] = e^{-3t} L^{-1}\left[\frac{s}{s^2+8^2}\right] \quad \text{by F.S.T}$$

Solution

$$= e^{-3t}$$

Cos 8t.

$$L^{-1}\left[\frac{1}{s^2+2s+5}\right]$$

$$L^{-1}\left[\frac{1}{s^2+2s+5}\right] = L^{-1}\left[\frac{1}{(s+1)^2+4}\right] = e^{-t} L^{-1}\left[\frac{1}{s^2+2^2}\right] = e^{-t}$$

$$L^{-1}\left[\frac{1}{(s+1)^2}\right]$$

$$L^{-1}\left[\frac{1}{(s+1)^2}\right] = L^{-1}\left[\frac{1}{(s+1)^2}\right] = e^{-t} L^{-1}\left[\frac{1}{s^2}\right] = e^{-t} t$$

2) Find

Solution :

$$\frac{1}{2} \sin 2t$$

3) Find

Solution :

4) Find Inverse L.T of $\frac{s}{(s+3)^2}$

$$\begin{aligned} L^{-1}\left[\frac{s}{(s+3)^2}\right] &= L^{-1}\left[\frac{s+3-3}{(s+3)^2}\right] = e^{-3t} L^{-1}\left[\frac{s-3}{s^2}\right] \\ &= e^{-3t} \left\{ L^{-1}\left[\frac{1}{s}\right] - 3 L^{-1}\left[\frac{1}{s^2}\right] \right\} = e^{-3t}(1-3t) \end{aligned}$$

Solution :

$$\begin{aligned} L^{-1}\left[\frac{s+3}{s^2-10s+29}\right] &= L^{-1}\left[\frac{s+3}{(s-5)^2+4}\right] = L^{-1}\left[\frac{(s-5)+5+3}{(s-5)^2+4}\right] \\ &= e^{5t} L^{-1}\left[\frac{s+8}{s^2+4}\right] \\ &= e^{5t} \left\{ L^{-1}\left[\frac{s}{s^2+4}\right] + 8 L^{-1}\left[\frac{1}{s^2+4}\right] \right\} \\ &= e^{5t} \left\{ L^{-1}\left[\frac{s}{s^2+2^2}\right] + 8 L^{-1}\left[\frac{1}{s^2+2^2}\right] \right\} \end{aligned}$$

5) Find

Solution :

]

(By F.S.T)

$$= e^{5t} [\cos 2t + 8 \times \frac{1}{2} \times \sin 2t] \quad a=2$$

$$= e^{5t}$$

SECOND SHIFTING THEOREM: $[\cos 2t + 4 \sin 2t]$

If $L^{-1}[f(s)] = f(t)$ then $L^{-1}[e^{-as} f(s)] = g(t)$ where $g(t) = f(t-a), t > a$
 $= 0, \quad t < a$

Proof: By S.S.T of L.T, $L[g(t)] = e^{-as} f(s)$ (write proof of SST)

$$\Rightarrow L^{-1}[e^{-as} f(s)] = g(t)$$

$$\Rightarrow L^{-1}[e^{-as} f(s)] = f(t-a), t > a$$

$$= 0, \quad t < a \text{ Note:}$$

We can also write as $L^{-1}[e^{-as} f(s)] = f(t-a) H(t-a)$

Problem:

Find $L^{-1}\left[\frac{e^{-\pi s}}{s^2+1}\right]$

$$L^{-1}\left[\frac{e^{-\pi s}}{s^2+1}\right] = L^{-1}\left[e^{-\pi s} \frac{1}{s^2+1}\right]_{\pi s}$$

Solution:

Let $f(s) = \frac{1}{s^2+1}$

$$L^{-1}[f(s)] = L^{-1}\left[\frac{1}{s^2+1}\right] = \sin t = f(t)$$

by S.S.T $L^{-1}[e^{-as} f(s)] = f(t-a), t > a$
 $= 0, t < a$

So $L^{-1}[e^{-\pi s} f(s)] = f(t-\pi), t > \pi$
 $= 0, t < \pi$

$$L^{-1}\left[e^{-\pi s} \frac{1}{s^2+1}\right] = \sin(t-\pi), t > \pi = 0, t < \pi$$

Chang of scale property :

If $L^{-1}[f(s)] = f(t)$ then $L^{-1}\left[f\left(\frac{s}{a}\right)\right] = a f(at)$
 (or) $L^{-1}[f(as)] = \frac{1}{a} f\left(\frac{t}{a}\right)$

Proof : By the change of scale property,

$$L[f(at)] = \frac{1}{a} f\left(\frac{s}{a}\right)$$

$$\Rightarrow L^{-1}\left[f\left(\frac{s}{a}\right)\right] = a f(at)$$

(or)

$$L^{-1}[f(as)] = \frac{1}{a} f\left(\frac{t}{a}\right)$$

Problem(1): If $L^{-1}\left[\frac{s^2-1}{(s^2+1)^2}\right] = t \cos t$, then find $L^{-1}\left[\frac{9s^2-1}{(9s^2+1)^2}\right]$

Solution : Given $L^{-1}\left[\frac{s^2-1}{(s^2+1)^2}\right] = t \cos t$

$$\text{i.e., } L^{-1}[f(s)] = f(t)$$

, Here $f(s) = \frac{s^2-1}{(s^2+1)^2}$ $f(t) = t \cos t$

$$L^{-1}\left[\frac{9s^2-1}{(9s^2+1)^2}\right] \quad \text{Now} \quad] = L^{-1}\left[\frac{(3s)^2-1}{\{(3s)^2+1\}^2}\right]$$

$$= L^{-1}[f(3s)]$$

By change of scale property ,

$$= \frac{1}{3} f\left(\frac{t}{3}\right)$$

$$L^{-1}[f(as)] = \frac{1}{a} f\left(\frac{t}{a}\right) = \frac{1}{3} \frac{t}{3} \cos \frac{t}{3} \quad a = 3$$

Inverse Laplace Transform of partial fractions :

Problems : (1) Find $L^{-1}\left[\frac{(s^2+1)(s-1)}{s^4}\right]$

Solution : Given
$$L^{-1}\left[\frac{(s^2+1)(s-1)}{s^4}\right] = L^{-1}\left[\frac{(s^3-s^2+s-1)}{s^4}\right]$$
$$= L^{-1}\left[\frac{1}{s}\right] - L^{-1}\left[\frac{1}{s^2}\right] + L^{-1}\left[\frac{1}{s^3}\right] - L^{-1}\left[\frac{1}{s^4}\right]$$
$$= 1 - t + \frac{1}{2}t^2 - \frac{t^3}{6}$$

(2). Find $L^{-1}\left[\frac{s+5}{s^2-3s+2}\right] \log\left(\frac{s+3}{s+4}\right)$ Solution : Here $f(s) = \frac{s+5}{s^2-3s+2}$
reduce into partial fractions

$$f(s) = \frac{s+5}{s^2-3s+2} = \frac{s+5}{(s-1)(s-2)} = \frac{A}{s-1} + \frac{B}{s-2} \text{ ---- (1)}$$

$$\Rightarrow s+5 = A(s-2) + B(s-1)$$

put $s=1$ on both sides $\Rightarrow A = -6$

put $s=2$ on both sides $\Rightarrow B = 7$

Therefore (1) $\Rightarrow f(s) = \frac{-6}{s-1} + \frac{7}{s-2}$

$$L^{-1}[f(s)] = L^{-1}\left[\frac{-6}{s-1} + \frac{7}{s-2}\right] = -6e^t + 7e^{2t}$$

Inverse Laplace Transform of derivatives :-

If $L^{-1}[f(s)] = f(t)$ then $L^{-1}\left[\frac{d^n}{ds^n} f(s)\right] = (-1)^n t^n f(t)$

Proof : By theorem of L.T. $L[t^n f(t)]$

$$(-1)^n \frac{d^n}{ds^n} f(s) = L[t^n f(t)]$$

$$\Rightarrow L^{-1}\left[\frac{d^n}{ds^n} f(s)\right] = (-1)^n t^n f(t)$$

Note:- $L^{-1}[f'(s)] = -t f(t)$

Problem (1):- Find

Solution : Let $f(s) =$

$$\log\left(\frac{s+3}{s+4}\right) = \log(s+3) - \log(s+4)$$

$$\frac{1}{s+3} - \frac{1}{s+4}$$

$$L^{-1}[f'(s)] = L^{-1}\left[\frac{1}{s+3} - \frac{1}{s+4}\right]$$

$$= e^{-3t} - e^{-4t}$$

By theorem, $\int_0^t f(t-\tau) f'(\tau) d\tau = \frac{e^{-3t} - e^{-4t}}{-t} e^{-4t}$ H.W. Find $L^{-1}\left[\log\left(\frac{s+1}{s-1}\right)\right]$ SO, $\frac{e^t - e^{-t}}{t}$

$f(t) = \text{Ans: } L^{-1}[f(s)] =$

[replace 3 by -3 and 4 by -4]

$$\Rightarrow L^{-1}[f(s)] = \frac{e^{-4t} - e^{-3t}}{t}$$

(2) Find $L^{-1}\left[\frac{s}{(s^2+a^2)^2}\right]$

Solution: W.K.T $L^{-1}\left[\frac{1}{(s^2+a^2)}\right] = \frac{1}{a} \sin at$

i.e $L^{-1}[f(s)] = f(t)$ 1 Let $f(s) = \frac{1}{(s^2+a^2)}$

$\therefore f(t) = \frac{1}{a} \sin at$

a

We have $L^{-1}[f'(s)] = -t f(t)$

$$L^{-1}\left[\frac{d}{ds}\left(\frac{1}{(s^2+a^2)}\right)\right] = -t \frac{1}{a} \sin at$$

$$L^{-1}\left[\frac{-2s}{(s^2+a^2)^2}\right] = -\frac{t}{a} \sin at$$

$$\Rightarrow L^{-1}\left[\frac{s}{(s^2+a^2)^2}\right] = \frac{t}{2a} \sin at$$

Inverse L.T. of integrals :-

If $L^{-1}[f(s)] = f(t)$ then $L^{-1}\left[\int_s^\infty f(s) ds\right] = \frac{f(t)}{t}$

Proof : We have $L\left[\frac{f(t)}{t}\right] = \int_s^\infty f(s) ds$ provided
exist

$$\Rightarrow L^{-1}\left[\int_s^\infty f(s) ds\right] = \frac{f(t)}{t}$$

Multiplication by powers of s :-

If $L^{-1}[f(s)] = f(t)$ and $f(0) = 0$, then $L^{-1}[s f(s)] = f'(t)$ Proof :

$$\text{W.K.T. } L[f'(t)] = s L[f(t)] - f(0)$$

$$= s f(s) - 0$$

$$\Rightarrow L^{-1}[s f(s)] = f'(t)$$

In general we have, $\Rightarrow L^{-1}[s^n f(s)] = f^n(t)$ if $f^n(0) = 0$

Problems :

$$L^{-1}\left[\frac{s^2}{(s^2+a^2)^2}\right]$$

(1) Find $L^{-1}\left[\frac{s}{(s^2+a^2)^2}\right] = L^{-1}\left[s \cdot \frac{s}{(s^2+a^2)^2}\right]$

2

solution :

s

Let $f(s) =$

$$L^{-1}[f(s)] = \frac{1}{2a} \left[\sin at + t a \cos at \right]$$

$$\text{We have } L^{-1}[s f(s)] = f'(t)$$

$$\Rightarrow L^{-1}\left[\frac{s^2}{(s^2+a^2)^2}\right] = \frac{1}{2a}$$

$$(2) \text{ Find } L^{-1}\left[\frac{s^2}{(s-1)^4}\right] \quad (\sin at + at \cos at)$$

Solution

$$\begin{aligned} : [f(s)] &= \frac{s}{(s-1)^4} \\ &= L^{-1}\left[\frac{s-1+1}{(s-1)^4}\right] \\ &= e^t L^{-1}\left[\frac{s+1}{s^4}\right] \\ &= e^t L^{-1}\left[\frac{1}{s^3} + \frac{1}{s^4}\right] \\ &= e^t \left(\frac{t^2}{2} + \frac{t^3}{6}\right) = f(t) \end{aligned}$$

$$\text{Let } f(s) = L^{-1}$$

] by F.S.T.

$$e^t \left(\frac{t^2}{2} + \frac{t^3}{6} \right) + e^t \left(t + \frac{t^2}{2} \right)$$

Now $f'(t) = e^t \left(t + t^2 + \frac{t^3}{6} \right)$

By theorem $L^{-1}[s f(s)] = f'(t)$

$$L^{-1} \left[s \frac{s}{(s-1)^4} \right] = e^t \left(t + t^2 + \frac{t^3}{6} \right) \text{ Division}$$

by power of S :

Theorem: If $L^{-1} f(s) = f(t)$ then $L^{-1} [s f(s)] = t f(t)$

Prof: we have by LT,

$$\int_0^t f(t) dt = \frac{f(s)}{s}$$

L[

$$\Rightarrow L^{-1} \left[\frac{f(s)}{s} \right] = \int_0^t f(t) dt$$

$$L^{-1} \left[\frac{f(s)}{s^2} \right] = \int_0^t \int_0^t f(t) dt dt$$

$$L^{-1} \left[\frac{f(s)}{s^2} \right] = \int_0^t \int_0^t f(t) dt dt$$

Note: ()

Problem:

1) Find $L^{-1} \left[\frac{1}{s(s+3)} \right]$

solution: Let $f(s) = \frac{1}{s+3}$

$$L^{-1} [f(s)] = L^{-1} \left[\frac{1}{s+3} \right] = e^{-3t} = f(t)$$

By theorem, $L^{-1} \left[\frac{1}{s} \cdot f(s) \right] = \int_0^t f(t) dt$

$$\Rightarrow L^{-1} \left[\frac{1}{s(s+3)} \right] = \int_0^t e^{-3t} dt = \left[\frac{e^{-3t}}{-3} \right]_0^t = \frac{1 - e^{-3t}}{3}$$

2) Find $L^{-1} \left[\frac{1}{s(s^2+a^2)} \right]$

Solution : let $f(s) = \frac{1}{s^2+a^2}$, $L^{-1} \left[\frac{1}{s^2+a^2} \right] = f(s) = \sin at = f(t)$

By
theorem

$$L^{-1} \left[\frac{1}{s} f(s) \right] = \int_0^t f(t) dt$$

$$\Rightarrow L^{-1} \left[\frac{1}{s(s^2+a^2)} \right] = \int_0^t \frac{1}{a} \sin at = \frac{1}{a} \left(1 - \frac{\cos at}{a} \right)$$

$$= \frac{1}{a^2} (1 - \cos at)$$

3) Find $L^{-1} \left[\frac{1}{s^2(s^2+a^2)} \right]$

1 1 solution : let f(s)

= , f(t) = sin at

a

theorem, $L^{-1} \left[\frac{1}{s^2} f(s) \right] =$

$$\int_0^t \int_0^t f(t) dt = \int_0^t \left[\int_0^t a \sin at dt \right] dt$$

$$= \int_0^t \frac{1}{a^2} (1 - \cos at) dt = \frac{1}{a^2} \left(t - \frac{\sin at}{a} \right)$$

Convolution :-

If $f(t)$ and $g(t)$ are two functions defined for $t \geq 0$, then the convolution of $f(t)$ and $g(t)$ is defined as , $f(t) * g(t) = \int_0^t f(u) g(t-u) du$.

$f(t) * g(t)$ can also be written as $(f * g)(t)$. Note:- The convolution operation is commutation

i.e. , $(f * g)(t) = (g * f)(t)$

$$\Rightarrow \int_0^t f(u) g(t-u) du = \int_0^t f(t-u) g(u) du$$

Convolution theorem :-

If $L[f(t)] = f(s)$ and $L[g(t)] = g(s)$ then $L[f(t) * g(t)] = L[f(t)] \cdot L[g(t)]$

(or)

$$= f(s) \cdot g(s)$$

So, $L[(f * g)(t)] = f(s) \cdot g(s)$

Corollary :- $L^{-1}[f(s) \cdot g(s)] = (f * g)(t)$

$$= \int_0^t f(u) g(t-u) du$$

$$= \int_0^t f(t-u) g(u) du.$$

Problems:

(1). Find $L^{-1}\left[\frac{1}{(s-2)(s^2+1)}\right]$ by using convolution theorem.

11 solution: Let $f(s) =$

$$\frac{1}{s-2}, \quad g(s) = \frac{1}{s^2+1}$$

$$L^{-1}[f(s)] = L^{-1}\left[\frac{1}{s-2}\right] = e^{2t}, \quad L^{-1}[g(s)] = L^{-1}\left[\frac{1}{s^2+1}\right] = \sin t$$

By convolution theorem,

$$L^{-1}[f(s) \cdot g(s)] = \int_0^t f(t-u) g(u) du$$

$$\Rightarrow L^{-1}\left[\frac{1}{(s-2)(s^2+1)}\right] = \int_0^t e^{2(t-u)} \sin u du$$

$$= e^{2t} \int_0^t e^{-2u} \sin u du$$

$$= e^{2t} \left[\frac{e^{-2u}}{(-2)^2+1^2} (-\sin u - \cos u) \right]_0^t$$

$\sin u -$

$\cos u)$

$$= e^{2t} \left[\frac{e^{-2t}}{5} (-2 \sin t - \cos t) - \frac{e^0}{5} (-1) \right]$$

$$= e^{2t} \left[\frac{e^{-2t}}{5} (-2 \sin t - \cos t) + \frac{1}{5} \right]$$

$$= \frac{1}{5} (-2 \sin t - \cos t) + \frac{1}{5} e^{2t}$$

$$= \frac{1}{5} [e^{2t} - 2 \sin t - \cos t]$$

2) Find $L^{-1}[\frac{1}{s(s^2-a^2)}]$ by convolution theorem

Solution : Let $f(s) = \frac{1}{s}$, $g(s) = \frac{1}{s^2-a^2}$

$$L^{-1}[f(s)] = L^{-1}\left[\frac{1}{s}\right] = 1 = f(t), \quad L^{-1}[g(s)] = L^{-1}\left[\frac{1}{s^2-a^2}\right] = \frac{1}{a} \sinh at = g(t) \text{ By convolution theorem,}$$

$$\begin{aligned} L^{-1}[f(s) \cdot g(s)] &= \int_0^t f(t-u) g(u) du \\ \Rightarrow L^{-1}\left[\frac{1}{s(s^2-a^2)}\right] &= \int_0^t 1 \cdot \frac{1}{a} \sinh au \, du \\ &= \frac{1}{a} \left[\frac{\cosh au}{a} \right], \text{ (apply limits 0 to t)} \\ &= \frac{1}{a^2} (\cosh at - 1) \end{aligned}$$

Application of L . T to Ordinary Differential Equations :

The L . T method is easier , time – saving and excellent tool for solving O.D.Es

Working rule for finding solution of D . E by L . T:

- 1) Write down the given equation and apply L . T O . B . S**
- 2) Use the given conditions**
- 3) Rearrange the given equation to given transformation of the solution**
- 4) Take inverse L.T O. B. S to obtain the desired observe Satisfying the given conditions**

The formulae to be used in this process are:

$$L [f' (t)] = s f (s) - f(0)$$

$$L [f'' (t)] = s^2 f (s) - s f(0) - f'(0)$$

$$L [f''' (t)] = s^3 f (s) - s^2 f(0) - s f'(0) - f''(0)$$

Note : let $f(t) = y(t)$ and $f(s) = y(s)$ Problems :

1) Solve $4 y'' + \pi^2 y = 0$, $y(0) = 2$, $y'(0) = 0$

Solution : Here $y = y(t)$

Given D . E $4 y^{11}(t) + \pi^2 y(t) = 0$ **Let L . T O.B.S**

$$\Rightarrow 4 [s^2 L(y)] - s y(0) - y'(0) + \pi^2 L[y] = L[0]^2$$

$$\Rightarrow L[y] [4s^2 + \pi^2] - L[y] = 0$$

$$\Rightarrow L[y] = \frac{8s}{4s^2 + \pi^2} \quad 4s(2) - 0 = 0$$

Let L^{-1} O . B . S, we get

$$y(t) = L^{-1} \left[\frac{s}{4(s^2 + \pi^2/4)} \right] = 8$$

$$= \frac{8}{4} L^{-1} \left[\frac{s}{s^2 + (\pi^2/2)^2} \right]$$

$$] = 2 \cdot \cos \pi/2t$$

$$\Rightarrow y(t) = 2 \cdot \cos \pi/2t \quad \text{is solution of}$$

gven D.E

3) Solve $y^{11} + 2y^{11} - y^1 - 2y = 0$ with $y(0) = y'(0) = 0$, $y^{11}(0) = 6$

Solution : given D . E

Let $L.T$ On Both Sides

$$L[y^{11}] + 2 L[y^{11}] - L[y^1] - 2 L[y] = 0$$

$$y^1(0)]$$

$$- s L[y] - y(0) - 2 L[y] = 0$$

$$\Rightarrow L[y] (s^3 + 2s^2 - s - 2) - 6 = 0$$

$$\Rightarrow L[y] = \frac{6}{s^3 + 2s^2 - s - 2}$$

$$\Rightarrow s^3 L[y] s^2 y(0) s y^1(0) y^{11}(0) + 2 [s^2 L[y]$$

$$L[y] = \frac{6}{(s-1)(s+1)(s+2)} = \frac{A}{s-1} + \frac{B}{s+1} + \frac{C}{s+2} \quad (1)$$

$$6 = A(s+1)(s+2) + B(s-1)(s+2) + C(s-1)(s+1)$$

$$(2) \text{ Put } s = 1 \text{ in } (2) \quad 6 = A(2)(3) \Rightarrow A = 1$$

$$\text{Put } s = -1 \text{ in } (2)$$

$$\Rightarrow 6 = B(-2)(1) \Rightarrow B = -3$$

$$\text{Put } s = -2 \text{ in } (2)$$

$$\Rightarrow 6 = C(-3)(-1) \Rightarrow C = 2$$

Substitute A , B , C in (1)

$$\Rightarrow L[y] = \frac{1}{s-1} - \frac{3}{s+1} + \frac{2}{s+2}$$

$$\Rightarrow y = L^{-1} \left[\frac{1}{s-1} - \frac{3}{s+1} + \frac{2}{s+2} \right]$$

$$\Rightarrow y(t) = e^t - 3e^{-t} + 2e^{-2t}$$

is the solution of given D . E

HW: Solve the D.E $\frac{d^2y}{dt^2} + 2\frac{dy}{dt} + 5y = e^{-t} \sin t$

$$\text{Ans: } y(t) = \frac{e^{-t}}{3} (\sin t - 2 \sin 2t)$$

UNIT – IV

FOURIER SERIES

Periodic Function :

Definition : A function $f(x)$ is said to be periodic with period T , if \forall
 x , $f(x+T) = f(x)$ where T is positive constant.

The least value of $T > 0$ is called the periodic function of $f(x)$.

Example: $\sin x = \sin(2\pi + x) = \sin(4\pi + x) = \dots$

Here $\sin x$ is periodic function with period 2π . **Def:**

Piecewise Continuous Function:

A function is said to be piece-wise continuous (or) Sectionally Continuous) over the closed interval $[a,b]$ if it is defined on that interval and is such that the interval can be divided into a finite number of sub intervals, in each of which $f(x)$ is continuous and both right and left hand limits at every end point of the sub intervals.

Dirichlet Conditions:

A function $f(x)$ satisfies Dirichlet conditions if

- (1) $f(x)$ is well defined and single valued except at a finite no. of points in $(-l,l)$

(2) $f(x)$ is periodic function with period $2l$

(3) $f(x)$ and $f'(x)$ are piece wise continuous in $(-l, l)$

Fourier Series: If $f(x)$ satisfies Dirichlet conditions, then it can be represented by an infinite series called Fourier Series in an interval $(-l, l)$ as

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \text{-----}$$

$$a_0 = \frac{1}{l} \int_{-l}^l f(x) dx, \quad a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx$$

$$b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx$$

(1) where

Here a_0 , a_n and b_n are called Fourier coefficients.

These are also

called Euler's formula.) (i.e., interval is $(-\pi, \pi)$)

Note (1): If $x \in (-\pi, \pi)$ $\frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$

$$\text{Then } f(x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx, \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

Where $a_0 =$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx$$

Note (2): In interval $(0, 2\pi)$, $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$

$$\text{Where } a_0 = \frac{1}{\pi} \int_0^{2\pi} f(x) \, dx, \quad a_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx$$

Note (3): The Fourier Series in $(-l, l)$, $(-\pi, \pi)$, $(c, c + 2\pi)$ are called Full range expansion series

Note (4): The above series (1) converges to $f(x)$ if x is a point of continuity

The above series (1) converges to $\frac{f(x+0) + f(x-0)}{2}$ if x is a point of discontinuity

$$f(\pi-0) + f(-\pi+0)$$

Note (5): At $x = \pm\pi$, $f(x) = \frac{f(\pi-0) + f(-\pi+0)}{2}$ here $x \in (-\pi, \pi)$

Even and odd functions:

Case (1): If the function $f(x)$ is an even function in the interval $(-l, l)$

$$\text{i.e., } f(-x) = f(x) \text{ then } a_0 = \frac{2}{l} \int_0^l f(x) \, dx$$

$$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx \quad (\text{since } f(x) \text{ \& } \cos \frac{n\pi x}{l} \text{ are even functions})$$

$$b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx \Rightarrow b_n = 0 \quad (\text{since } f(x) \cdot \sin \frac{n\pi x}{l} \text{ is odd function})$$

Therefore, in this case we get (only) Fourier cosine series only.

Case (2): If function $f(x)$ is odd i.e., $f(-x) = -f(x)$ then

$$a_n = 0 \quad (\text{since } f(x) \cos \frac{n\pi x}{l} \text{ is odd}) \quad (a_0 = 0 \text{ also})$$

$$\text{And } b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

In this case we get Fourier sine series only.

[only for intervals $(-l, l)$, $(-\pi, \pi)$]

Problems

:

1) Find Fourier series for the function $f(x) = e^{ax}$ in $(0, 2\pi)$ Solution : Given

function $f(x) = e^{ax}$ in $(0, 2\pi)$

$$\frac{1}{\pi} \int_0^{2\pi} f(x) dx = \frac{1}{\pi} \int_0^{2\pi} e^{ax} dx = \frac{1}{\pi} \left(\frac{e^{ax}}{a} \right)$$

$ax \text{ } a_0 =) \text{ apply limits } 0$

to 2π

$$= \frac{1}{a\pi} (e^{2\pi a} - 1)$$

$$= \frac{1}{\pi} \int_0^{2\pi} e^{ax} \cos nx \, dx$$

an

=

$$= \frac{1}{\pi} \left[\frac{e^{ax}}{a^2 + n^2} (a \cos nx + n \sin nx) \right]$$

apply limits 0 to 2π

$$= \frac{1}{\pi} \left[\frac{e^{2\pi a}}{a^2 + n^2} (a \cos 2n\pi + 0) - \frac{e^0}{a^2 + n^2} \right]$$

$$= \frac{1}{\pi} \frac{1}{a^2 + n^2} [e^{2\pi a} a - 1 \cdot a]$$

$$= \frac{a}{\pi(a^2 + n^2)} (e^{2\pi a} - 1)$$

$$= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx \, dx$$

$$= \frac{1}{\pi} \int_0^{2\pi} e^{ax} \sin nx \, dx$$

$$= \frac{1}{\pi} \left[\frac{e^{ax}}{a^2 + n^2} (a \sin nx + n \cos nx) \right] \quad \text{apply limits 0 to } 2\pi$$

$$= \frac{1}{\pi} \left[\frac{e^{2\pi a}}{a^2 + n^2} (0 - n \cos 2n\pi) - \frac{e^0}{a^2 + n^2} (0 - n) \right]$$

$$= \frac{1}{\pi} \frac{n}{a^2 + n^2} (1 - e^{2\pi a}) = \frac{-n}{\pi(a^2 + n^2)} (e^{2\pi a} - 1)$$

$$\frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx \, dx$$

(a + 0)] apply limits 0 to 2π

b_n

Now the fourier series is $f(x) =$

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx$$

$$= \frac{\frac{1}{a\pi} (e^{2\pi a} - 1)}{2} + \sum_{n=1}^{\infty} \frac{a}{\pi(a^2 + n^2)} (e^{2\pi a} - 1)$$

$$\frac{-n}{\pi(a^2 + n^2)} (e^{2\pi a} - 1) \sin nx \quad \cos nx + \sum_{n=1}^{\infty}$$

(2): Find Fourier series for the function $f(x) = e^x$ in $(0, 2\pi)$

Solution : Given function $f(x) = e^x$ in $(0, 2\pi)$ $a_0 =$

apply $\frac{1}{\pi} \int_0^{2\pi} f(x) dx = \frac{1}{\pi} \int_0^{2\pi} e^x dx$ limits 0 to 2π

$$= \frac{1}{\pi} (e^x)$$

apply limits 0 to 2π

$$= \frac{1}{\pi} (e^{2\pi} - 1)$$

an = $\frac{1}{\pi} \int_0^{2\pi} f(x) \cos nx dx$

$$= \frac{1}{\pi} \int_0^{2\pi} e^x \cos nx dx$$

bn

$$= \frac{1}{\pi} \left[\frac{e^x}{1+n^2} (1 \cos nx + n \sin nx) \right]$$

$$= \frac{1}{\pi} \left[\frac{e^{2\pi}}{1+n^2} (\cos 2n\pi + 0) - \frac{e^0}{1+n^2} (\cos 0 + 0) \right]$$

$$= \frac{1}{\pi} \frac{1}{1+n^2} [e^{2\pi} - 1]$$

$$= \frac{1}{\pi(1+n^2)} (e^{2\pi} - 1)$$

$= \frac{1}{\pi} \int_0^{2\pi} f(x) \sin nx dx$

$$= \frac{1}{\pi} \int_0^{2\pi} e^x \sin nx dx$$

$$= \frac{1}{\pi} \left[\frac{e^x}{1+n^2} (\sin nx + n \cos nx) \right]$$

apply limits 0 to 2π

$$= \frac{1}{\pi} \left[\frac{e^{2\pi}}{1+n^2} (0 - n \cos 2n\pi) - \frac{e^0}{1+n^2} (0 - n) \right]$$

$$= \frac{1}{\pi} \frac{n}{1+n^2} (1 - e^{2\pi}) = \frac{-n}{\pi(1+n^2)} (e^{2\pi} - 1)$$

Now the fourier series is $f(x) =$

$$\begin{aligned} & \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + \sum_{n=1}^{\infty} b_n \sin nx \\ &= \frac{\frac{1}{\pi}(e^{2\pi}-1)}{2} + \sum_{n=1}^{\infty} \frac{1}{\pi(1+n^2)} (e^{2\pi}-1) \cos nx + \sum_{n=1}^{\infty} \frac{-n}{\pi(1+n^2)} (e^{2\pi}-1) \sin nx \end{aligned}$$

Problem (3): H.W

Find Fourier series for the function $f(x) = e^{-x}$ in $(0, 2\pi)$

(Hint:- put $a = -1$ in problem (1) we get the solution.)

(4) Express $f(x) = x - \pi$ as Fourier Series in the interval $-\pi < x < \pi$ Solution:

Given function $f(x) = x - \pi$ a₀

$$\begin{aligned} &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_{-\pi}^{\pi} (x - \pi) dx \\ &= \frac{1}{\pi} \int_{-\pi}^{\pi} x dx - \frac{1}{\pi} \int_{-\pi}^{\pi} \pi dx \end{aligned}$$

$= 0 - [x]$ with limits $-\pi$ to π

$$= 0 - [\pi + \pi] = -2\pi \quad \text{an} \quad =$$

$$\begin{aligned}
 \text{even)} \quad \frac{dx}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx &= \frac{1}{\pi} \int_{-\pi}^{\pi} (x - \pi) \cos nx \, dx && \text{(since)} \\
 &= \frac{1}{\pi} \int_{-\pi}^{\pi} x \cos nx \, dx - \frac{1}{\pi} (-\pi) \int_{-\pi}^{\pi} \cos nx \, dx \\
 &= \frac{1}{\pi} (0) \text{ (since } x \cos nx \text{ is odd)} + 2 \int_0^{\pi} \cos nx \\
 &= 0 + 2 \left[\frac{\sin nx}{n} \right]_0^{\pi} \text{ 0 to } \pi \text{ limits apply we get an} = \\
 &0+0 = 0
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \int_{-\pi}^{\pi} (x - \pi) \sin nx \, dx \\
 &= \frac{1}{\pi} \int_{-\pi}^{\pi} x \sin nx \, dx - \frac{1}{\pi} (-\pi) \int_{-\pi}^{\pi} \sin nx \, dx
 \end{aligned}$$

$$\begin{aligned}
 &\text{(even)} && \text{(odd)} \\
 &= \frac{1}{\pi} 2 \int_0^{\pi} x \sin nx && dx - 0 \text{ (since } \sin nx \text{ is odd)} \\
 &= \frac{2}{\pi} \left[\left\{ x \left(-\frac{\cos nx}{n} \right) \right\} - \int_0^{\pi} \frac{-\cos nx}{n} dx \right] \\
 &= \frac{2}{\pi} \left[-\pi \frac{\cos n\pi}{n} + 0 + \frac{1}{n} \left(\frac{\sin nx}{n} \right) \right] && \text{apply limits 0 to } \pi \\
 &= \frac{2}{\pi} \left[-\pi \frac{\cos n\pi}{n} + 0 + \frac{1}{n} (0) \right] = -\frac{2}{n} \cos n\pi = -\frac{2}{n} (-1)^n = \frac{2}{n} (-1)^{n+1}, n=1,2,3,\dots
 \end{aligned}$$

Now the Fourier Series of $f(x)$ is $f(x)$

$$\begin{aligned}
&= \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)_{f(x)} \\
&= \frac{2\pi}{2} + \sum_{n=1}^{\infty} \left[(0) \cos nx + \frac{2}{n} (-1)^{n+1} \sin nx \right] \\
&= \pi + \sum_{n=1}^{\infty} \left[\frac{2}{n} (-1)^{n+1} \sin nx \right]
\end{aligned}$$

(5) Obtain the Fourier series for $f(x) = x - x^2$ in the interval $[-\pi, \pi]$

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = \frac{\pi^2}{12}$$

Hence show

that (or)

$$\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots = \frac{\pi^2}{12}$$

Solution : Given function is $f(x) = x - x^2$ in $[-\pi, \pi]$

$$\begin{aligned}
a_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_{-\pi}^{\pi} (x - x^2) dx \\
&= \frac{1}{\pi} \int_{-\pi}^{\pi} x dx - \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 dx \\
&= 0 \text{ (odd)} - \frac{1}{\pi} \left[\frac{x^3}{3} \right]_{-\pi}^{\pi} = -\frac{2\pi^2}{3}
\end{aligned}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx$$

$$= \frac{1}{\pi} \int_{-\pi}^{\pi} (x - x^2) \cos nx \, dx$$

$$= \frac{1}{\pi} \int_{-\pi}^{\pi} x \cos nx \, dx - \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \cos nx \, dx$$

(odd)

(even)

$$= 0 - \frac{1}{\pi} 2 \int_0^{\pi} x^2 \cos nx \, dx \quad x^2, \quad dv = \cos nx \, dx$$

$$= -\frac{2}{\pi} \left[\left(\frac{x^2 \sin nx}{n} \right) - \frac{2}{n} \int_0^{\pi} x \sin nx \, dx \right]$$

$$du = 2x \, dx, \quad dv = -\cos nx \, dx$$

apply limits 0 to π

$$= -\frac{2}{\pi} \left[0 - \frac{2}{n} \left(-x \cos nx \right) + 0 \right] \cos nx = \frac{4}{n} \sin nx \, dx$$

apply limits 0 to π

$$= \frac{4}{\pi n} \left[-\pi \frac{(-1)^n}{n} + \frac{1}{n^2} \right]$$

$$= \frac{4}{n^2} (-1)^{n+1}$$

$$a_n = \frac{4}{n^2} (-1)^{n+1} \quad (\sin nx) \quad \int u dv = uv - \int v du$$

$$a_1 = \frac{4}{1^2} = 4 \quad \text{if } n \text{ is odd}$$

$$n^2$$

$$-\frac{4}{n^2} \text{ if } n \text{ is even}$$

$$a_2 = \frac{4}{2^2} = 1$$

$$a_3 = \frac{4}{3^2} = 4/9$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} (x - x^2) \cos nx \, dx$$

$$= \frac{1}{\pi} \left[\int_{-\pi}^{\pi} x \sin nx \, dx - \int_{-\pi}^{\pi} x^2 \sin nx \, dx \right]$$

$$= \frac{2}{\pi} \left[\left(\frac{-x \cos nx}{n} \right) + \frac{1}{n} \int_0^{\pi} \cos nx \, dx \right]$$

(even) (odd)

$$= \frac{2}{n} \left[-\pi \frac{(-1)^n}{n} + \frac{1}{n^2} \right]$$

$$\sin nx \,] \, b_1 = 2/1 = 2 = \frac{2}{n} (-1)^{n+1} = \frac{2}{n} \text{ if } n \text{ is odd}$$

$$b_2$$

$$= -$$

$$2/2 = -1$$

$$b_3 = 2/3$$

$$= -\frac{2}{n} \text{ if } n \text{ is even}$$

Now

$$= \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) \text{-----(1) substitute}$$

, f(x)

$$\Rightarrow f(x) = \frac{-\pi^2}{3} + 4 \left(\frac{\cos x}{1^2} - \frac{\cos 2x}{2^2} + \frac{\cos 3x}{3^2} - \dots \dots \right)$$

in

$$+ 2 \left(\frac{\sin x}{1} - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} + \dots \dots \right) \text{-----(2)}$$

(1)

put $x = 0$ in (2)

$$f(0) = 0 = \frac{-\pi^2}{3} + 4\left(\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots\right)$$

$$\Rightarrow \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \dots = \frac{\pi^2}{12}$$

Half range series

(1) The half range cosine series in $(0, l)$ is $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l}$

$$a_0 = \frac{2}{l} \int_0^l f(x) dx, \quad a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx$$

(2) The half range sine series in $(0, l)$ is $f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l}$

$$\text{where } b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

Note :1) The half range cosine series in $(0, \pi)$ is $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx$

$$a_0 = \frac{2}{\pi} \int_0^{\pi} f(x) dx, \quad a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos n\pi dx$$

where

Note :2) The half range sine series in $(0, \pi)$ is $f(x) = \sum_{n=1}^{\infty} b_n \sin nx$ where

$$b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx dx$$

(1) Express $f(x) = \pi - x$ as Fourier cosine and sine series in $(0, \pi)$

Solution :

The half range cosine series for $f(x)$ is $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx$ (1)

$$\begin{aligned} \text{where } a_0 &= \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} \pi - x dx \\ &= \frac{2}{\pi} \left[\pi x - \frac{x^2}{2} \right] \text{ apply limits 0 to } \pi \\ &= \frac{2}{\pi} \left[\pi^2 - \frac{\pi^2}{2} - (0-0) \right] = \frac{2}{\pi} \left(\frac{\pi^2}{2} \right) = \pi \\ a_n &= \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx \\ &= \frac{2}{\pi} \int_0^{\pi} (\pi - x) \cos nx dx \\ &= \frac{2}{\pi} \left[\left\{ (\pi - x) \frac{\sin nx}{n} \right\} + \int_0^{\pi} \frac{\sin nx}{n} dx \right] \end{aligned}$$

$$\begin{aligned} &\quad \text{(apply 0 to } \pi) \\ &= \frac{2}{\pi} \left[(0-0) + \frac{1}{n} \left(-\frac{\cos nx}{n} \right) \right] \\ &= -\frac{2}{\pi n^2} [\cos n\pi - \cos 0] \\ &= -\frac{2}{\pi n^2} [\cos n\pi - \cos 0] \quad \text{apply 0 to } \pi \\ &= -\frac{2}{\pi n^2} [(-1)^n - 1] = \frac{2}{\pi n^2} [1 - (-1)^n] \end{aligned}$$

$$\text{Now (1)} \Rightarrow \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{2}{\pi n^2} [1 - (-1)^n] \cos nx: f(x) =$$

H.W.) Express $f(x) = \pi - x$ as fourier sine series in $(0, \pi)$ Ans : $2 \sum_{n=1}^{\infty} \frac{\sin nx}{n}$ ($b_n = \frac{2}{n}$)

2) Find the half range sine series of $f(x) = x$ in the range $0 < x < \pi$

π^2

Hence deduce that $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots =$ 8

Solution : The half range cosine series for $f(x)$ is $f(x)$

$$= \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx \dots\dots\dots(1)$$

where $a_0 = \frac{2}{\pi} \int_0^{\pi} f(x) dx = \frac{2}{\pi} \int_0^{\pi} x dx = \frac{2}{\pi} \left[\frac{x^2}{2} \right] \text{ apply limits 0 to } \pi$

$$= \pi$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \cos nx dx$$

$$= \frac{2}{\pi} \int_0^{\pi} (x) \cos nx dx$$

$$= \frac{2}{\pi} \left[\left\{ (x) \frac{\sin nx}{n} \right\} - \int_0^{\pi} \frac{\sin nx}{n} dx \right]$$

(apply 0 to π)

$$= \frac{2}{\pi} \left[(0-0) - \frac{1}{n} \left(-\frac{\cos nx}{n} \right) \right]$$

$$= \frac{2}{\pi n^2} [\cos n\pi - \cos 0]$$

$$= \frac{2}{\pi n^2} [(-1)^n - 1] \text{ apply 0 to } \pi$$

$a_n = 0$ if n is even

$$= -\frac{4}{\pi n^2}$$

if n is odd

Now

(1)

$$\Rightarrow: f(x) = \frac{\pi}{2} + \sum_{n=1}^{\infty} \frac{-4}{\pi n^2} \cos nx \quad \text{if } n \text{ is odd}$$

$$x = \frac{\pi}{2} - \frac{4}{\pi} \left(\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} - \dots \right)$$

Put x=0 on both sides

$$\Rightarrow 0 = \frac{\pi}{2} - \frac{4}{\pi} \left(1^2 + \frac{1}{3^2} + \frac{1}{5^2} - \dots \right)$$

$$\Rightarrow \frac{4}{\pi} \left(1^2 + \frac{1}{3^2} + \frac{1}{5^2} - \dots \right) = \frac{\pi}{2}$$

$$\Rightarrow \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$$

3) Express $f(x) = \cos x$, $0 < x < \pi$ in half range sine series

π

$$\sum_{n=1}^{\infty} bn \sin nx \text{ -----(1)}$$

$$= \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx \, dx$$

$$= \frac{2}{\pi} \int_0^{\pi} \cos x \sin nx \, dx$$

$$= \frac{2}{\pi} \frac{1}{2} \int_0^{\pi} [\sin(n+1)x + \sin(n-1)x] \, dx$$

$$= \frac{1}{\pi} \left[\frac{-\cos(n+1)x}{n+1} - \frac{\cos(n-1)x}{n-1} \right] \text{ apply limits 0 to } \pi$$

$$= \frac{1}{\pi} \left[\frac{-\cos(n+1)\pi}{n+1} - \frac{\cos(n-1)\pi}{n-1} + \frac{1}{n+1} + \frac{1}{n-1} \right]$$

$$= \frac{1}{\pi} \left[\frac{-(-1)^{n+1}}{n+1} + \frac{(-1)^n}{n-1} + \frac{1}{n+1} + \frac{1}{n-1} \right]$$

$$= \frac{1}{\pi} \left[\left(\frac{(-1)^2(-1)^n}{n+1} + \frac{(-1)^n}{n-1} + \frac{1}{n+1} + \frac{1}{n-1} \right) \right]$$

$$= \frac{1}{\pi} [(-1)^n \left\{ \frac{1}{n+1} + \frac{1}{n-1} \right\} + \left\{ \frac{1}{n+1} + \frac{1}{n-1} \right\}]$$

$$= \frac{1}{\pi} \left[\{(-1)^{n+1}\} \left(\frac{1}{n+1} + \frac{1}{n-1} \right) \right]$$

$$= \frac{2n}{\pi} \left[\frac{1+(-1)^n}{n^2-1} \right] \text{ (n not equal to 1)}$$

Solution : The half range sine series in (0,) is $f(x) =$

where

$$b_n$$

$$]$$

$$], n \text{ is not equal to } 1$$

$$b_n = 0 \text{ if } n \text{ is odd.}$$

$$= \frac{4n}{\pi(n^2-1)} \text{ if } n \text{ is even}$$

$$b_1 = b_3 = b_5 = \dots = 0$$

$$(1) \Rightarrow f(x) = \sum_{n=2}^{\infty} \frac{4n}{\pi(n^2-1)} \sin nx, \text{ for } n \text{ is even}$$

$$4)\text{Find half range sine series for } f(x) = x(\pi - x), \text{ in } 0 < x < \pi$$

$$\frac{1}{1^3} - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} \dots \pi^3$$

$$\text{Deduce that } +\dots =$$

$$32$$

Solution : Fourier series is $f(x) = \sum_{n=1}^{\infty} b_n \sin nx \dots (1)$

$$\begin{aligned} & \frac{2}{\pi} \int_0^{\pi} f(x) \sin nx \, dx \\ &= \frac{2}{\pi} \int_0^{\pi} x(\pi - x) \sin nx \, dx \\ &= \frac{2}{\pi} \pi \int_0^{\pi} x \sin nx \, dx - \frac{2}{\pi} \int_0^{\pi} x^2 \sin nx \, dx \\ &= 2 \left[\left(\frac{-x \cos nx}{n} \right) - \int_0^{\pi} \frac{-\cos nx}{n} \, dx \right] - \frac{2}{\pi} \left[\left(\frac{-x^2 \cos nx}{n} \right) - \int_0^{\pi} \frac{-\cos nx}{n} 2x \, dx \right] \end{aligned}$$

(apply

0 to π)
$$= 2 \left[\left(\frac{-\pi \cos n\pi}{n} \right) + 0 + \frac{1}{n} \left(\frac{\sin nx}{n} \right) 0 \text{ to } \pi \right] - \frac{2}{\pi} \left[\left(\frac{-\pi^2 \cos n\pi}{n} \right) + 0 + \frac{2}{n} \int_0^{\pi} x \cos nx \, dx \right]$$

(apply

0 to π)
$$= 2 \left[-\pi \frac{(-1)^n}{n} + 0 \right] + \frac{2}{\pi} \cdot \pi^2 \frac{(-1)^n}{n} - \frac{4}{\pi n} \left[\left(\frac{x \sin nx}{n} \right) 0 \text{ to } \pi - \int_0^{\pi} \frac{\sin nx}{n} \, dx \right]$$

$$= 2 \left[-\pi \frac{(-1)^n}{n} \right] + 2\pi \frac{(-1)^n}{n} + \frac{4}{\pi n^2} \left(\frac{-\cos nx}{n} \right)$$

$$= \frac{4}{\pi n^3} [-\cos n\pi + \cos 0] \quad) 0 \text{ to } \pi$$

$$= \frac{4}{\pi n^3} [1 - (-1)^n] \quad \text{sub in (1)}$$

b_n

$$(1) \Rightarrow f(x) = \sum_{n=1}^{\infty} \frac{4}{\pi n^3} [1 - (-1)^n] \sin nx$$

$$(1) \Rightarrow f(x) = b_1 \sin x + b_2 \sin 2x + b_3 \sin 3x + \dots$$

$$= \frac{4}{\pi} (2) \sin x + 0 + \frac{4}{\pi \cdot 3^3}$$

$$\Rightarrow x(\pi - x) = \frac{8}{\pi} \left[\frac{\sin x}{1^3} + \frac{\sin 3x}{3^3} + \dots \right] (2) \sin 3x + \dots \text{ Put}$$

$x = \pi/2$ on both sides

$$\frac{\pi \pi}{2} \left[1 - \frac{1}{3^3} + \dots \right] \Rightarrow$$

$$(2) =$$

$$\Rightarrow \frac{\pi^2}{4} = \frac{8}{\pi} \left[\frac{1}{1^3} - \frac{1}{3^3} + \frac{1}{5^3} - \dots \right]$$

$$\Rightarrow \left[\frac{1}{1^3} - \frac{1}{3^3} + \frac{1}{5^3} - \dots \right] = \frac{\pi^2}{32}$$

- **FOURIER SERIES IN AN ARBITRARY INTERVAL I.e in $(-l, l)$ & $(0, 2l)$**

- **Problem : 1) Obtain the half range sine series for e^x in $0 < x < 1$ Solution :** Given

$$f(x) = e^x \text{ in } (0,1)$$

The half range sine series for $f(x)$ in $(0,l)$ is $f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \dots\dots(1)$

$$l=1 \text{ Where } b_n = \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

$$= \frac{2}{1} \int_0^1 f(x) \sin n\pi x dx \quad b_n$$

$$= 2 \int_0^1 e^x \sin(n\pi x) dx$$

$$= 2 \frac{e^x}{(1)^2 + (n\pi)^2} (\sin n\pi x - n\pi \cdot \cos n\pi x) \text{ apply limits 0 to 1}$$

$$= \frac{2}{1+n^2\pi^2} [e^1(0 - n\pi \cdot \cos n\pi) - e^0(0 - n\pi \cdot \cos 0)]$$

$$= \frac{2}{1+n^2\pi^2} [-n\pi \cdot e \cdot \cos n\pi + n\pi]$$

$$= \frac{2}{1+n^2\pi^2} [-n\pi e(-1)^n + n\pi]$$

$$= \frac{2n\pi}{1+n^2\pi^2} [1 - e(-1)^n] \quad b_n$$

$$(1) \Rightarrow \sum_{n=1}^{\infty} \frac{2n\pi}{1+n^2\pi^2} [1 - e(-1)^n] \sin n\pi x \quad f(x)=$$

2)

$$\sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \dots\dots(1)$$

$$= \frac{2}{l} \int_0^l f(x) \sin \frac{n\pi x}{l} dx$$

Find the half
range sine
series of $f(x) =$

1 in (0,l) **Solution :** The half range sine series in
(0,l) is $f(x) =$

where b_n

$$= \frac{2}{l} \int_0^l 1 \sin \frac{n\pi x}{l} dx$$

$$= \frac{2}{l} \left[\frac{-\cos \left(\frac{n\pi x}{l} \right)}{\frac{n\pi}{l}} \right] \text{ apply limits 0 to l}$$

$$= -\frac{2}{l} \cdot \frac{l}{n\pi} [\cos n\pi - \cos 0]$$

$$= -\frac{2}{n\pi} [(-1)^n - 1]$$

$b_n = 0$ if n is even

if n is odd

Now (1) $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n\pi} \sin \frac{n\pi x}{l}$
 If n is odd

3) Find the half range cosine series of $f(x) = x(2-x)$ in the range $0 \leq x \leq 2$

Hence find sum of series $\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$

Solution : Given function $f(x) = x(2-x) = 2x - x^2$

The half range cosine series for $f(x)$ is $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l} \dots\dots\dots(1)$

where $a_0 = \frac{2}{l} \int_0^l f(x) dx = \frac{2}{2} \int_0^2 f(x) dx = \int_0^2 (2x - x^2) dx$
 $= \frac{2}{2} [\frac{2x^2}{2} - \frac{2x^3}{3}] \text{ apply 0 to 2 } = -\frac{4}{3}$

$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx$
 $= \frac{2}{2} \int_0^2 f(x) \cos \frac{n\pi x}{2} dx \quad (l=2)$
 $= \int_0^2 (2x - x^2) \cos \frac{n\pi x}{2} dx \quad \text{(using integration by parts)}$
 $= [(2x - x^2) \frac{2}{n\pi} \{ \sin \frac{n\pi x}{2} + (2-2x) \frac{4}{n^2\pi^2} \cos \frac{n\pi x}{2} + (2) \frac{8}{n^3\pi^3} \sin \frac{n\pi x}{2}]$

apply limits 0 to 2
 $= \frac{-8}{n^2\pi^2} \cos n\pi - \frac{8}{n^2\pi^2} = \frac{-8}{n^2\pi^2} [1 - (-1)^n]$

$\frac{-16}{n^2\pi^2}$ when n is even $a_n =$

= 0 when n is odd

Substitute the values of a_0 and a_n in (1) we get

$$\begin{aligned}
 (1) \Rightarrow 2x - x^2 &= \frac{2}{3} - \frac{16}{\pi^2} \sum_{n=2,4,6}^{\infty} \left(\frac{1}{n^2} \cos \frac{n\pi x}{2} \right) \\
 &= \frac{2}{3} - \frac{16}{\pi^2} \left(\frac{1}{2^2} \cos \pi x + \frac{1}{4^2} \cos 2\pi x + \frac{1}{6^2} \cos 3\pi x + \dots \right) \\
 &= \frac{2}{3} - \frac{16}{\pi^2} \cdot \frac{1}{2^2} \left(\cos \pi x + \frac{1}{2^2} \cos 2\pi x + \frac{1}{3^2} \cos 3\pi x + \dots \right) \\
 \Rightarrow 2x - x^2 &= \frac{2}{3} - \frac{4}{\pi^2} \left(\cos \pi x + \frac{1}{2^2} \cos 2\pi x + \frac{1}{3^2} \cos 3\pi x + \dots \right) \text{-----(2)}
 \end{aligned}$$

Putting $x = 1$

in (2) we get

$$\begin{aligned}
 2 - 1 &= \frac{2}{3} - \frac{4}{\pi^2} \left(\cos \pi + \frac{1}{2} \cos 2\pi + \frac{1}{3^2} \cos 3\pi + \frac{1}{4^2} \cos 4\pi + \dots \right) \\
 \Rightarrow 1 - \frac{2}{3} &= - \frac{4}{\pi^2} \left(-1 + \frac{1}{2^2} - \frac{1}{3^2} + \frac{1}{4^2} \right. \\
 &\quad \left. - \dots \right) \\
 \Rightarrow \frac{1}{3} &= \frac{4}{\pi^2} \left(1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} \right. \\
 &\quad \left. + \dots \right) \\
 + \Rightarrow \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} &\quad \dots \dots \dots) = \\
 12
 \end{aligned}$$

(4) Expand $f(x) = e^{-x}$ as Fourier series in $(-1,1)$

Solution : Here $l = 1$

$$a_0 = \frac{1}{l} \int_{-l}^l f(x) dx$$

$$= \frac{1}{1} \int_{-1}^1 e^{-x} dx = \left(\frac{e^{-x}}{-1} \right) \text{apply limits -1 to 1}$$

$$= -e^{-1} + e^1 = e - \frac{1}{e} = 2 \sinh 1$$

$$\frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx$$

$$= 1 \int_{-1}^1 e^{-x} \cos(n\pi x) dx$$

$$= \frac{e^{-x}}{(-1)^2 + (n\pi)^2} (-\cos n\pi x + n\pi$$

an =

. sin $n\pi x$) apply limits -1 to 1

$$= \frac{1}{1+n^2\pi^2} [e^{-1}\{-(-1)^n + 0\} - e^1\{-(-1)^n + 0\}]$$

$$= \frac{1}{1+n^2\pi^2} (-1)^n (e - e^{-1})$$

$$= \frac{1}{1+n^2\pi^2} (-1)^n 2\sinh 1$$

$$b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx$$

$$= \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx$$

$$= \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx \quad (l=1)$$

$$= \int_{-1}^1 e^{-x} \sin(n\pi x) dx$$

$$= \frac{e^{-x}}{(-1)^2 + (n\pi)^2} \left(\right.$$

$$= \frac{1}{1+n^2\pi^2} [e^{-1}(0 - n\pi \cdot \cos n\pi) - e^1(0 - n\pi \cdot \cos n\pi)]$$

$$= \frac{1}{1+n^2\pi^2} n\pi \cdot \cos n\pi (e - e^{-1})$$

$$= \frac{1}{1+n^2\pi^2} n\pi (-1)^n 2\sinh 1$$

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos n\pi x + \sum_{n=1}^{\infty} b_n \sin n\pi x \dots\dots\dots(1)$$

- sin nπx -

nπ . cos

nπx)

apply

limits -1

to 1

Now Fourier series of f(x)

in (-l,l) is

f(x) =

$$f(x) = \frac{2 \sinh 1}{2} + \sum_{n=1}^{\infty} \frac{1}{1+n^2\pi^2} (-1)^n 2 \sinh 1 \cos n\pi x + \sum_{n=1}^{\infty} \frac{1}{1+n^2\pi^2} n\pi (-1)^n 2 \sinh 1 \sin n\pi x$$

$$\Rightarrow f(x) = 2 \sinh 1 + \left[\frac{1}{2} + \sum_{n=1}^{\infty} \frac{1}{1+n^2\pi^2} (-1)^n \{ \cos n\pi x + n\pi \sin n\pi x \} \right]$$

• Functions having points of discontinuity: Problems:

(1) If $f(x)$ is a function with period 2π is defined by $f(x) =$

0, for $-\pi < x \leq 0$

$= x$, for $0 \leq x < \pi$ then write the fourier series for $f(x)$

Hence deduce that $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$

Solution : The Fourier series in $(-\pi, \pi)$ is $f(x) =$

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) \text{ -----(1)}$$

$$\text{Where } a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_{-\pi}^0 f(x) dx + \int_0^{\pi} f(x) dx$$

$$= \frac{1}{\pi} \left[0 + \int_0^{\pi} x dx \right] = \frac{1}{\pi} \left(\frac{x^2}{2} \right) \text{ 0 to } \pi = \frac{\pi}{2}$$

$$\begin{aligned}
 a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx \\
 &= \frac{1}{\pi} \left[0 + \int_0^{\pi} x \cos nx \, dx \right] \quad \int u \, dv = uv - \int v \, du \\
 &= \frac{1}{\pi n^2} [(-1)^n - 1] \quad u = x, \quad dv = \cos nx \, dx = 0, \text{ if } n \text{ is even} \\
 &= -\frac{2}{\pi n^2}, \text{ if } n \text{ is odd}
 \end{aligned}$$

$$\begin{aligned}
 b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx \\
 &= \frac{1}{\pi} \left[0 + \int_0^{\pi} x \sin nx \, dx \right] \\
 &= \frac{1}{\pi} \left[\left(\frac{-x \cos nx}{n} \right) - \int_0^{\pi} \frac{-\cos nx}{n} \, dx \right] \quad (\text{apply } 0 \text{ to } \pi) \\
 &= \frac{1}{\pi} \left[\left(\frac{-\pi \cos n\pi}{n} \right) + 0 + \frac{1}{n} \left(\frac{\sin nx}{n} \right) 0 \text{ to } \pi \right] \\
 &= \frac{1}{\pi} \left[\frac{-\pi(-1)^n}{n} + 0 + 0 = -\frac{(-1)^n}{n} \right] \\
 b_n &= \frac{1}{n}, \text{ if } n \text{ is odd} \\
 &= -\frac{1}{n}, \text{ if } n \text{ is even}
 \end{aligned}$$

$$(1) \Rightarrow f(x) = \frac{1}{2} \frac{\pi}{2} - \frac{2}{\pi} \left[\left(\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \dots \right) + \left(\frac{\sin x}{1} - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \dots \right) \right] \quad (2)$$

$$\text{Put } x = 0 \text{ on both sides} \quad f(0) = 0$$

$$(2) \Rightarrow 0 = \frac{\pi}{4} - \frac{2}{\pi} \left(\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right)$$

$$\frac{2}{\pi} \left(\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right) = \frac{\pi}{4}$$

$$\Rightarrow \left(\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \right) = \frac{\pi^2}{8}$$

) + 0

Problem (2) : Find Fourier series to represent the function $f(x)$ given by

$$f(x) = -k, \text{ for } -\pi < x < 0$$

$$k, \text{ for } 0 < x < \pi \text{ hence show}$$

$$\text{that } 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}$$

Solution : In

$$-\pi < x < 0$$

$$\text{i.e., } x \in (-\pi, 0), \quad f(x) = -k$$

$$f(-x) = -f(x) \text{ in } (0, \pi)$$

$$\text{In } 0 < x < \pi \text{ i.e., } x \in (0, \pi) \quad f(x)$$

$$= k \quad f(-x) = k = -$$

$$(-k)$$

$$= -f(x) \text{ in } (-$$

$\pi, 0)$ Therefore $f(x)$ is odd function in $(-\pi, \pi)$

$$\text{so } a_0 = 0, \quad a_n = 0$$

$$b_n = \frac{2}{\pi} \int_0^\pi f(x) \sin nx \, dx$$

$$\begin{aligned}
 &= \frac{2}{\pi} \int_0^{\pi} k \sin nx \, dx \\
 &= \frac{2k}{\pi} \left(\frac{-\cos nx}{n} \right) \\
 &= \frac{2k}{\pi n} [(-1)^n - 1]
 \end{aligned}$$

b_n

) apply limits 0 to π

= 0, if n is even

= $\frac{4k}{\pi n}$, if n is odd

Now $f(x) = \sum_{n=1}^{\infty} b_n \sin nx$

= $b_1 \sin 1x + b_2 \sin 2x + b_3 \sin 3x + b_4 \sin 4x + \dots$ -----f(x)

$\frac{4k}{\pi}$ $\frac{4k \sin 3x}{\pi}$

= $\pi \sin x + 0 + \frac{\pi}{3} + 0 + \dots$ -----(1)

Deduction : put $x = \frac{\pi}{2}$ on both sides in (1)

$$(1) \Rightarrow k = \frac{4k}{\pi} (1) + \frac{4k}{\pi} \left(-\frac{1}{3}\right) + \frac{4k}{\pi} \left(\frac{1}{5}\right) + \dots$$

$$\Rightarrow k = \frac{4k}{\pi} \left[1 - \frac{1}{3} + \frac{1}{5} - \dots \right]$$

$$\Rightarrow \frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \dots$$

Parseval's Formula :-

Prove That $\int_{-l}^l [f(x)]^2 dx = l \left[\frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2) \right]$

Proof :- We know that the Fourier series of $f(x)$ in $(-l, l)$ is **$f(x)$**

$$= \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{l} \text{ -----(1)}$$

Multiplying on both sides of (1) by $f(x)$ and integrate term by

$$\begin{aligned} \text{term from } -l \text{ to } l \text{ we get } \int_{-l}^l [f(x)]^2 dx = \\ \frac{a_0}{2} \int_{-l}^l f(x) dx + \sum_{n=1}^{\infty} a_n \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx \\ + \sum_{n=1}^{\infty} b_n \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx \text{ -----(2)} \end{aligned}$$

Now $a_0 = \frac{1}{l} \int_{-l}^l f(x) dx \Rightarrow \int_{-l}^l f(x) dx = l a_0$

$$a_n = \frac{1}{l} \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx \Rightarrow \int_{-l}^l f(x) \cos \frac{n\pi x}{l} dx = l a_n$$

and $b_n = \frac{1}{l} \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx \Rightarrow \int_{-l}^l f(x) \sin \frac{n\pi x}{l} dx = l b_n$

Substitute these in (2)

$$(2) \Rightarrow \int_{-l}^l [f(x)]^2 dx = \frac{a_0^2}{2} \cdot l + \sum_{n=1}^{\infty} \frac{2l}{n^2} (a_n^2 + b_n^2)$$

This is called parseval's formula.

Note 1): In (0,2l) the parseval's formula is

$$\int_0^{2l} [f(x)]^2 dx = l \left[\frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2) \right]$$

Note :2) If $0 < x < l$ (for half range cosine series of f(x)) parseval's formula is

$$\int_0^l [f(x)]^2 dx = \frac{l}{2} \left[\frac{a_0^2}{2} + \sum_{n=1}^{\infty} a_n^2 \right]$$

Note :3) If $0 < x < l$ (for half range sine series of f(x)) parseval's formula is

$$\int_0^l [f(x)]^2 dx = \frac{l}{2} \left[\sum_{n=1}^{\infty} b_n^2 \right]$$

Problem : prove that in $0 < x < l$, $x = \frac{l}{2} - \frac{4l}{\pi^2} \left(\frac{\cos \frac{\pi x}{l}}{1^2} + \frac{\cos \frac{3\pi x}{l}}{3^2} + \dots \right)$

$$\frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots = \frac{\pi^4}{96}$$

4

) and hence

deduce that

Solution : Let $f(X) = x$, $0 < X < l$

The Fourier cosine series for $f(x)$ in $(0,l)$ is

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{l} \text{----- (1)}$$

$$\text{Here } a_0 = \frac{2}{l} \int_0^l f(x) dx$$

$$= \frac{2}{l} \int_0^l x dx$$

$$= \frac{2}{l} \left[\frac{x^2}{2} \right] \text{ apply limits 0 to l}$$

=

$$dv = \cos \frac{n\pi x}{l} dx$$

$$a_n = \frac{2}{l} \int_0^l f(x) \cos \frac{n\pi x}{l} dx$$

$$= \frac{2}{l} \int_0^l x \cos \frac{n\pi x}{l} dx$$

u = x,

$\frac{n\pi x}{l}$

$$dx] = \frac{2}{l} \left[\left\{ \frac{x \sin \frac{n\pi x}{l}}{\frac{n\pi}{l}} \right\} 0 \text{ to } l - \int_0^l \frac{\sin \frac{n\pi x}{l}}{\frac{n\pi}{l}} \right]$$

$$= \frac{2}{l} \cdot \frac{l}{n\pi} [(0 - 0) - \left\{ \frac{-\cos \frac{n\pi x}{l}}{\frac{n\pi}{l}} \right\} 0 \text{ to } l]$$

$$= \frac{2}{n\pi} \cdot \frac{l}{n\pi} [\cos n\pi - \cos 0]$$

$$= \frac{2l}{n^2\pi^2} [(-1)^n - 1]$$

$$-4l \quad -4l \text{ an} = 0 ,$$

$$n \text{ is even } \quad a_1 = \frac{-4l}{\pi^2 \cdot 1^2}, \quad a_3 = \frac{-4l}{\pi^2 \cdot 3^2}$$

$$= \frac{-4l}{n^2 \pi^2}, \quad n \text{ is odd} \quad a_2 = 0, \quad a_4 = 0 \dots\dots\dots$$

Substitute a_0, a_n in (1)

$$(1) \Rightarrow \frac{l}{2} \cdot \frac{-4l}{\pi^2} \left(\frac{\cos \frac{\pi x}{l}}{1^2} + \frac{\cos \frac{3\pi x}{l}}{3^2} + \dots\dots\dots \right)$$

$$\text{Now } a_0 = l, \quad a_1 = \frac{-4l}{\pi^2 \cdot 1^2}, \quad a_3 = \frac{-4l}{\pi^2 \cdot 3^2} \dots\dots\dots$$

From Parseval's formula, we have

$$\int_0^l [f(x)]^2 dx = \frac{l}{2} \left[\frac{a_0^2}{2} + \frac{l^2}{2} \left(\frac{l}{2} + \frac{16l}{\pi^4 \cdot 1^4} + a_1^2 + \frac{16l}{\pi^4 \cdot 3^4} + \dots\dots\dots \right) a_2^2 + a_3^2 + \dots\dots\dots \right]$$

$$\Rightarrow \left(\frac{x^3}{3} \right)_0^l = \frac{l^2}{2} \left[\frac{1}{2} + \frac{16}{\pi^4 \cdot 1^4} + \frac{16}{\pi^4 \cdot 3^4} + \dots\dots\dots \right]$$

$$\Rightarrow \frac{1}{3} (2l^3) \cdot \frac{2}{l^3} = \frac{1}{2} + \frac{16}{\pi^4 \cdot 1^4} + \frac{16}{\pi^4 \cdot 3^4} + \dots$$

$$\Rightarrow \frac{2}{3} - \frac{1}{2} = \frac{16}{\pi^4} \left(\frac{1}{1^4} + \frac{1}{3^4} + \dots \right)$$

$$\Rightarrow \frac{1}{6} \cdot \frac{\pi^4}{16} = \frac{1}{1^4} + \frac{1}{3^4} + \dots$$

$$\text{There fore } \frac{1}{1^4} + \frac{1}{3^4} + \dots = \frac{\pi^4}{96}$$

COMPLEX FOURIER SERIES in $(-l, l)$ or $(0, 2l)$:-

The complex form of Fourier series of a periodic function $f(x)$ of period $2l$ is defined by

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{\frac{in\pi x}{l}} \quad \text{--- (1) where } c_n = \frac{1}{2l} \int_{-l}^l f(x) e^{\frac{-in\pi x}{l}} dx, n=0, -1, 1, 2, \dots$$

Note (1) : If period of function is 2π , i.e., in $(-\pi, \pi)$ or $(0, 2\pi)$ then complex fourier series is $f(x) = \sum_{n=-\infty}^{\infty} c_n e^{inx}$ ----(2)

$$\text{Where } c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx, n = 0, -1, 1, -2, 2, \dots$$

Problem : Find complex fourier series of $f(x) = e^x$ if $-\pi < x < \pi$ and $f(x) = f(x + 2\pi)$

$$\text{Solution : Complex fourier series of } f(x) = e^x \text{ is } f(x) = \sum_{n=-\infty}^{\infty} c_n e^{inx} \text{ ----(1)}$$

$$\text{When } c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx$$

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^x e^{-inx} dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{(1-in)x} dx$$

$$= \frac{1}{2\pi} \left[\frac{e^{(1-in)x}}{1-in} \right] \text{ limits } (-\pi, \pi) = \frac{1}{2\pi(1-in)} [e^{(1-in)\pi} - e^{(1-in)(-\pi)}]$$

$$= \frac{1}{2\pi(1-in)} [e^{\pi} \cdot e^{-in\pi} - e^{-\pi} \cdot e^{in\pi}]$$

$$= \frac{1}{2\pi} \cdot \frac{1}{(1-in)} [e^{\pi} \cdot (-1)^n - e^{-\pi} \cdot (-1)^n]$$

$$= \frac{1}{2\pi(1-in)} (-1)^n (e^{\pi} - e^{-\pi}) = \frac{(-1)^n}{1-in} \cdot \frac{e^{\pi} - e^{-\pi}}{2}$$

$$] \quad e^{in\pi} = \cos n\pi + i \sin n\pi$$

$$\sin n\pi = 0$$

$$(1-in) \cdot \frac{1}{1-in} = 1$$

$$= \frac{(-1)^n}{2\pi} \cdot \frac{1+in}{(1+n^2)} \cdot (2 \sin h \pi) \quad (\sin h \pi) \quad \text{sub in (1)}$$

$$\text{Therefore } c_n = (-1)^n \cdot \frac{1+in}{\pi(1+n^2)} \quad (\sin h \pi) e^{inx}$$

Problem : Find the complex form of the fourier

$$(1) \Rightarrow f(x) = \sum_{n=-\infty}^{\infty} (-1)^n \cdot \frac{1+in}{\pi(1+n^2)} \quad \text{series of } f(x) = \quad , -1 \leq x \leq 1$$

here(l=1)

Solution : The complex fourier series of f(x) in (-1,1) is

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{\frac{in\pi x}{l}} \text{ -----(1)}$$

$$\text{Where } c_n = \frac{1}{2} \int_{-1}^1 e^{-x} e^{-in\pi x} dx = \frac{1}{2} \int_{-1}^1 e^{-(1+in\pi)x} dx$$

$$= \frac{1}{2} \left[\frac{e^{-(1+in\pi)x}}{-(1+in\pi)} \right]_{-1}^1$$

$$= -\frac{1}{2} \cdot \frac{1}{1+in\pi} [e^{-(1+in\pi)} - e^{(1+in\pi)}]$$

$$= \frac{1}{2} \left[\frac{1-in\pi}{1+\pi^2 n^2} \right] [e^{(1+in\pi)} - e^{-(1+in\pi)}]$$

$$= \frac{1}{2} \left[\frac{1-in\pi}{1+\pi^2 n^2} \right] [e \cdot e^{in\pi} - e^{-1} \cdot e^{-in\pi}]$$

$$= \frac{1}{2} \left[\frac{1-in\pi}{1+\pi^2 n^2} \right] [(-1)^n (e - e^{-1})]$$

$$= \frac{1}{2} (-1)^n \left[\frac{1-in\pi}{1+\pi^2 n^2} \right] 2 \sin h$$

$$(1) \Rightarrow f(x) = \sum_{n=-\infty}^{\infty} (-1)^n \left[\frac{1-in\pi}{1+\pi^2 n^2} \right] \sin h \cdot e^{-in\pi x} \quad] \text{ limits } (-1,1)$$

UNIT V

FOURIER TRANSFORMS

&

Z- TRANSFORMS

-
- **FOURIER TRANSFORMS**

Fourier Integral Theorem:-

Statement : If $f(x)$ is a given function defined in $(-l,l)$ and satisfies Dirichlet's condition then $f(x) = \frac{1}{\pi} \int_0^{\infty} \int_{-\infty}^{\infty} f(t) \cos \lambda(t-x) dt d\lambda$.

The representation of $f(x)$ is known as Fourier Integral of $f(x)$

Problems on integral theorem:

(1) Express the function $f(x) = 1, |x| \leq 1$

$$= 0, -\infty < x < -1 =$$

$$0, 1 < x < \infty$$

as fourier integral and hence evaluate (i) $\int_0^{\infty} \frac{\sin \lambda \cos \lambda x}{\lambda} d\lambda$

$$(ii) \int_{-\infty}^{\infty} \frac{\sin x}{x} dx = \pi$$

• **Solution:** The Fourier Integral theorem is given by f(x)

$$= \frac{1}{\pi} \int_0^{\infty} \int_{-\infty}^{\infty} f(t) \cos \lambda(t-x) dt d\lambda.$$

$$= \frac{1}{\pi} \int_0^{\infty} \left[\int_{-1}^1 \cos \lambda(t-x) dt \right] d\lambda$$

$$= \frac{1}{\pi} \int_0^{\infty} \left[\frac{\sin \lambda(t-x)}{\lambda} \right] d\lambda \quad \text{limits } (-1 \text{ to } 1) \text{ for } t$$

$$= \frac{1}{\pi} \int_0^{\infty} \left[\frac{\sin \lambda(1-x) - \sin \lambda(-1-x)}{\lambda} \right] d\lambda$$

$$= \frac{1}{\pi} \int_0^{\infty} \left[\frac{\sin (\lambda-\lambda x) + \sin (\lambda+\lambda x)}{\lambda} \right] d\lambda$$

$$= \frac{1}{\pi} \int_0^{\infty} 2 \cdot \left[\frac{\sin \lambda \cdot \cos \lambda x}{\lambda} \right] d\lambda$$

$$\text{therefore } f(x) = \frac{2}{\pi} \int_0^{\infty} \left[\frac{\sin \lambda \cdot \cos \lambda x}{\lambda} \right] d\lambda \text{ -----(1)}$$

Deduction :

$$\begin{aligned}
 (1) \int_0^{\infty} \frac{\sin \lambda \cos \lambda x}{\lambda} d\lambda &= \frac{\pi}{2} f(x) \\
 &= \frac{\pi}{2}, \quad |x| \leq 1 \\
 &= 0, \quad |x| > 1 \quad \text{-----} \\
 &\quad (2)
 \end{aligned}$$

Put $x = 0$

$$\begin{aligned}
 (2) \Rightarrow \int_0^{\infty} \frac{\sin \lambda \cos 0}{\lambda} d\lambda &= \frac{\pi}{2} \\
 \Rightarrow \int_0^{\infty} \frac{\sin \lambda}{\lambda} d\lambda &= \frac{\pi}{2} \\
 \Rightarrow \int_0^{\infty} \frac{\sin x}{x} dx &= \frac{\pi}{2}
 \end{aligned}$$

Fourier cosine & sine Integrals:

1) Fourier cosine Integral of $f(x)$ is

$$f(x) = \frac{2}{\pi} \int_0^{\infty} \cos \lambda x \int_0^{\infty} f(t) \cos \lambda t dt d\lambda$$

2) Fourier sine Integral of $f(x)$ is

$$f(x) = \frac{2}{\pi} \int_0^{\infty} \sin \lambda x \int_0^{\infty} f(t) \sin \lambda t dt d\lambda$$

Problems:-

2) Express $f(x) = 1, 0 \leq x \leq \pi$

$0, x > \pi$ as a fourier sine integral and

Hence evaluate $\int_0^\infty \left(\frac{1 - \cos \lambda \pi}{\lambda}\right) \sin \lambda x \, d\lambda$

Solution : Fourier sine integral of $f(x)$ is given by

$$\begin{aligned}
 f(x) &= \frac{2}{\pi} \int_0^\infty \sin \lambda x \left[\int_0^\infty f(t) \sin \lambda t \, dt \right] d\lambda \\
 &= \frac{2}{\pi} \int_0^\infty \sin \lambda x \left[\int_0^\pi \sin \lambda t \, dt \right] d\lambda \\
 &= \frac{2}{\pi} \int_0^\infty \sin \lambda x \left(\frac{-\cos \lambda t}{\lambda} \right) (0 \text{ to } \pi) d\lambda \\
 &= \frac{2}{\pi} \int_0^\infty \left(\frac{1 - \cos \lambda \pi}{\lambda} \right) \sin \lambda x \, d\lambda \quad f(x) = \\
 &\Rightarrow \int_0^\infty \left(\frac{1 - \cos \lambda \pi}{\lambda} \right) \sin \lambda x \, d\lambda \quad \pi = \\
 &\quad \quad \quad 2 \quad f(x) .
 \end{aligned}$$

$$= \frac{\pi}{2} \cdot 1, 0 \leq x \leq \pi$$

$$0, x > \pi$$

Problem : 3) Using Fourier Integral show that

$$\int_0^\infty \frac{1 - \cos \lambda \pi}{\lambda} \sin \lambda x \, d\lambda = \frac{\pi}{2}, 0 < x < \pi$$

$$0, x > \pi$$

Solution : Let $f(x) = 1$, $0 \leq x \leq \pi$

0 , $x > \pi$

then write above solution (problem.(2) solution).

Problem :4) Using Fourier Integral, show that $e^{-ax} = \frac{2a}{\pi} \int_0^{\infty} \frac{\cos \lambda x}{\lambda^2 + a^2} d\lambda$

Solution : Let $f(x) = e^{-ax}$

The Fourier Cosine Integral is given by $f(x)$

$$= \frac{2}{\pi} \int_0^{\infty} \cos \lambda x \left[\int_0^{\infty} f(t) \cos \lambda t dt \right] d\lambda$$

Now $f(t) = e^{-at}$

$$e^{-ax} = \frac{2}{\pi} \int_0^{\infty} \cos \lambda x \left[\int_0^{\infty} e^{-at} \cos \lambda t dt \right] d\lambda \text{ ----(1)}$$

$$\int_0^{\infty} e^{-at} \cos \lambda t dt = \left[\frac{e^{-at}}{a^2 + \lambda^2} (-a \cos \lambda t + \lambda \sin \lambda t) \right]_0^{\infty}$$

Therefore

Now $-a \cos \lambda t + \lambda \sin \lambda t$ (0 to ∞)

$$-a.1 + 0 = \frac{-a}{a^2 + \lambda^2}$$

sub in (1)

$$(1) \Rightarrow e^{-ax} = \frac{2}{\pi} \int_0^{\infty} \cos \lambda x \cdot \frac{a}{a^2 + \lambda^2} d\lambda$$

$$= \frac{2a}{\pi} \int_0^{\infty} \frac{\cos \lambda x}{a^2 + \lambda^2} d\lambda$$

$$\frac{\pi}{2} e^{-x} = \int_0^{\infty} \frac{\cos \lambda x}{a^2 + \lambda^2} d\lambda$$

Problem 5

above problem(4)

Solution : Let $f(x) = e^{-x}$

Problem 6): Using Fourier Integral , show that

$$e^{-ax} - e^{-bx} = \frac{2(b^2 - a^2)}{\pi} \int_0^{\infty} \frac{\lambda \sin \lambda x}{(\lambda^2 + a^2)(\lambda^2 + b^2)} d\lambda \quad (a, b > 0)$$

Solution : Let $f(x) = e^{-ax}$

The Fourier Sine integral is given by f(x)

$$\frac{2}{\pi} \int_0^{\infty} \sin \lambda x \left[\int_0^{\infty} f(t) \sin \lambda t dt \right] d\lambda$$

$$\frac{2}{\pi} \int_0^{\infty} \sin \lambda x \left[\int_0^{\infty} e^{-at} \sin \lambda t dt \right] d\lambda \text{ ----(1)}$$

$$\int_0^{\infty} e^{-at} \sin \lambda t dt = \left[\frac{e^{-at}}{a^2 + \lambda^2} (-a \sin \lambda t - \lambda \cos \lambda t) \right]_0^{\infty}$$

$$= 0 - \frac{1}{a^2 + \lambda^2} (-\lambda) = \frac{\lambda}{a^2 + \lambda^2}$$

sub in (1)

$$(1) \Rightarrow f(x) = \frac{2}{\pi} \int_0^{\infty} \sin \lambda x \cdot \frac{\lambda}{a^2 + \lambda^2} d\lambda$$

$$\Rightarrow e^{-ax} = \frac{2}{\pi} \int_0^{\infty} \frac{\lambda \sin \lambda x}{\lambda^2 + a^2} d\lambda \text{ -----(2)}$$

similarly , $e^{-bx} = \frac{2}{\pi} \int_0^{\infty} \frac{\lambda \sin \lambda x}{\lambda^2 + b^2} d\lambda \text{ -----(3)}$

$$\begin{aligned} (2) - (3) &= e^{-ax} - e^{-bx} = \frac{2}{\pi} \int_0^{\infty} \lambda \sin \lambda x \left(\frac{1}{\lambda^2 + a^2} - \frac{1}{\lambda^2 + b^2} \right) d\lambda \\ &= \frac{2}{\pi} \int_0^{\infty} \lambda \sin \lambda x \left[\frac{b^2 - a^2}{(\lambda^2 + a^2)(\lambda^2 + b^2)} \right] d\lambda \\ &= \frac{2}{\pi} (b^2 - a^2) \int_0^{\infty} \frac{\lambda \sin \lambda x}{(\lambda^2 + a^2)(\lambda^2 + b^2)} d\lambda \end{aligned}$$

There fore , $e^{-ax} - e^{-bx} = \frac{2(b^2 - a^2)}{\pi} \int_0^{\infty} \frac{\lambda \sin \lambda x}{(\lambda^2 + a^2)(\lambda^2 + b^2)} d\lambda$

FOURIER TRANSFORMATION:

Definition : 1) The fourier transform of $f(x)$, $-\infty < x < \infty$ is denoted by $f(s)$ or $F\{f(x)\}$ and is defined as ,

$$F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx = f(s) \text{ -----(1)}$$

The inverse fourier transform is given by

$$f(x) = F^{-1}\{f(s)\} = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} f(s) ds \text{ -----(2)}$$

$$F\{f(x)\} = f(s)$$

Note 2): Some authors also defined as

$$F\{f(x)\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isx} f(x) dx$$

$$\text{and inverse fourier transform as } f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-isx} f(s) ds$$

$$\text{Def : 3) : } F\{f(x)\} = \int_{-\infty}^{\infty} e^{-isx} f(x) dx \quad \text{and}$$

$$\text{Inverse Fourier Transform as } f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{isx} f(s) ds$$

Def: **Fourier Sine Transform:-**

The Fourier Sine Transform of $f(x)$, $0 < x < \infty$ is denoted by $fs(s)$ or $Fs\{f(x)\}$ and defined by

$$Fs\{f(x)\} = \int_0^{\infty} f(x) \sin_{sx} dx = fs(s) \text{ -----(3)}$$

$$Fs\{f(x)\} = \int_0^{\infty} f(x) \sin_{sx} dx = fs(s) \text{ -----(3)}$$

The inverse Fourier Sine Transform is given by

$$f(x) = \frac{2}{\pi} \int_0^{\infty} fs(s) \sin_{sx} ds \text{ -----(4)}$$

Note : Some authors also defined as

$$Fs\{f(x)\} = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \sin_{sx} dx = fs(s)$$

and inverse fourier sine transform as $f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(s) \sin sx \, ds$

Def : **Fourier Cosine Transform :-**

The Fourier Cosine Transform of $f(x)$, $0 < x < \infty$ is denoted by $fc(s)$ or $Fc\{f(x)\}$ and defined by

$$Fc\{f(x)\} = \int_0^{\infty} f(x) \cos_{sx} \, dx = fc(s) \text{ -----(5) and}$$

The inverse Fourier Cosine Transform is given by,

$$f(x) = \frac{2}{\pi} \int_0^{\infty} fc(s) \cos_{sx} \, ds \text{ -----(6)}$$

Note : Some authors also defined as

$$Fc\{f(x)\} = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \cos_{sx} \, dx$$

and inverse fourier cosine transform as $f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} fc(s) \cos_{sx} \, ds$

Linear Property: If $f(s)$, $g(s)$ are Fourier Transform of $f(x)$ & $g(x)$ then

$$\begin{aligned} F\{c_1 f(x) + c_2 g(x)\} &= c_1 F\{f(x)\} + c_2 F\{g(x)\} \\ &= c_1 f(S) + c_2 g(s) \end{aligned}$$

Proof:- The definition of Fourier Transform is

$$F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx = f(s) \text{ -----(1)}$$

$$\begin{aligned} \text{By definition } F\{c_1 f(x) + c_2 g(x)\} &= \int_{-\infty}^{\infty} e^{isx} [c_1 f(x) + c_2 g(x)] dx \\ &= c_1 \int_{-\infty}^{\infty} e^{isx} f(x) dx + c_2 \int_{-\infty}^{\infty} e^{isx} g(x) dx \\ &= c_1 f(s) + c_2 g(s) \quad \text{by (1)} \end{aligned} \quad \textbf{Note:-}$$

Linear Property:

$$(I) \quad F_s\{c_1 f(x) + c_2 g(x)\} = c_1 f_s(s) + c_2 g_s(s)$$

$$(II) \quad F_c\{c_1 f(x) + c_2 g(x)\} = c_1 f_c(s) + c_2 g_c(s)$$

Proof:- (I) The definition of Fourier Sine Transform is

$$F_s\{f(x)\} = \int_0^{\infty} f(x) \sin_{sx} dx = f_s(s) \text{ -----(1)}$$

$$\begin{aligned} \text{By the definition, } F_s\{c_1 f(x) + c_2 g(x)\} &= \int_0^{\infty} [c_1 f(x) + c_2 g(x)] \sin_{sx} dx \\ &= c_1 \int_0^{\infty} f(x) \sin_{sx} dx + c_2 \int_0^{\infty} g(x) \sin_{sx} dx \\ &= c_1 f_s(s) + c_2 g_s(s) \quad \text{by (1)} \end{aligned} \quad \textbf{Change}$$

of scale property:

$$\text{Statement : If } F\{f(X)\} = f(s) \text{ then } F\{f(ax)\} = \frac{1}{a} f\left(\frac{s}{a}\right)$$

Proof :- The definition of Fourier Transform of f(x) is

$$F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx = f(s) \text{ -----(1)}$$

By definition $F\{f(ax)\} = \int_{-\infty}^{\infty} e^{isx} f(ax) dx$ let $ax = t$ $x = t/a$

$$= \int_{-\infty}^{\infty} e^{is \frac{t}{a}} \frac{1}{a} f(t) dt$$

1 $dx = dt$

$$= \frac{1}{a} \int_{-\infty}^{\infty} e^{i\left(\frac{s}{a}\right)t} f(t) dt$$

$$= \frac{1}{a} \int_{-\infty}^{\infty} e^{i\left(\frac{s}{a}\right)x} f(x) dx$$

$f(x) dx$ (by property $= \frac{1}{a} f\left(\frac{s}{a}\right)$ of def. integral)

Note : 1) If $Fs\{f(x)\} = \frac{1}{a} fs\left(\frac{s}{a}\right)$ then Fs

2) If $Fc\{f(x)\} = fc(s)$ then $Fc\{f(ax)\} = \frac{1}{a} fc\left(\frac{s}{a}\right)$

Proof: (I) The definition of Fourier Sine Transform is

$$Fs\{f(x)\} = \int_0^{\infty} f(x) \sin sx dx = fs(s) \text{ -----(1)}$$

$$= \int_0^{\infty} f(ax) \sin \left(\frac{s}{a}t\right) \frac{1}{a} dt$$

By definition $= \int_0^{\infty} f(t) \sin \left(\frac{s}{a}t\right) \frac{1}{a} dt$ let $ax = t$

1

$s \left(dt \right) dx = dt$

a

$$\begin{aligned}
 & \int_0^\infty f(t) \sin\left(\frac{s}{a}t\right) dt \\
 &= \frac{1}{a} \int_0^\infty f(x) \sin\left(\frac{s}{a}x\right) dx \\
 &= \frac{1}{a} f\left(\frac{s}{a}\right) \text{ by (1)}
 \end{aligned}$$

Shifting Property:-

If $F\{f(x)\} = f(s)$ then $F\{f(x-a)\} = e^{isa} f(s)$

Proof : $F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx = f(s)$ ----(1)

By definition $F\{f(x-a)\} = \int_{-\infty}^{\infty} e^{isx} f(x-a) dx$ let $x-a=t$ then $x=t+a$ and $dx=dt$

$$\begin{aligned}
 &= \int_{-\infty}^{\infty} e^{is(t+a)} f(t) dt \\
 &= e^{isa} \int_{-\infty}^{\infty} e^{ist} f(t) dt \\
 &= e^{isa} f(s) \text{ by (1)}
 \end{aligned}$$

Modulation Theorem :-

If $F\{f(x)\} = f(s)$ then $F\{f(x) \cos ax\} = \frac{1}{2} \{f(s-a) + f(s+a)\}$

$$= \frac{1}{2} \left[\int_{-\infty}^{\infty} e^{i(s+a)x} f(x) dx + \int_{-\infty}^{\infty} e^{i(s-a)x} f(x) dx \right]$$

Proof: The definition of Fourier

$$\text{Transform is } \cos ax \} = \int_{-\infty}^{\infty} e^{isx} f(x) \cos ax \, dx \quad F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) \, dx$$

$$= f(s) \text{-----(1)} \quad \text{By} \quad = \int_{-\infty}^{\infty} e^{isx} \frac{e^{iax} + e^{-iax}}{2} \quad \text{definition } F\{f(x)$$

$$f(x) \, dx$$

$$f(x) \, dx + f(x) \, dx$$

$$= \frac{1}{2} \{f(s-a) + f(s+a)\}$$

Note: If $F_s(s)$ & $F_c(s)$ are Fourier Sine & Cosine Transform of $f(x)$ respectively

Then (i) $F_s\{f(x) \cos ax\} = \frac{1}{2} \{F_s(s+a) + F_s(s-a)\}$

(ii) $F_s\{f(x) \sin ax\} = \frac{1}{2} \{F_s(s+a) - F_s(s-a)\}$

(iii) $F_c\{f(x) \sin ax\} = \frac{1}{2} \{F_c(s+a) - F_c(s-a)\}$

Proof: The definition of Fourier Sine Transform of $f(x)$ is

$$F_s\{f(x)\} = \int_0^{\infty} f(x) \sin sx \, dx = f(s) \text{-----(1)}$$

By definition $F_s\{f(x) \cos ax\} = \int_0^{\infty} f(x) \cos ax \sin sx \, dx$

$$= \int_0^{\infty} f(x) \sin sx \cos ax \, dx$$

$$= \int_0^\infty f(x) \cdot \frac{1}{2} \cdot (2 \sin(s-a)x \, dx]$$

$$= \frac{1}{2} f(x) \int_0^\infty [\sin(sx + ax) + \sin(sx - ax)] dx$$

$$= \frac{1}{2} \left[\int_0^\infty f(x) \sin(s+a)x \, dx + \int_0^\infty f(x) \sin(s-a)x \, dx \right]$$

$$= \frac{1}{2} [Fs(s+a) + Fs(s-a)]$$

Similarly we get (ii) & (iii) Problems:

1) Find Fourier Transform of $f(x) = e^{ikx}$, $a < x < b$

$$0, \quad x < a, \quad x > b$$

Solution : By definition, $F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) \, dx$

$$= \int_a^b e^{isx} e^{ikx} \, dx$$

$$= \int_a^b e^{i(s+k)x} \, dx$$

$$= \left[\frac{e^{i(s+k)x}}{i(s+k)} \right] \quad \text{(apply limits a to b)}$$

$$= \frac{e^{i(s+k)b} - e^{i(s+k)a}}{i(s+k)}$$

2) Find, $F\{f(x)\}$ if $f(x) = x$, $|x| < a$

$$0, |x| > a \quad |x| < a \text{ means } -a < x < a$$

Solution : By definition , $F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx$

$$= \int_{-a}^a e^{isx} x dx$$

use $\int_{-a}^a x \cdot e^{isx} dx$, integration by parts ,

$\int u dv = \left(\frac{xe^{isx}}{is}\right) - \frac{1}{is} \int_{-a}^a e^{isx}$ $uv - \int v du$

(apply -a to a)

$u=x, \quad dv=e^{isx}dx$

$$= \frac{1}{is} (a \cdot e^{ias} + a \cdot e^{-ias}) - \frac{1}{is} \left(\frac{e^{isx}}{is} \right)$$

$$= \frac{2a \cos as}{is} + \frac{1}{s^2} (e^{ias} - e^{-ias})$$

$$= \frac{-2ia \cos as}{s} + \frac{2i \sin as}{s^2}$$

$= \frac{e^{isx}}{is}$

) (apply -a to a)

$du=dx, \quad v= \int e^{isx} dx$

3) If $f(x) = 1, |x| < a$

0

, $|x| > a$, Find Fourier Transform of $f(x)$

Deduce that $\int_{-\infty}^{\infty} \frac{\sin as \cos sx}{s} ds$ (ii) $\int_{-\infty}^{\infty} \frac{\sin s}{s} ds$

(i) $\int_{-\infty}^{\infty} e^{isx} f(x) dx$

Solution : $F\{f(X)\} =$

$|x| < a$ means $-a < x < a$

$= \int_{-a}^a e^{isx} .1. dx$

$= \frac{e^{isx}}{is} (-a \text{ to } a)$

$$= \frac{1}{is}(e^{ias} - e^{-ias})$$

$$= \frac{1}{is} 2 \sin as$$

$$f(s) = \frac{1}{s} \quad F\{f(x)\} = f(s)$$

Deduction :

Inverse Fourier Transform is defined by $f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} f(s) ds$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} (\cos sx - i \sin sx) \frac{2 \sin as}{s} ds \quad f(x) =$$

$$= \frac{2}{2\pi} \left[\int_{-\infty}^{\infty} (\cos sx) \frac{\sin as}{s} ds - i \int_{-\infty}^{\infty} (\sin sx) \frac{\sin as}{s} ds \right]$$

(even)

(odd)

$$\Rightarrow f(x) = \frac{1}{\pi} \left[2 \int_{-\infty}^{\infty} (\cos sx) \frac{\sin as}{s} ds - 0 \right]$$

$$(i) \int_{-\infty}^{\infty} \frac{\sin as \cos sx}{s} ds = \frac{\pi}{2} \cdot f(x)$$

$$= \frac{\pi}{2} \begin{cases} 1, & |x| < a \\ 0, & |x| > a \end{cases}$$

(ii) Put $a = 1$, $x = 0$ in (i) we get

$$\int_0^\infty \frac{\sin s}{s} ds = \frac{\pi}{2} \cdot 1$$

$$\Rightarrow \int_0^\infty \frac{\sin s}{s} ds = \frac{\pi}{2}$$

4) Find Fourier Transform of $f(x) = 1 - x^2$, $|x| \leq 1$

$$0, |x| > 1$$

$$\int_0^\infty \left(\frac{x \cos x - \sin x}{x^3} \right) \cos \frac{x}{2} dx$$

Evaluate

$$\begin{aligned} \text{Solution:- } F\{f(x)\} &= \int_{-\infty}^{\infty} e^{isx} f(x) dx \\ &= \int_{-1}^1 e^{isx} (1 - x^2) dx \\ &= \int_{-1}^1 (1 - x^2) e^{isx} dx \\ &= \left[(1 - x^2) \cdot \frac{e^{isx}}{is} \right] - \int_{-1}^1 \frac{e^{isx}}{is} (-2x) dx \end{aligned}$$

$$u dv = uv - \int v du$$

(limits -1 to 1)

$$u = (1 - x^2) \quad dv = e^{isx} dx$$

$$0 + \frac{2}{is} \int_{-1}^1 x \cdot e^{isx}$$

$$= [0 - dx] \quad du = -2x$$

$$dx, \quad v = \frac{e^{isx}}{is}$$

$$= \frac{e^{isx}}{is}$$

$$\int_{-1}^1 \frac{e^{isx}}{is} dx]$$

$$= \frac{2}{is} \left[\left(\frac{x e^{isx}}{is} \right) (-1 \text{ to } 1) - \right]$$

$$= \frac{4}{s^3} \left[\sin s - s \cos s \right] = f(s)$$

Deduction : $= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} f(s) ds$ Inverse

$$\text{Fourier} = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} f(s) ds$$

Transform is defined by $f(x) = \int_{-\infty}^{\infty} f(s) ds$

$$\begin{aligned} & \frac{4}{s^3} [\sin s - s \cos s] ds \\ &= \frac{1}{2\pi} \cdot 4 \int_{-\infty}^{\infty} (\cos sx - i \sin sx) \frac{(\sin s - s \cos s)}{s^3} ds = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} f(s) ds \\ &= \frac{2}{\pi} \left[\int_{-\infty}^{\infty} \cos sx \frac{(\sin s - s \cos s)}{s^3} ds - i \int_{-\infty}^{\infty} \sin sx \frac{(\sin s - s \cos s)}{s^3} ds \right] \end{aligned}$$

(even function)

(odd function)

$$\begin{aligned} f(x) &= \frac{2}{\pi} \left[\int_{-\infty}^{\infty} \cos sx \frac{(\sin s - s \cos s)}{s^3} ds - 0 \right] \\ \Rightarrow \int_{-\infty}^{\infty} \cos sx \frac{(\sin s - s \cos s)}{s^3} ds &= f(x) \end{aligned}$$

$$ds = f(x)$$

$$= \frac{\pi}{2} (1 - x^2), |x| \leq 1$$

$$0, |x| > 1$$

At $x = \frac{1}{2}$, $\Rightarrow \int_{-\infty}^{\infty} \cos \frac{s}{2} \frac{(\sin s - s \cos s)}{s^3} ds = \frac{\pi}{2} (1 - \frac{1}{4})$ put

$s = x$

$$\Rightarrow \int_{-\infty}^{\infty} \cos \frac{x}{2} \frac{(\sin x - x \cos x)}{x^3} dx = \frac{\pi}{2} (1 - \frac{1}{4}) = \frac{3\pi}{8}$$

$$\Rightarrow 2 \int_0^{\infty} \cos \frac{x}{2} \frac{(\sin x - x \cos x)}{x^3} dx = \frac{3\pi}{8}$$

$$\int_0^{\infty} \cos \frac{x}{2} \left[\frac{(x \cos x - \sin x)}{x^3} \right] dx = -\frac{3\pi}{16}$$

5) Find Fourier Transform of $f(x) = \frac{1}{2a}$ if $|x| \leq a$
 0 , if $|x| > a$

Solution : By definition,

$$\begin{aligned}
 F\{f(x)\} = f(s) &= \int_{-\infty}^{\infty} e^{isx} f(x) dx \\
 &= \int_{-\infty}^{-a} e^{isx} f(x) dx + \int_{-a}^a e^{isx} f(x) dx + \int_a^{\infty} e^{isx} f(x) dx \\
 &= \int_{-a}^a \frac{1}{2a} e^{isx} dx = \frac{1}{2a} \frac{e^{isx}}{is} \quad (\text{apply limits}) = \frac{1}{2a} \frac{(e^{isa} - e^{-isa})}{is} \\
 &= \frac{\sin as}{ias}
 \end{aligned}$$

6) Find Fourier Transform of $f(x) = \sin x$, if $0 < x < \pi$
 0 , otherwise

Solution : By definition,

$$\begin{aligned}
 F\{f(x)\} = f(s) &= \int_{-\infty}^{\infty} e^{isx} f(x) dx \\
 &= \int_{-\infty}^0 e^{isx} f(x) dx + \int_0^{\pi} e^{isx} f(x) dx + \int_{\pi}^{\infty} e^{isx} f(x) dx \\
 &= \int_0^{\pi} e^{isx} \sin x dx \\
 &= \frac{e^{isx}}{(is)^2 + 1^2} [is \sin x - 1 \cdot \cos x] \quad \text{apply 0 to } \pi \\
 &= \frac{1}{1-s^2} [e^{is\pi} (0 - \cos \pi) - e^0 (0 - 1)] \\
 &= \frac{1}{1-s^2} [e^{is\pi} (1) - 1(0 - 1)] \\
 &= \frac{e^{is\pi} + 1}{1-s^2}
 \end{aligned}$$

7) Find Fourier Transform of $f(x) = xe^{-x}$, $0 < x < \infty$

Solution : By definition,

$$F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx \quad f(s) =$$

$$= \int_0^{\infty} e^{isx} x e^{-x} dx$$

$$= \int_0^{\infty} x e^{(is-1)x} dx$$

$$= \left[\frac{x e^{(is-1)x}}{is-1} - 1 \cdot \frac{e^{(is-1)x}}{(is-1)^2} \right] (0 \text{ to } \infty)$$

$$= \left[\frac{x \{e^{isx} - e^{-x}\}}{is-1} \right] (0 \text{ to } \infty) - \frac{1}{(is-1)^2} (e^{isx} - e^{-x})$$

$$= [(0-0) - \frac{1}{(is-1)^2} (0-1)]$$

$$= \frac{1}{(is-1)^2}$$

$$= \frac{1}{(is-1)^2} \cdot \frac{(is+1)^2}{(is+1)^2}$$

$$= \frac{(1+is)^2}{(1+s)^2}$$

$$-x^2$$

$$-x^2$$

8) Find Fourier Transform of $e^{-\frac{x^2}{2}}$. Show that $e^{-\frac{s^2}{2}}$ is reciprocal Solution : By definition,

$$\begin{aligned}
 F\{f(x)\} &= f(s) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isx} f(x) dx \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isx} e^{-\frac{x^2}{2}} dx \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\frac{-1}{2}(x^2 - 2isx)} dx \quad (x-is)^2/2 = y^2 \\
 &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{\frac{-1}{2}[(x-is)^2 + s^2]} dx \quad x-is = 2y \\
 &= \frac{1}{\sqrt{2\pi}} e^{\frac{-s^2}{2}} \int_{-\infty}^{\infty} e^{\frac{-1}{2}(x-is)^2} dx \quad dx = 2dy \\
 &= \frac{1}{\sqrt{2\pi}} e^{\frac{-s^2}{2}} \int_{-\infty}^{\infty} e^{-y^2} \sqrt{2} dy \\
 &= \frac{1}{\sqrt{\pi}} e^{\frac{-s^2}{2}} \int_{-\infty}^{\infty} e^{-y^2} dy \\
 &= \frac{1}{\sqrt{\pi}} e^{\frac{-s^2}{2}} 2 \int_0^{\infty} e^{-y^2} dy \\
 &= e^{\frac{-s^2}{2}} \cdot \frac{2}{\sqrt{\pi}} \cdot \frac{\sqrt{\pi}}{2} \\
 &= e^{\frac{-s^2}{2}} = f(s)
 \end{aligned}$$

Therefore Function is self reciprocal

9) Find the inverse Fourier Transform of $f(x)$ of $f(s) = e^{-|s|y}$

Solution : We have $|s| = -s$, if $s < 0$

s , if $s > 0$

From inverse Fourier Transform, we have

$$\begin{aligned}
 f(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-isx} f(s) ds \\
 &= \frac{1}{2\pi} \left[\int_{-\infty}^0 e^{-isx} f(s) ds + \int_0^{\infty} e^{-isx} f(s) ds \right] \\
 &= \frac{1}{2\pi} \left[\int_{-\infty}^0 e^{-isx} e^{sy} ds + \int_0^{\infty} e^{-isx} e^{-sy} ds \right] \\
 &= \frac{1}{2\pi} \left[\int_{-\infty}^0 e^{(y-ix)s} ds + \int_0^{\infty} e^{-(y+ix)s} ds \right] \\
 &= \frac{1}{2\pi} \left[\frac{e^{(y-ix)s}}{y-ix} \right]_{-\infty}^0 + \frac{1}{2\pi} \left[\frac{e^{-(y+ix)s}}{-(y+ix)} \right]_0^{\infty} \\
 &= \frac{1}{2\pi} \left[\frac{1}{y-ix} \right] + \frac{1}{2\pi} \left[\frac{1}{y+ix} \right] \\
 &= \frac{1}{2\pi} \left[\frac{y+ix+y-ix}{(y-ix)(y+ix)} \right] = \frac{1}{2\pi} \frac{2y}{y^2-x^2} \\
 &= \frac{1}{\pi} \frac{y}{y^2+x^2}
 \end{aligned}$$

] (0 to ∞)

Problems on sine and cosine Transform:-

1) Find Fourier cosine Transform of $f(x)$ defined by $f(x) = \cos x$, $0 < x < a$
 $= 0$, $x > a$

Solution : $F_c\{f(x)\} = \int_0^{\infty} f(x) \cos_{sx} dx$
 $= \int_0^a \cos x \cos_{sx} dx = \frac{1}{2} \int_0^a 2 \cos x \cos_{sx} dx$

$$\begin{aligned} &= \frac{1}{2} \int_0^a [\cos(x + sx)] \\ &= \frac{1}{2} \left[\int_0^a \cos(1 + s)x \, dx + \int_0^a \cos(1 - s)x \right. \\ &= \frac{1}{2} \left[\frac{\sin(1+s)x}{1+s} + \frac{\sin(1-s)x}{1-s} \right] \quad (\text{apply 0 to a}) \\ &= \frac{1}{2} \left[\frac{\sin(1+s)a}{1+s} + \frac{\sin(1-s)a}{1-s} \right] \end{aligned}$$

$$2\cos A \cos B = \cos(A+B) + \cos(A-B)$$

$$+ \cos(x - sx)] \, dx$$

$$A=x, B=sx$$

$$dx]$$

2) Find Fourier cosine Transform of $f(x)$ defined by $f(x) = x, 0 < x < 1$
 $2-x, 1 < x < 2$
 $0, x > 2$

Solution : $F_c\{f(x)\} = \int_0^{\infty} f(x) \cos sx \, dx$

$$= \int_0^1 x \cos sx \, dx + \int_1^2 (2-x) \cos sx \, dx$$

$$= \left[x \frac{\sin sx}{s} - 1 \left(-\frac{\cos sx}{s^2} \right) \right] \text{ (apply 0 to 1) } + \left[(2-x) \frac{\sin sx}{s} - (-1) \left(-\frac{\cos sx}{s^2} \right) \right]$$

$$= \left(\frac{\sin s}{s} + \frac{\cos s}{s^2} - 0 - \frac{1}{s^2} \right) + \left(0 - \frac{\cos 2s}{s^2} - \frac{\sin s}{s} + \frac{\cos s}{s^2} \right)$$

$$= \frac{2\cos s - \cos 2s - 1}{s^2}$$

$$= \frac{2\cos s - (2\cos^2 s - 1) - 1}{s^2}$$

$$= \frac{1}{s^2} (2\cos s - 2\cos^2 s)$$

$$= \frac{2}{s^2} \cos s (1 - \cos s)$$

$$= \int_0^1 f(x) \cos sx \, dx + \int_1^2 f(x) \cos sx \, dx + \int_2^{\infty} 0 \cos sx \, dx$$

)] (1to 2)

3) Find Fourier sine & cosine Transform of $2e^{-5x} + 5e^{-2x}$

Solution : Given $f(x) = 2e^{-5x} + 5e^{-2x}$

$$\begin{aligned} \text{Fs}\{f(x)\} &= \int_0^{\infty} f(x) \sin_{sx} dx \\ &= \int_0^{\infty} (2e^{-5x} + 5e^{-2x}) \sin_{sx} dx \\ &= [2 \int_0^{\infty} e^{-5x} \sin_{sx} dx + 5 \int_0^{\infty} e^{-2x} \sin_{sx} dx] \\ &= [2 \left\{ \frac{e^{-5x}}{25+s^2} (-5 \sin sx - s \cos sx) \right\} \text{ (apply 0 to } \infty) \right] sx \\ &\quad dx \end{aligned}$$

$$\begin{aligned}
 & + 5 \left\{ \frac{e^{-2x}}{4+s^2} (-2 \sin sx - s \cos sx) \right\} \text{ (apply 0 to } \infty) \} \\
 & 0) \} + 5 \{ 0 - \frac{e^0}{25+s^2} (0 - s \cos -\frac{e^0}{-4+s^2}(-s)) \} \\
 & = \left[2 \left\{ 0 - \frac{e^0}{25+s^2} (0 - s \cos -\frac{e^0}{-4+s^2}(-s)) \right\} \right] \\
 & = \left[\frac{2s}{25+s^2} + \frac{5s}{4+s^2} \right]
 \end{aligned}$$

Similarly $\frac{10}{s^2+25} + \frac{10}{s^2+4}$] (ii) $F_c\{f(x)\} = [$

4) Find Fourier cosine Transform of (i) e^{-ax}

$\cos ax$, (ii) $e^{-ax} \sin ax$ Solution

: Given $f(x) = e^{-ax} \cos ax$ (i)

$$\begin{aligned}
 F_c\{f(x)\} &= \int_0^\infty f(x) \cos sx \, dx \\
 &= \int_0^\infty e^{-ax} \cos ax \cos sx \, dx \\
 &= \frac{1}{2} \int_0^\infty e^{-ax} 2 \cos ax \cos sx \, dx \\
 &= \frac{1}{2} \left[\int_0^\infty e^{-ax} \cos (a+s)x \, dx + \int_0^\infty e^{-ax} \cos (a-s)x \, dx \right] \\
 &= \frac{1}{2} \left[\frac{e^{-ax}}{a^2 + (a+s)^2} \{-a \cos (a+s)x + (a+s) \sin (a+s)x\} \right. \\
 &\quad \left. + \frac{e^{-ax}}{a^2 + (a-s)^2} \{-a \cos (a-s)x + (a-s) \sin (a-s)x\} \right] \text{ (apply 0 to } \infty) \\
 &= \frac{1}{2} \left[\left\{ 0 - \frac{e^0}{a^2 + (a+s)^2} (-a \cos 0) \right\} + \left\{ 0 - \frac{e^0}{a^2 + (a-s)^2} (-a \cos 0) \right\} \right]
 \end{aligned}$$

(ii) Similarly $F_s\{f(x)\} = F_s\{(e^{-ax} \sin ax)\} = \frac{1}{2} \left[\frac{a}{a^2 + (s-a)^2} - \frac{a}{a^2 + (a+s)^2} \right]$

5) Find Fourier cosine & sine Transform of e^{-ax} , $a > 0$ hence

deduce (i) $\int_0^\infty \frac{\cos sx}{a^2+s^2} ds$ (ii) $\int_0^\infty \frac{s \sin sx}{a^2+s^2} ds$

Solution : Let $f(x) = e^{-ax}$

$$\begin{aligned} Fc\{f(x)\} &= \int_0^\infty f(x) \cos sx \, dx \\ &= \int_0^\infty e^{-ax} \cos sx \, dx \\ &= \left[\frac{e^{-ax}}{a^2+s^2} (-a \cos sx + s \sin sx) \right] \text{ (apply 0 to } \infty) \end{aligned}$$

$$= \left[0 - \frac{e^0}{a^2+s^2} (-a + 0) \right] = \frac{a}{a^2+s^2} = Fc(s) \text{-----(1)}$$

$$\begin{aligned} Fs\{f(x)\} &= \int_0^\infty f(x) \sin sx \, dx \\ &= \int_0^\infty e^{-ax} \sin sx \, dx \end{aligned}$$

$$= \left[\frac{e^{-ax}}{a^2+s^2} (-a \sin sx - s \cos sx) \right] \text{ (apply 0 to } \infty) = -\frac{s}{a^2+s^2} \text{-----(2)}$$

$$Fs\{f(x)\} =$$

By Inverse cosine Transform

$$\begin{aligned} f(x) &= \frac{2}{\pi} \int_0^\infty f(s) \cos sx \, ds \\ &= \frac{2}{\pi} \int_0^\infty \frac{a}{a^2+s^2} \cos sx \, ds \end{aligned}$$

$$\Rightarrow \int_0^{\infty} \frac{1}{a^2+s^2} \cos sx \, ds = \frac{e^{-ax}}{a} \cdot \frac{\pi}{2}$$

By inverse sine Transform ,

$$\begin{aligned} f(x) &= \frac{2}{\pi} \int_0^{\infty} f(s) \sin sx \, ds \\ &= \frac{2}{\pi} \int_0^{\infty} \frac{s}{a^2+s^2} \sin sx \, ds \\ \Rightarrow \int_0^{\infty} \frac{s}{a^2+s^2} \sin sx \, ds &= \frac{\pi}{2} \cdot e^{-ax} \end{aligned}$$

6) Find Fourier sine Transform of $f(x) =$

$$\begin{aligned} &\int_0^{\infty} f(x) \sin sx \, dx \\ &= \int_0^{\infty} \frac{\sin sx}{x} \, dx \text{ ----(1)} \end{aligned}$$

Solution : $F_s\{f(x)\} = \frac{\pi}{2}$

7) Find Fourier sine Transform of e^{-ax} , hence deduce that

$$x$$

Solution : $Fs\{f(x)\} = \int_0^{\infty} f(x) \sin_{sx} dx$
 $= \int_0^{\infty} \frac{e^{-ax}}{x} \sin_{sx} dx = I \text{ ---(1)}$

$$\begin{aligned} \frac{dt}{ds} &= \int_0^{\infty} \frac{e^{-ax}}{x} \cdot x \cdot \cos_{sx} dx \\ &= \int_0^{\infty} e^{-ax} \cos_{sx} dx \\ + \quad &= \left[\frac{e^{-ax}}{a^2 + s^2} (-a \cos_{sx} \quad s \sin_{sx}) \right] \text{ (apply 0} \\ \text{to} \quad &= \left[0 - \frac{e^0}{a^2 + s^2} \quad \infty \right) \\ \Rightarrow \frac{dt}{ds} &= \frac{a}{a^2 + s^2} \quad (-a + 0) \end{aligned}$$

Integrate on both sides w.r.t. s we get

$$\begin{aligned} I &= a \int \frac{1}{a^2 + s^2} ds = a \cdot \frac{1}{a} \cdot \tan^{-1} \frac{s}{a} + c \\ &= \tan^{-1} \left(\frac{s}{a} \right) + c \text{ -----(2)} \end{aligned}$$

put $s = 0$ on both sides we get {in (1) & (2)}

$$0 = \tan^{-1}(0) + c \Rightarrow 0 = 0 + c \Rightarrow c=0$$

$$I = \tan^{-1}\left(\frac{s}{a}\right) = \text{Fs}\{f(x)\}$$

8) Find Fourier cosine Transform of $\frac{1}{1^2+x^2}$, and

(ii) Fourier sine Transform of $\frac{x}{1^2+x^2}$

Solution : Let $f(x) = \frac{1}{1^2+x^2}$, We will find $\text{Fc}\{f(x)\} = \text{Fc}\left\{\frac{1}{1^2+x^2}\right\}$

$$= \int_0^\infty f(x) \cos sx \, dx$$

$$= \int_0^\infty \frac{1}{1^2+x^2} \cos sx \, dx \quad \text{Fc}\{f(x)\}$$

$$= I \quad \text{-----(1)}$$

Differentiate on both sides w.r.t s

$$\frac{dI}{ds} = \int_0^{\infty} -\frac{x \sin sx}{1+x^2} dx \text{ ----(2)}$$

$$= - \int_0^{\infty} \frac{x^2 \sin sx}{x(1+x^2)} dx$$

$$= - \int_0^{\infty} \frac{(1+x^2-1) \sin sx}{x(1+x^2)} dx$$

$$= - \left[\int_0^{\infty} \frac{\sin sx}{s} dx - \int_0^{\infty} \frac{\sin sx}{x(1+x^2)} dx \right]$$

$$\frac{dI}{ds} = -\frac{\pi}{2} + \int_0^{\infty} \frac{\sin sx}{x(1+x^2)} dx \text{ ----(3) } \quad dx \text{ Diff}$$

on both sides w.r.t 's'

We get $\frac{d^2 I}{ds^2} = \int_0^{\infty} \frac{x \cos sx}{x(1+x^2)} dx$

$$\Rightarrow \frac{d^2 I}{ds^2} = I \text{ by (1)} \Rightarrow \frac{d^2 I}{ds^2} - I = 0$$

$$\Rightarrow (D^2 - 1)I = 0 \text{ This is D.E}$$

$$\text{A.E. is } m^2 - 1 = 0$$

$$m = \pm 1$$

$$\text{solution is } I = c_1 e^s + c_2 e^{-s} \text{ ----- (4)}$$

$$\frac{dI}{ds} = c_1 e^s - c_2 e^{-s} \text{ ----- (5)}$$

$$\text{From (1) \& (4), } c_1 e^s + c_2 e^{-s} = \int_0^\infty \frac{1}{1+x^2} \cdot \cos sx \, dx$$

Put $s = 0$ on both sides

$$\begin{aligned}\Rightarrow c_1 + c_2 &= \int_0^{\infty} \frac{1}{1+x^2} dx \\ &= (\tan^{-1})(0 \text{ to } \infty) = \tan^{-1} \infty - \tan^{-1} 0 \\ &= \frac{\pi}{2} - 0\end{aligned}$$

there fore , $c_1 + c_2 = \frac{\pi}{2}$ -----(6)

From (3) & (5) , $c_1 e^s - c_2 e^{-s} = -\frac{\pi}{2} + \int_0^{\infty} \frac{\sin sx}{x(1+x^2)} dx$

$$\Rightarrow c_1 - c_2 = -\frac{\pi}{2}$$
 -----(7)

solve (6) & (7) we get $c_1 = 0$, $c_2 = \frac{\pi}{2}$ sub in (4)

$$(4) \Rightarrow I = \frac{\pi}{2} \cdot e^{-s}$$

i.e., $\text{Fc}\{f(x)\} = \text{Fc}\left\{\frac{1}{1+x^2}\right\} = \frac{\pi}{2} \cdot e^{-s}$

Now $I = \frac{\pi}{2} \cdot e^{-s}$

$$\frac{dI}{ds} = -\frac{\pi}{2} \cdot e^{-s}$$
 -----(8)

From (2) & (8) , we have

$$-\int_0^{\infty} \frac{x \sin sx}{1+x^2} dx = -\frac{\pi}{2} \cdot e^{-s}$$

$$\Rightarrow \int_0^{\infty} \left(\frac{x}{1+x^2} \right) \sin sx \, dx = \frac{\pi}{2} \cdot e^{-s}$$

Therefore $\text{Fs} \left\{ \frac{x}{1+x^2} \right\} = \frac{\pi}{2} \cdot e^{-s}$

9) Find the Inverse Fourier Cosine Transform of $f(x)$ of $f_c(s) = \frac{1}{2a} \left(a - \frac{s}{2} \right)$, $s < 2a$
 0 , $s \geq 2a$

Solution : From the inverse Fourier Cosine Transform, we have

$$\begin{aligned} f(x) &= \frac{2}{\pi} \int_0^{\infty} f_c(s) \cos sx \, ds \\ &= \frac{2}{\pi} \left[\int_0^{2a} f_c(s) \cos sx \, ds + \int_{2a}^{\infty} f_c(s) \cos sx \, ds \right] \\ &= \frac{2}{\pi} \frac{1}{2a} \int_0^{2a} \left(a - \frac{s}{2} \right) \cos sx \, ds \\ &= \frac{1}{\pi a} \left[\left\{ \left(a - \frac{s}{2} \right) \cdot \frac{\sin sx}{x} \right\} (0 \text{ to } 2a) - \int_0^{2a} \frac{\sin sx}{x} \left(-\frac{1}{2} \right) ds \right] \\ &= \frac{1}{\pi a} \left[(0-0) + \frac{1}{2} \cdot \frac{1}{x^2} (-\cos sx) \right. \\ &= \frac{1}{2\pi a x^2} (-\cos 2ax + \cos 0) \\ &= \frac{1 - \cos 2ax}{2\pi a x^2} = \frac{\sin^2 ax}{\pi a x^2} \quad (0 \text{ to } 2a) \end{aligned}$$

10) Find $f(x)$ if its Fourier Sine Transform is e^{-as}

Solution : Given $f(s) = e^{-as}$

By definition of inverse sine transform

$$\begin{aligned}
 f(x) &= \frac{2}{\pi} \int_0^{\infty} f(s) \sin_{sx} ds \\
 &= \frac{2}{\pi} \int_0^{\infty} e^{-as} \sin_{sx} ds \\
 &= \frac{2}{\pi} \left[\frac{e^{-as}}{a^2 + x^2} (-a \sin_{sx} - x \cos_{sx}) \right] (0 \text{ to } \infty) \\
 &= \frac{2}{\pi} \left[0 - \frac{1}{a^2 + x^2} (-x) \right] \\
 &= \frac{2x}{\pi(a^2 + x^2)}
 \end{aligned}$$

11) Find the Inverse Fourier Sine Transform $f(x)$ of $F_s(s) = \frac{s}{1+s^2}$
(or)

Find $f(x)$ if its Fourier sine Transform is $\frac{s}{1+s^2}$

Solution : By Fourier Inverse sine Transform $f(x) = \frac{2}{\pi} \int_0^{\infty} f(s) \sin_{sx} ds = 1$

$$f(x) = \frac{2}{\pi} \int_0^{\infty} \frac{s}{1+s^2} \sin sx \, ds = I \text{ -----(1)}$$

$$= \frac{2}{\pi} \int_0^{\infty} \left(\frac{1}{s} - \frac{1}{s(s^2+1)} \right) \sin sx \, ds$$

$$= \frac{2}{\pi} \left[\int_0^{\infty} \frac{\sin sx}{s} \, ds - \int_0^{\infty} \frac{\sin sx}{s(s^2+1)} \, ds \right]$$

$$= \frac{2}{\pi} \left[\frac{\pi}{2} - \int_0^{\infty} \frac{\sin sx}{s(s^2+1)} \, ds \right]$$

$$f(x) = 1 - \frac{2}{\pi} \int_0^{\infty} \frac{\sin sx}{s(s^2+1)} \, ds = I \text{ -----(2)}$$

diff on both sides w.r.t. x

$$\text{We get } \frac{dI}{dx} = -\frac{2}{\pi} \int_0^{\infty} \frac{s \cos sx}{s(s^2+1)} \, ds \text{ -----(3)}$$

Diff w.r.t. x

$$\frac{d^2 I}{dx^2} = -\frac{2}{\pi} \int_0^{\infty} -s \frac{\cos sx}{(s^2+1)} \, ds$$

$$= \frac{2}{\pi} \int_0^{\infty} s \frac{\cos sx}{(s^2+1)} \, ds$$

$$\frac{d^2 I}{dx^2} = I \text{ from (1)} \Rightarrow (D^2 - 1)I = 0 \text{ -----(4) is D.E.}$$

$$\text{Solution of (4) is } I = c_1 e^x + c_2 e^{-x} \text{ -----(5)}$$

$$\frac{dI}{dx} = c_1 e^x - c_2 e^{-x} \text{ -----(6)}$$

From (2) & (5)

$$\text{If } x = 0, I = 1,$$

$$\Rightarrow c_1 + c_2 = 1 \quad (5)$$

Substitute in (3) & (6)

$$(5) \Rightarrow f(x) =$$

$$c_2 e^{-x}$$

$$\Rightarrow f(x) =$$

$$\text{If } x = 0, (3)$$

$$\text{if } x = 0, (6) \Rightarrow c_1 - c_2 = -\frac{2}{\pi} (\tan^{-1} s)(0 \text{ to } \infty)$$

$$= -\frac{2}{\pi} \frac{\pi}{2} = -1$$

Now solve $c_1 + c_2 = 1$ &

$$c_1 - c_2 = -1 \quad \text{we get } c_1 = 0 \text{ \& } c_2 = 1$$

From

$$\Rightarrow \frac{dI}{dx} = -\frac{2}{\pi} \int_0^\infty \frac{1}{1+s^2} ds$$

(5)

$$I = 0 +$$

$$e^{-x}$$

Convolution: The convolution of two functions $f(x)$ & $g(x)$ over the interval

$(-\infty, \infty)$ is defined as $f * g = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) g(x-u) du$

CONVOLUTION THEOREM: If $F\{f(x)\}$ and $F\{g(x)\}$ are Fourier Transform of functions $f(x)$ and $g(x)$, then

$$\begin{aligned} F\{f(x) * g(x)\} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isx} \{f(x) * g(x)\} dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isx} \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(u) g(x-u) du \right] dx \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(u) \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isx} g(x-u) dx \right] du \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(u) \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{is(u+y)} g(y) dy \right] du \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isu} f(u) du \cdot \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{isy} g(y) dy \\ &= F\{f(x)\} * F\{g(x)\} \end{aligned}$$

Relation between Fourier and Laplace Transform:

Statement: If $f(t) = e^{-xt} g(t)$, $t > 0$ then $F\{f(t)\} = L\{g(t)\}$
 0 , $t < 0$

Proof: $F\{f(t)\} = \int_{-\infty}^{\infty} e^{ist} f(t) dt$

$$\begin{aligned}
&= \int_{-\infty}^0 e^{ist} f(t) dt + \int_0^{\infty} e^{ist} f(t) dt \\
&= 0 + \int_0^{\infty} e^{ist} e^{-xt} g(t) dt \\
&= \int_0^{\infty} e^{-(x-is)t} g(t) dt \\
&= \int_0^{\infty} e^{\rho t} g(t) dt \\
&= L\{g(t)\}
\end{aligned}$$

Fourier Transform of derivatives of a function:

Statement: If $F\{f(x)\} = f(s)$ then $F\{f^n(x)\} = (-is)^n f(s)$, if the 1^{st} (n-1) derivatives of $f(x)$ vanish identically as $x \rightarrow \pm\infty$

Proof: By definition $F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx$ -----(1)

$$\begin{aligned}
F\{f'(x)\} &= F\left\{\frac{d}{dx} f(x)\right\} \\
&= \int_{-\infty}^{\infty} e^{isx} f'(x) dx \\
&= [e^{isx} f(x)](-\infty \text{ to } \infty) - \int_{-\infty}^{\infty} f(x) \cdot is \cdot e^{isx} dx \\
&= 0 - is \int_{-\infty}^{\infty} e^{isx} f(x) dx
\end{aligned}$$

Therefore $F\{f'(x)\} = -is F\{f(x)\}$

$$F\{f'(x)\} = -is f(x) \text{ -----(2)}$$

$$\text{Now } F\{f''(x)\} = \int_{-\infty}^{\infty} e^{isx} f''(x) dx$$

$$= [e^{isx} f'(x)](-\infty \text{ to } \infty) - \int_{-\infty}^{\infty} f'(x) \cdot is \cdot e^{isx} dx$$

$$= 0 - is \int_{-\infty}^{\infty} e^{isx} f'(x) dx$$

$$= -is \cdot F\{f'(x)\}$$

$$= -is (-is) f(s) \quad \text{by (2)}$$

Therefore $F\{f''(x)\} = (-is)^2 f(s)$

Similarly we can show that $F\{f^n(x)\} = (-is)^n f(s)$

Finite Fourier Transforms :-

Definition : The Finite Fourier sine Transform of $f(x)$, $0 < x < l$ is defined by

$$F_s\{f(x)\} = (s) = \int_0^l f(x) \sin \frac{s\pi x}{l} dx \quad f_s$$

$$\text{If } 0 < x < \pi, \quad (s) = \int_0^\pi f(x) \sin \quad F_s\{f(x)\} = f_s \quad sx dx$$

The function $f(x)$ is called the inverse finite Fourier sine transform of $f_s(s)$ and is

given by $f(x) = \int ds$

$$\text{If } 0 < x < \pi, f(x) \frac{2}{l} \sum_{s=1}^{\infty} f(s) \sin \frac{s\pi x}{l} = sx$$

Definition : The finite Fourier sine Transform of $f(x)$, $0 < x < l$ is defined by

$$Fc\{f(x)\} = fc(s) = \int_0^l f(x) \cos \frac{s\pi x}{l} dx$$

$$\text{If } 0 < x < \pi, Fc\{f(x)\} = \int_0^{\pi} f(x) \cos sx dx$$

The function $f(x)$ is called inverse finite Fourier cosine transform of $fc(s)$ and is given

$$\text{by } f(x) = Fc^{-1}\{fc(s)\} = \frac{1}{l} fc(0) + \frac{2}{l} \sum_{s=1}^{\infty} fc(s) \cos \frac{s\pi x}{l}$$

$$= Fc^{-1}\{fc(s)\} = \frac{1}{\pi} fc(0) + \frac{2}{\pi} \sum_{s=1}^{\infty} fc(s) \cos sx, (0, \pi)$$

Problem :

1) Find the Fourier Finite cosine transform of $f(x) = x$, $0 < x < \pi$ **Solution** : $Fc\{f(x)\}$

$$= fc(s) = \int_0^{\pi} f(x) \cos sx dx$$

$$= \int_0^{\pi} x \cos sx dx = \left(\frac{x \sin sx}{s} \right) (0 \text{ to } \pi) - \frac{1}{s} \int_0^{\pi} \sin sx dx$$

$$= (0 - 0) - \frac{1}{s} \left(\frac{-\cos sx}{s} \right) (0 \text{ to } \pi)$$

$$s = 1, 2, 3, \dots \quad = \frac{1}{s^2} [\cos s\pi - 1]$$

$$\text{If } s = 0, f_c(s) = \frac{1}{s^2} [(-1)^s - 1]$$

$$\text{Therefore} \quad f_c(s) =$$

$$\int_0^\pi x^2 dx = \frac{x^3}{3} (0 \text{ to } \pi) = \frac{\pi^3}{3}$$

$$\frac{1}{s^2} [(-1)^s - 1], \quad s > 0$$

2) Find the Fourier

$$= \frac{\pi^2}{2}, \quad s = 0$$

Finite sine transform of $f(x)$

$$\text{Solution: } F_s(n) = \int_0^\pi x \sin nx \, dx = \frac{1}{n} \int_0^\pi x \sin nx \, dx$$

$$= \frac{1}{n} \left[x \left(\frac{-\cos nx}{n} \right) \right] (0 \text{ to } \pi) - \frac{1}{n^2} \left(\frac{-\sin nx}{1} \right) (0 \text{ to } \pi)$$

$$= \frac{1}{n} \left[-\frac{\pi}{n} \cos n\pi + 0 - 0 - 0 \right] = -\frac{1}{n^2} \cos n\pi = -\frac{1}{n^2} (-1)^n = \frac{(-1)^{n+1}}{n^2}$$

3) Find the Fourier Finite sine transform of $f(x) = x^3$ in $(0, \pi)$ Solution: By definition the finite Fourier sine Transform is

$$F_s\{f(x)\} = \int_0^\pi f(x) \sin sx \, dx$$

$$= \int_0^\pi x^3 \sin sx \, dx$$

$$\begin{aligned}
 u &= x^3 - 3x^2 + 6x - 6 \quad dv = \sin nx \, dx \quad \frac{-\cos nx}{n} - \frac{\sin nx \cos nx}{n^2} + \frac{\sin nx \cos nx}{n^3} - \frac{\sin nx}{n^4} \\
 &= \left[-x^3 \frac{\cos nx}{n} - 3x^2 \left(\frac{-\sin nx}{n^2} \right) + 6x \left(\frac{\cos nx}{n^3} \right) - 6 \left(\frac{\sin nx}{n^4} \right) \right] (0 \text{ to } \pi) \\
 &= \left[-\pi^3 \frac{\cos n\pi}{n} - 0 + 6\pi \frac{\cos n\pi}{n^3} - 0 \right] - 0 \\
 &= \frac{-\pi^3}{n} (-1)^n + \frac{6\pi}{n^3} (-1)^n \\
 &= (-1)^n \frac{\pi}{n} \left[\frac{6}{n^2} - \pi^2 \right], \quad n = 1, 2, 3, \dots
 \end{aligned}$$

4) Find Finite sine Transform of $f(x) = x$ in $0 < x < 4$

$$\begin{aligned}
 \text{Solution : Let } f(x) \text{ is } F_s\{f(x)\} &= \int_0^4 f(x) \sin \frac{n\pi x}{4} \, dx \\
 &= \left[x \left(-\frac{\cos \frac{n\pi x}{4}}{\frac{n\pi}{4}} \right) - \left(-\frac{\sin \frac{n\pi x}{4}}{\frac{n^2\pi^2}{16}} \right) \right] (0 \text{ to } 4) \\
 &= -\frac{4}{n\pi} \cdot 4 \cdot \cos n\pi - 0 + \frac{16}{n^2\pi^2} (0 - 0) \\
 &= -\frac{16}{n\pi} \cos n\pi = -\frac{16}{n\pi} (-1)^n
 \end{aligned}$$

$$\text{Similarly } F_c\{f(x)\} = \frac{16}{n^2\pi^2} [(-1)^n - 1] = f_c(n)$$

if $n = 0$, $\int_0^4 x \, dx = \left(\frac{x^2}{2}\right) (0 \text{ to } 4) = 8$

Parseval's Identity for Fourier Transforms :-

Statement : If $f(s)$ & $g(s)$ are Fourier Transform of $f(x)$ & $g(x)$ respectively then

(i) $\frac{1}{2\pi} = \int_{-\infty}^{\infty} f(x) g(x) \, dx$

(ii) $\frac{1}{2\pi} \int_{-\infty}^{\infty} |f(s)|^2 \, ds = \int_{-\infty}^{\infty} |f(x)|^2 \, dx$

Now (iii) $\frac{2}{\pi} \int_{-\infty}^{\infty} f^c(s) g_c(s) \, ds = \int_0^{\infty} f(x) g(x) \, dx$

Proof : By the inverse Fourier Transform we have

$$g(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} g(s) e^{-isx} \, ds \text{ -----(1)}$$

Taking conjugate Complex on both sides in (1)

$$(1) \Rightarrow g(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} g(s) e^{isx} \, ds$$

$$\begin{aligned} \int_{-\infty}^{\infty} f(x) g(x) \, dx &= \int_{-\infty}^{\infty} f(x) \left[\frac{1}{2\pi} \int_{-\infty}^{\infty} g(s) e^{isx} \, ds \right] \, dx \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} g(s) \left[\int_{-\infty}^{\infty} f(x) e^{isx} \, dx \right] \, ds \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} g(s) f(s) \, ds \end{aligned}$$

$$\Rightarrow \frac{1}{2\pi} \int_{-\infty}^{\infty} f(s) g(s) \, ds = \int_{-\infty}^{\infty} f(x) g(x) \, dx \text{ -----(2)}$$

dx dx]

ds

(ii) Putting $g(x) = f(x)$ in (2) we get

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} f(s) f(s) ds = \int_{-\infty}^{\infty} f(x) f(x) dx$$

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} |f(s)|^2 ds = \int_{-\infty}^{\infty} |f(x)|^2 dx \text{ ----- Therefore (3)}$$

For Sine Transform:

$$(2) \Rightarrow \frac{2}{\pi} \int_0^{\infty} f(s) g(s) ds = \int_0^{\infty} f(x) g(x) dx$$

$$\frac{2}{\pi} \int_0^{\infty} |f(s)|^2 ds = \int_0^{\infty} |f(x)|^2 dx$$

Similarly for Cosine

Problem 1):) If $f(x) = 1$, $|x| < a$

0, $|x| > a$, Find Fourier Transform of $f(x)$

$$\int_0^{\infty} \frac{\sin ax}{x^2} dx = \frac{\pi a}{2}$$

Deduce that

Solution : $F\{f(x)\} = \int_{-\infty}^{\infty} e^{isx} f(x) dx$ $|x| < a$ means $-a < x < a$

$$= \int_{-a}^a e^{isx} . 1. dx$$

$$= \frac{e^{isx}}{is} \Big|_{-a}^a$$

$$= \frac{1}{is} (e^{ias} - e^{-ias}) = \frac{1}{is} (2i \sin as)$$

$$= \frac{2 \sin as}{s} = f(s)$$

$$F\{f(x)\} = f(s)$$

By Parseval's identity for Fourier Transform

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |f(s)|^2 ds$$

$$\Rightarrow \int_{-a}^a 1 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} \left(\frac{2 \sin as}{s}\right)^2 ds$$

$$\Rightarrow \int_{-a}^a 1 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{4 \sin^2 as}{s^2} ds$$

$$\Rightarrow 2a = \frac{2}{\pi} \int_{-\infty}^{\infty} \frac{\sin^2 as}{s^2} ds$$

$$\Rightarrow \int_{-\infty}^{\infty} \frac{\sin^2 as}{s^2} ds = a\pi$$

$$\Rightarrow \int_{-\infty}^{\infty} \left(\frac{\sin as}{s}\right)^2 ds = a\pi$$

$$\Rightarrow 2 \cdot \int_0^{\infty} \frac{\sin^2 as}{s^2} ds = a\pi$$

$$\int_0^{\infty} \frac{\sin^2 as}{s^2} ds = \frac{a\pi}{2}$$

Therefore $\int_{-\infty}^{\infty} \frac{\sin^2 as}{s^2} ds = a\pi$

2) Find Fourier Transform of $f(x) = 1 - x^2$, $|x| \leq 1$

$$0, |x| > 1 \quad \text{is} \quad \frac{4}{s^3} [\sin s - s \cos s]$$

Using Parseval's

Identity Prove That

$$\int_0^\infty \left[\frac{(\sin x - x \cos x)}{x^3} \right]^2 dx = \frac{\pi}{15}$$

Solution :-

$$\int_{-\infty}^{\infty} e^{isx} f(x) dx$$

$$= \int_{-1}^1 e^{isx} (1 - x^2) dx$$

$$= \int_{-1}^1 (1 - x^2) e^{isx} dx$$

$$F\{f(x)\} =$$

$$\int u dv = uv - \int v du \quad = \left[(1 - x^2) \cdot \frac{e^{isx}}{is} \right] - \int_{-1}^1 \frac{e^{isx}}{is} (-2x) dx \quad v du$$

(limits -1 to 1)

$$u = (1 - x^2) \quad dv = e^{isx} dx$$

$$du = -2x dx, \quad v = \frac{e^{isx}}{is}$$

$$= \left[0 - 0 + \frac{2}{is} \int_{-1}^1 x \cdot e^{isx} dx \right]$$

$$= \frac{e^{isx}}{is}$$

$$= \frac{2}{is} \left[\left(\frac{x e^{isx}}{is} \right) (-1 \text{ to } 1) - \int_{-1}^1 \frac{e^{isx}}{is} dx \right]$$

$$= \frac{2}{is} \left[1 \cdot \left(\frac{e^{is} + e^{-is}}{is} \right) - \frac{1}{is} \frac{e^{isx}}{is} \right] (-1 \text{ to } 1)$$

$$= \frac{2}{is} \left[\frac{2 \cos s}{is} - \frac{1}{is} \left(\frac{e^{is} - e^{-is}}{is} \right) \right]$$

$$= \frac{2}{is} \cdot \frac{1}{is} (2 \cos s - \frac{1}{is} 2i \sin s)$$

$$= -\frac{2}{s^2} \cdot 2 \left[\cos s - \frac{\sin s}{s} \right]$$

$$= \frac{4}{s^3} [\sin s - s \cos s] = f(s)$$

$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |f(s)|^2 ds \quad \int_{-1}^1 (1 - x^2)^2 dx =$$

]

By parseval's identity for
Fourier Transform

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \left[\frac{4}{s^3} (\sin s - s \cos s) \right]^2 ds \quad 2. \int_0^1 (1 - x^2)^2 dx = \frac{1}{2\pi} \Rightarrow \int_0^{\infty} \left[\frac{(\sin s - s \cos s)}{s^3} \right]^2 ds =$$

$$. 2. 16 \int_0^{\infty} \left[\frac{(\sin s - s \cos s)}{s^3} \right]^2 ds \quad ds =$$

$$\Rightarrow \frac{16}{\pi} \int_0^{\infty} \left[\frac{(\sin s - s \cos s)}{s^3} \right]^2 ds = 2 \cdot \frac{8}{15} \Rightarrow \int_0^{\infty} \left[\frac{(\sin x - x \cos x)}{x^3} \right]^2 dx =$$

$$\frac{\pi}{\pi} \int_0^{\pi} \left[\frac{(\sin x - x \cos x)}{x^3} \right]^2 dx =$$

$$15$$

Shifting Properties:-

1. Shifting $f(n)$ to the right :-


If $Z[f(n)] = F(Z)$ then $Z[f(n-k)] = Z^{-k}F(Z)$

Proof: we know that

$$Z[f(n)] =$$

$$\begin{aligned} & \sum_{n=0}^{\infty} f(n-k)Z^{-n} && (k, n \text{ are different forms}) \\ &= \sum_{n=k}^{\infty} f(n-k)Z^{-n} && (\text{since we are shifting } f(n) \text{ to right}) \\ &= f(0)Z^{-k} + f(1)Z^{-(k+1)} + f(2)Z^{-(k+2)} + \dots \\ &= Z^{-k}[f(0) + f(1)Z^{-1} + f(2)Z^{-2} + \dots] \\ &= Z^{-k} \sum_{n=0}^{\infty} f(n)Z^{-n} \\ &= Z^{-k}F(Z) \end{aligned}$$

$$\sum_{n=0}^{\infty} f(n)Z^{-n} \text{ consider } Z[f(n-k)] = Z^{-k}F(Z)$$



$$Z[f(n-k)] = Z^{-k}F(Z)$$

NOTE :- $Z[f(n-k)] = Z^{-k}F(Z)$ putting $k=1$, we have

$$Z[f(n-1)] = Z^{-1}F(Z) \text{ putting } k=2, \text{ we have } Z[f(n-2)] = Z^{-2}F(Z)$$

putting $k=3$, we have

$$Z[f(n-3)] = Z^{-3}F(Z)$$

2. Shifting $f(n)$ to left :-

$$\text{If } Z[f(n)] = F(Z) \text{ then } Z[f(n+k)] = Z^k[F(Z) - f(0) - f(1)Z^{-1} - f(2)Z^{-2} - \dots - f(k-1)Z^{-(k-1)}]$$

$$\text{Proof: we know that } Z[f(n)] = \sum_{n=0}^{\infty} f(n)Z^{-n} \quad [Z^{-n} = Z^k \cdot Z^{-(n+k)}]$$


$$Z[f(n+k)] = \sum_{n=0}^{\infty} f(n+k)Z^{-(n+k)}$$

$$\text{consider } Z[f(n+k)] = Z^k \sum_{n=k}^{\infty} f(n+k)Z^{-(n+k)}$$

$$= Z^k \sum_{n=k}^{\infty} f(n)Z^{-n} \quad (\text{replace } (n+k) \text{ by } n)$$

$$= Z^k [\sigma_{n=0}^{\infty} f(n)Z^{-n} - \sigma_{n=0}^{(k-1)} f(n)Z^{-n}]$$

$$= Z^k [Z[f(n)] - \sigma_{n=0}^{(k-1)} f(n)Z^{-n}]$$



$$Z[f(n+k)] = Z^k[F(Z) - f(0) - f(1)Z^{-1} - f(2)Z^{-2} - \dots - f(k-1)Z^{-(k-1)}] \quad \text{which is Recurrence formula} \quad \therefore$$

In particular

$$(a) \text{ If } k=1 \text{ then } Z[f(n+1)] = Z[F(Z) - f(0)]$$

$$(b) \text{ If } k=2 \text{ then } Z[f(n+2)] = Z^2[F(Z) - f(0) - f(1)Z^{-1}]$$

(c) If $k=3$ then $Z[f(n+3)] = Z^3[F(Z) - f(0) - f(1)Z^{-1} - f(2)Z^{-2}]$ ----- and so on.

Problems: 1. Prove $Z\left(\frac{1}{n+1}\right) = Z \log\left(\frac{Z}{Z-1}\right)$

$$\frac{1}{(n+1)} = Z \log\left(\frac{Z}{Z-1}\right)$$

we know that $Z[f(n)] = \sum_{n=0}^{\infty} f(n)Z^{-n}$ Solution- let $f(n) = Z\left(\frac{1}{n+1}\right)$

$$\frac{1}{n+1} = \sum_{n=0}^{\infty} \frac{1}{n+1} Z^{-n}$$

$$= \sum_{n=0}^{\infty} \frac{1}{n+1} \cdot \frac{1}{Z^n}$$

$$= \frac{1}{1} \cdot \frac{1}{1} + \frac{1}{2} \cdot \frac{1}{Z} + \frac{1}{3} \cdot \frac{1}{Z^2} + \dots$$

$Z[$

expansion needs 'Z' in

denominator's, for this, multiply & divide with 'Z'

$]$

$$\left[x + \frac{x^2}{2} + \frac{x^3}{3} + \dots = -\log(1-x) \right]$$

$$= Z\left[\frac{1}{Z} + \frac{1}{2} \cdot \frac{1}{Z^2} + \frac{1}{3} \cdot \frac{1}{Z^3} + \frac{1}{4} \cdot \frac{1}{Z^4} + \dots\right] \quad \text{evaluate (a) } Z\left(\frac{1}{Z} + \frac{1}{2} \cdot \frac{1}{Z^2} + \frac{1}{3} \cdot \frac{1}{Z^3} + \frac{1}{4} \cdot \frac{1}{Z^4} + \dots\right)$$

$$= Z\left[\frac{1}{Z} + \frac{\left(\frac{1}{Z}\right)^2}{2} + \frac{\left(\frac{1}{Z}\right)^3}{3} + \dots\right]$$

$$= Z\left[-\log\left(1 - \frac{1}{Z}\right)\right]$$

$$= Z\left[\log\left(1 - \frac{1}{Z}\right)^{-1}\right]$$

$$= Z\log\left(\frac{Z-1}{Z}\right)^{-1}$$

$$= Z\log\left(\frac{Z}{Z-1}\right)$$

\therefore hence proved

2. Find $Z\left[\frac{1}{n!}\right]$ and using shifting theorem

$$\frac{1}{(n+1)!} \text{ and (b) } Z\left(\frac{1}{(n+2)!}\right)$$

$$= 1 + \frac{1}{1!} Z^{-1} + \frac{1}{2!} Z^{-2} + \frac{1}{3!} Z^{-3} + \dots$$

$$= 1 + \frac{1}{Z} + \frac{\left(\frac{1}{Z}\right)^2}{2!} + \frac{\left(\frac{1}{Z}\right)^3}{3!} + \dots$$

$$= e^{\frac{1}{Z}}$$

$$- [e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \dots]$$

$= F(Z)$ (say) By

shifting theorem

$$\rightarrow Z[f(n+1)] = Z[F(Z) - F(0)]$$

$$Z \left[\left(\frac{1}{(n+1)!} \right) - \left(\frac{1}{n!} \right) \right] \quad (1)$$

$$(1) \quad Z \left[\frac{1}{(n+1)!} \right] = Z \left[e^{\frac{1}{Z}} - 1 \right] \quad [f(0) = \frac{1}{0!} = 1]$$

$$(2) \quad Z \left[\frac{1}{(n+2)!} \right] = Z^2 \left[e^{\frac{1}{Z}} - 1 - \frac{1}{1!} Z^{-1} \right]$$

$$= Z^2 \left[e^{\frac{1}{Z}} - 1 - Z^{-1} \right]$$

$$f(n) = \frac{1}{n!} \rightarrow Z[f(n+2)] = Z \left[\frac{1}{(n+2)!} \right] = Z^2 \left[e^{\frac{1}{Z}} - 1 - \frac{1}{1!} Z^{-1} \right]$$

$$f(n+1) = \frac{1}{(n+1)!}$$

$$f(n+2) = \frac{1}{(n+2)!}$$

!

$$f(n)] = -Z \frac{d}{dz} [F(Z)]$$

Proof:- we know that $Z[f(n)] = \sum_{n=0}^{\infty} f(n)Z^{-n}$

$$\therefore Z[nf(n)] = -Z \frac{d}{dz} Z[f(n)] = -Z \sum_{n=0}^{\infty} n f(n) Z^{-n-1}$$

$$= -Z \sum_{n=0}^{\infty} \frac{d}{dz} [f(n) Z^{-n}]$$

pb) If $F(Z) = \frac{(Z-1)^4}{Z^2(5+3Z^{-1}+12Z^{-2})}$ then find the values of $f(2)$ and $f(3)$

$$= -Z \frac{d}{dz} [Z f(n)]$$

$$F(Z)$$

$$= -Z \frac{d}{dz} \frac{(Z-1)^4}{Z^2(5+3Z^{-1}+12Z^{-2})}$$

$$= \frac{1}{Z^2} \frac{(5+3Z^{-1}+12Z^{-2})}{(1-Z^{-1})^4}$$

Solution: Given $F(Z) =$

By Initial value theorem we have

Multiplication by 'n': If $Z[f(n)] = F(Z)$ then

$Z[nf$

$$[Z^{-n} = Z^1 \cdot Z^{-n-1}]$$

$$\left[\frac{d}{dz} (Z^{-n}) = (-n) Z^{-n-1} \right]$$

$$5Z^2+3Z+12$$

$$5Z^2+3Z+12$$

$$f(0)=\lim_{Z\rightarrow\infty} F(Z) = 0 \quad \left(\frac{1}{\infty} = 0\right) \quad \xrightarrow{\hspace{1cm}} \quad 1$$

$$f(1)=\lim_{Z\rightarrow\infty} Z[f(Z)-f(0)] = 0$$

$$f(2)=\lim_{Z\rightarrow\infty} Z^2[F(Z) - f(0) - f(1)Z^{-1}]$$

$$=5 - 0 - 0$$

$$=5$$

$$f(3)=\lim_{Z\rightarrow\infty} Z^3[F(Z)-f(0)-f(1)Z^{-1} -f(2)Z^{-2}]$$

$$= \lim_{Z\rightarrow\infty} Z^3[F(Z)- (0) -(0.Z^{-1}) - 5Z^{-2}]$$

$$= \lim_{Z\rightarrow\infty} Z^3\left[\frac{5Z^2+3Z+12}{(Z-1)^4} - \frac{5}{Z^2}\right]$$

$$= \lim_{Z\rightarrow\infty} Z^3 5Z^4+3Z^3+2(12Z-Z1^2)-45 Z-1^4$$

$$= \lim_{Z \rightarrow \infty} Z^3 23 - 18Z^3 - [1 + -20Z - Z^{-2} 4 - 5Z^{-3}] = 23$$

$$\begin{aligned} \rightarrow (Z-1)^4 &= (z-1)^2 \cdot (z-1)^2 \\ &= (Z^2+1-2Z)(Z^2+1-2Z) \\ &= Z^4+Z^2-2Z^3+Z^2+1-2Z-2Z^3-2Z+4Z^2=Z^4+ \\ &\quad 6Z^2-4Z^3-4Z+1 \end{aligned}$$

$$[g(0) + g(1)Z^{-1} + g(2)Z^{-2} + g(3)Z^{-3} + \dots + g(n)Z^{-n} + \dots]$$

$$= \sum_{n=0}^{\infty} [f(0)g(n) + f(1)g(n-1) + f(2)g(n-2) + \dots + f(n)g(0)]Z^{-n}$$

*We have $Z[f(n)] = F(Z)$ which can be also written as $f(n) = Z^{-1}[F(Z)]$.

Then $f(n)$ is called inverse Z-transform of $F(Z)$

*Thus finding the sequence $\{f(n)\}$ from $F(Z)$ is defined as Inverse Z-Transform.

If $Z^{-1}[F(Z)] = f(n)$ and $Z^{-1}[G(Z)] = g(n)$ then

$$Z^{-1}[F(Z).G(Z)] = f(n) * g(n) = \sum_{m=0}^n f(m)g(n-m)$$

*The symbol Z^{-1} is the Inverse Z – Transform.

CONVOLUTION

Proof:- We have $F(Z) = \sum_{n=0}^{\infty} f(n)Z^{-n}$ and $G(Z) = \sum_{n=0}^{\infty} g(n)Z^{-n}$ then

THEOREM(v.v.imp):-

[where * is convolution operator]

$$F(Z).G(Z) = [f(0) + f(1)Z^{-1} + f(2)Z^{-2} + f(3)Z^{-3} + \dots + f(n)Z^{-n} + \dots]$$

$$= Z[f(0)g(n) + f(n)g(n-1) + \dots + f(n)g(0)] Z^{-1}[F(Z).G(Z)]$$

$$= f(0)g(n) + f(n)g(n-1) + \dots + f(n)g(0)$$

$$= \sum_{m=0}^n f(m)g(n-m)$$

$$\therefore Z^{-1}[F(Z).G(Z)] = f(n) * g(n) = \sum_{m=0}^n f(m)g(n-m)$$

Problems:-

1. Evaluate (a) $Z^{-1} \left[\left(\frac{Z}{Z-a} \right)^2 \right]$ (b) $Z^{-1} \left[\frac{Z^2}{(Z-a)(Z-b)} \right]$

Solution:-

(a) $Z^{-1} \left[\left(\frac{Z}{Z-a} \right)^2 \right]$

$$= Z^{-1} \left[\frac{Z}{Z-a} \cdot \frac{Z}{Z-a} \right]$$

$$F(Z) = \frac{Z}{Z-a} \Rightarrow f_n = Z^{-1} \frac{Z}{Z-a} = a_n$$

$$G(Z) = \frac{Z}{Z-a} \Rightarrow g_n = Z^{-1} \frac{Z}{Z-a} = a_n$$

by convolution theorem ,

$$Z^{-1} \left[\left(\frac{Z}{Z-a} \right) \left(\frac{Z}{Z-a} \right) \right]$$

$$= \sum_{m=0}^n a_m \cdot a_{n-m}$$

$$= \sum_{m=0}^n a_m \cdot a_{n-m}$$

$$g((n)) = \sum_{m=0}^n Z^{-1} F$$

$$f(m)g(n-m) \quad (1)$$

$$Z^{-1}[F(Z).G(Z)] = f(n) * g(n) = \sum_{m=0}^n f(m)g(n-m)$$

$$= \sigma_{n_m=0} a_m . b_{n-m}$$

$$= \sum_{m=0}^n b^m (ab)^m$$

$$= b^n \sum_{m=0}^n (ab)^m$$

$$= b^n \left[\frac{(a)^0}{b} + \frac{(a)^1}{b} + \frac{(a)^2}{b} + \frac{(a)^3}{b} + \dots + \frac{(a)^n}{b} \right]$$

this is in geometric progression,

$$1 + ar + ar^2 + \dots + ar^{n-1} + \dots = \frac{a(1-r^n)}{1-r}, \quad r < 1$$

$$= \frac{a(r^n - 1)}{1-r}, \quad r > 1$$

$$\frac{b^n \left[1 - \frac{a}{b} \right]}{1 - \frac{a}{b}}$$

Put a=3

$$Z^{-1} \left[\frac{b^n \left[\frac{b^{n+1} - a^{n+1}}{n+1} \right]}{(Z-a)(Z-b)} \right] = \frac{b^n \cdot \frac{b^{n+1} - a^{n+1}}{n+1}}{b^{n+1} - a^{n+1}} \cdot \frac{b}{b-a}$$

show that

$$\frac{1}{(Z-a)(Z-b)} = \frac{1}{b-a} \left[\frac{b-a}{Z-a} - \frac{b-a}{Z-b} \right]$$

Solution: $f(n) = \frac{b^n}{n!}$ $g(n) = \frac{a^n}{n!}$

$$= \frac{b^n}{b-a}$$

and b=4 we get

$$4^{n+1} - 3^{n+1} = 4^{n+1} - 3^{n+1} \quad Z^{-3} Z^{-4} 4^{-3}$$

2. Using Convolution theorem

$$Z^{-1} \left[\frac{1}{n!} * \frac{1}{n!} \right] = \frac{1}{n!} \quad \text{where } * \text{ is convolution operator}$$

$$\begin{aligned}
f(n) * g(n) &= \sum_{m=0}^n f(m)g(n-m) \\
&= \sum_{m=0}^n \frac{1}{m!} \cdot \frac{1}{(n-m)!} \\
&= 1 \cdot \frac{1}{n!} + \frac{1}{1!} \cdot \frac{1}{(n-1)!} + \frac{1}{2!} \cdot \frac{1}{(n-2)!} + \dots + \frac{1}{n!} \cdot \frac{1}{(0)!} \\
&= \frac{1}{n!} + \frac{1}{(n-1)!} + \frac{1}{2!} \cdot \frac{1}{(n-2)!} + \dots + \frac{1}{n!} \quad \left[\frac{1}{(n-1)!} = \frac{n}{n(n-1)!} = \frac{1}{(n-1)!} \right] \\
&= \frac{1}{n!} + \frac{1}{1!} \frac{n}{n!} + \frac{1}{2!} \frac{n(n-1)}{n!} + \dots + \frac{1}{n!} \quad ! \\
&= \frac{1}{n!} \left[1 + \frac{n}{1!} + \frac{n(n-1)}{2!} + \dots \right] \\
&= \frac{1}{n!} (1 + 1)^n \\
&= \frac{2^n}{n!} \quad \text{- to (n+1) terms]
\end{aligned}$$

3. Evaluate $Z^{-1} \left[\frac{Z^2}{(Z-4)(Z-5)} \right]$

Solution- Given $Z^{-1} \left[\frac{Z}{Z-4} \cdot \frac{Z}{Z-5} \right]$

$$F(Z) = \frac{Z}{Z-4} \Rightarrow f(n) = Z^{-1} \left[\frac{Z}{Z-4} \right] = 4^n \quad [G(Z)] = f(n) * g(n) = \sum_{m=0}^n f(m)g(n-m)$$

$$[G(Z) = \frac{Z}{Z-5}] \Rightarrow g(n) = Z^{-1} \left[\frac{Z}{Z-5} \right] = 5^n$$

by convolution theorem, $Z^{-1}[F(Z) \cdot G(Z)] = \sum_{m=0}^n 4^m \cdot 5^{n-m}$

$$Z^{-1} \left[\frac{Z}{Z-4} \cdot \frac{Z}{Z-5} \right] = \sum_{m=0}^n 5^n \cdot (45)_m$$

$$= 5^n \sum_{m=0}^n (45)_m$$

$$= 5^n \left[\binom{4}{5}^0 + \binom{4}{5}^1 + \binom{4}{5}^1 + \binom{4}{5}^3 + \dots + \binom{4}{5}^n \right]$$

$$= 5^n \left[1 + \frac{4}{5} + \left(\frac{4}{5} \right)^2 + \left(\frac{4}{5} \right)^3 + \dots + \left(\frac{4}{5} \right)^n \right]$$

this is in geometric progression,

$$\frac{1 + ar^3 + \dots + ar^{n-1} + \dots}{1-r} = a(1-r^n), \quad r < 1 \quad a + ar$$

$$a(r^n - 1)$$

$$\begin{aligned}
&= \frac{5^n \left[1 - \left(\frac{4}{5} \right)^{n+1} \right]}{1 - \frac{4}{5}} \\
&= \frac{5^n \left[1 - \frac{4^{n+1}}{5^{n+1}} \right]}{\frac{5-4}{5}} \\
&= \frac{5^n \left[\frac{5^{n+1} - 4^{n+1}}{5^{n+1}} \right]}{\frac{5-4}{5}} \\
&= 5^n \cdot \frac{5^{n+1} - 4^{n+1}}{5^{n+1}} \cdot \frac{5}{1} \\
&= \frac{5^{n+1} - 4^{n+1}}{1} \\
&= 5^{n+1} - 4^{n+1}
\end{aligned}$$

$$\left[\frac{\quad}{Z_{2+11}Z_{2+24}} \right] (\text{non repeated linear factors}) \ (v. imp)$$

1. Find Z^{-1} $\left[\begin{array}{c|c} \hline \hline \end{array} \right]$ _____

Solution:- let $F(Z) = \frac{1}{Z^2+11Z+24} = \frac{1}{(Z+3)Z(Z+8)}$

$$\text{then } \frac{F(Z)}{Z} = \frac{1}{(Z+3)(Z+8)} = \frac{A}{(Z+3)} + \frac{B}{(Z+8)} \rightarrow 1$$

$$= \frac{1}{(Z+3)(Z+8)} = \frac{A(Z+8)+B(Z+3)}{(Z+3)(Z+8)}$$

$$= 1 = A(Z+8)+B(Z+3) \rightarrow 2$$

$$\text{put } Z=-8 \Rightarrow 1 = A(-8+8) + B(-8+3)$$

$$1 = B(-5)$$

$$B = -\frac{1}{5}$$

$$\text{put } Z=-3 \Rightarrow 1 = A(-3+8) + B(-3+3)$$

$$1 = A(5)$$

$$A = \frac{1}{5}$$

$$5$$

$$\{Z+8=0 \Rightarrow Z=-8 \text{ \& } Z+3=0 \Rightarrow z=-3\}$$

now substitute A and B values in equation -1 we get

$$\left[\frac{F(Z)}{Z} \right] = \frac{1}{5(Z+3)} - \frac{1}{5(Z+8)}$$

$$\begin{aligned} F(Z) &= \frac{Z}{5(Z+3)} - \frac{Z}{5(Z+8)} \\ &= Z^{-1} \left[\frac{Z}{5(Z+3)} - \frac{Z}{5(Z+8)} \right] \\ &= \frac{1}{5} \left[Z^{-1} \left[\frac{Z}{Z+3} \right] - Z^{-1} \left[\frac{Z}{Z+8} \right] \right] \\ &= \frac{1}{5} [(-3)^n - (-8)^n] \end{aligned}$$

2. Find $\therefore Z^{-1} \left[\frac{Z}{Z^2 + 11Z + 24} \right] = \frac{1}{5} [(-3)^n - (-8)^n]$ the Inverse Z-Transform of _____

$$\frac{(Z-1)(Z-2)}{Z}$$

Solution:- let $F(Z) = \frac{Z}{(Z-1)(Z-2)}$ here we can resolve $F(Z)$ into partial fractions directly as follows

$$\begin{aligned} F(Z) &= Z \left[\frac{1}{(Z-1)(Z-2)} \right] = Z \left[\frac{1}{Z-2} - \frac{1}{Z-1} \right] \\ F(Z) &= \frac{Z}{Z-2} - \frac{Z}{Z-1} \end{aligned}$$

$$\begin{aligned} \text{hence } Z^{-1}[F(Z)] &= Z^{-1} \left[\frac{Z}{Z-2} \right] - Z^{-1} \left[\frac{Z}{Z-1} \right] \\ &= 2_n - 1_n \end{aligned}$$

$$Z^{-2} Z^{-1}$$

$$\left[\frac{1}{(5Z-1)(5Z+2)} \right]$$

Solution:- let $F(Z) = \frac{Z(3Z+1)}{(5Z-1)(5Z+2)}$ then

$$\frac{F(Z)}{Z} = \frac{3Z+1}{(5Z-1)(5Z+2)} = \frac{A}{5Z-1} + \frac{B}{5Z+2} \rightarrow 1 \text{ (by partial fractions)}$$

$$\frac{3Z+1}{(5Z-1)(5Z+2)} = \frac{A(5Z+2)+B(5Z-1)}{(5Z-1)(5Z+2)}$$

$$3Z+1 = A(5Z+2)+B(5Z-1)$$

$$\text{put } Z = -\frac{2}{5} \Rightarrow A = \frac{8}{15}$$

$$\text{put } Z = \frac{1}{5} \Rightarrow B = \frac{1}{15}$$

substituting A and B values in equation-1 we get

$$\frac{F(Z)}{Z} = \frac{8}{15} \frac{1}{5Z-1} + \frac{1}{15} \frac{1}{5Z+2}$$

$$\frac{F(Z)}{Z} = \frac{8}{15} \frac{1}{\left(Z - \frac{1}{5}\right)} + \frac{1}{15} \frac{1}{\left(Z + \frac{2}{5}\right)}$$

3. Find Z^{-1} of $\frac{3Z^2+Z}{3Z^2+Z}$

$$\text{hence } F(Z) = \frac{8}{75} \cdot \frac{Z}{\left(Z - \frac{1}{5}\right)} + \frac{1}{75} \cdot \frac{Z}{\left(Z + \frac{2}{5}\right)}$$

$$Z^{-1}[F(Z)] = Z^{-1} \left[\frac{8}{75} \left(\frac{Z}{Z-0.2} \right) + \frac{1}{75} \left(\frac{Z}{Z+0.4} \right) \right]$$

$$\frac{8}{75} Z^{-1} \left(\frac{Z}{Z-0.2} \right) + \frac{1}{75} Z^{-1} \left(\frac{Z}{Z-(-0.4)} \right)$$

$$\frac{8}{75} (0.2)^n + \frac{1}{75} (-0.4)^n$$

$$\therefore Z^{-1} \left[\frac{3Z^2 + Z}{(5Z-1)(5Z+2)} \right] = \frac{8}{75} (0.2)^n + \frac{1}{75} (-0.4)^n$$

=

=

Geometric Progression:a)

Finite –

$$a + ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n = \frac{a(1-r^{n+1})}{1-r}$$

b)

$$a + ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n + \dots = \frac{a}{a-r}$$

Infinite –

$$1 + r + r^2 + r^3 + \dots + r^n + \dots = \frac{1}{1-r}$$

eg; 1

4. Find $Z^{-1} \left[\frac{Z}{(Z+3)^2(Z-2)} \right]$ (repeated Linear factor of form $(ax + b)^2$ times)

Solution:- let $F(Z) = \frac{Z}{(Z+3)^2(Z-2)}$

$$\frac{F(Z)}{Z} = \frac{1}{(Z+3)^2(Z-2)}$$

$$\frac{F(Z)}{Z} = \frac{1}{(Z+3)^2(Z-2)} = \frac{A}{Z-2} + \frac{B}{Z+3} + \frac{c}{(Z+3)^2} \rightarrow 1$$

$$\frac{1}{(Z+3)^2(Z-2)} = \frac{A(Z+3)^2 + B(Z-2)(Z+3) + c(Z-2)}{(Z-2)(Z+3)^2}$$

$$1 = A(Z+3)^2 + B(Z-2)(Z+3) + c(Z-2) \quad \{ Z-2=0 \Rightarrow Z=2 \text{ \& } Z+3=0 \Rightarrow Z=-3 \}$$

put $Z=2 \Rightarrow 1=A(2+3)^2$

$$1 = A(25)$$

$$A = \frac{1}{25}$$

$$\text{put } Z=-3 \Rightarrow 1=c(-3-2)$$

$$1 = -5c \quad c =$$

$$\frac{-1}{5}$$

now comparing the co-efficients of Z^2 on both sides

$$0=A+B$$

$B = \frac{-1}{25}$ substituting A, B and C

values in equation-1, we get

$$\frac{F(Z)}{Z} = \frac{1}{(Z+3)^2(Z-2)} = \frac{1}{25} \cdot \frac{1}{Z-2} - \frac{1}{25} \cdot \frac{1}{Z+3} - \frac{1}{5} \cdot \frac{1}{(Z+3)^2}$$

$$F(Z) = \frac{1}{25} \cdot \frac{Z}{Z-2} - \frac{1}{25} \cdot \frac{Z}{Z+3} - \frac{1}{5} \cdot \frac{Z}{(Z+3)^2}$$

$$Z^{-1} \left[\frac{Z}{(Z+3)^2(Z-2)} \right] = Z^{-1} \left[\frac{1}{25} \cdot \frac{Z}{Z-2} - \frac{1}{25} \cdot \frac{Z}{Z+3} - \frac{1}{5} \cdot \frac{Z}{(Z+3)^2} \right]$$

$$= \frac{1}{25} \left[\frac{Z}{Z-2} \right] - \frac{1}{25} \left[\frac{Z}{Z+3} \right] - \frac{1}{5} \left[\frac{Z}{(Z+3)^2} \right]$$

$$\therefore Z^{-1} \left[\frac{Z}{(Z+3)^2(Z-2)} \right] = \frac{1}{25} \left[\frac{Z}{Z-2} \right] - \frac{1}{25} \left[\frac{Z}{Z+3} \right] - \frac{1}{5} \left[\frac{Z}{(Z+3)^2} \right]$$

Solutions Of Difference Equations

Difference Equations:-

Just as the Differential equations are used for dealing with continuous process in nature, the difference equations are used for dealing of discrete process.

Definition:-

A difference equation is a relation between the difference of an unknown function at one (or) more general value of the argument.

thus $\Delta y_n + 2y_n = 0$ and
 $\Delta^2 y_n + 5\Delta y_n + 6y_n = 0$ are difference equations

Solution:-

The solution of a difference equation is an expression for y_n which satisfies the given difference equation

General Solution:-

The general solution of a difference equation is that in which the number of arbitrary constants is equal to the order of the difference equation.

Linear Difference Equation:-

The Linear difference equation is that in which $y_{n+1}, y_{n+2}, y_{n+3} - - - - -$ etc occur to the 1st degree only and are not multiplied together.

The difference equation is called Homogeneous if $f(n)=0$, Otherwise it is called as NonHomogeneous equation (i.e:- $f(n) \neq 0$)

Working rule (or) Working Procedure:-

To solve a given linear difference equation with constant co-efficient by Z-transforms.

Step-1 :- Let $Z(y_n) = Z[y(n)] = Y(Z)$

Step-2 :- Take Z-Transform on both sides of the given difference equation.

Step-3 :-Use the formulae $Z(y_n) = Y(Z)$

$$Z[y_n + 1] = Z[Y(Z) - y_0]$$

$$Z[y_n + 2] = Z^2[Y(Z) - y_0 - y_1 Z^{-1}]$$

Step-4:-Simplify and find $Y(Z)$ by transposing the terms to the right and dividing by the co-efficient of $y(Z)$.

Step-5:-Take the Inverse Z-Transform of $Y(Z)$ and find the solution y_n

This gives y_n as a function of n which is the desired solution. Problems:-

1.Solve $y_{n+1} - 2y_n = 0$ using Z –Transforms.

Solution:-let $Z[y_n] = Y(Z)$

$$Z[y_{n+1}] = Z[Y(Z) - y_0]$$

taking Z-Transform of the given equation we get $Z[y_{n+1}] - 2Z[y_n] = 0$

$$Z[Y(Z) - y_0] - 2Y(Z) = 0$$

$$Y(Z)[Z - 2] = y_0$$

$$Y(Z) = \frac{y_0}{Z - 2}$$

$$[Z(a^n) = \frac{Z}{Z - a}]$$

$$Y(Z) = \frac{y_0}{Z - 2}$$

$$Z^{-1}[Y(Z)] = Z^{-1}\left[\frac{y_0}{Z - 2}\right]$$

$$y_n = 2^n y_0$$

$$\Rightarrow Z[Y(n)] = Y(Z)$$

$$Z^{-1}[Y(Z)] = y_n$$

2.Solve the difference equation using Z-Transforms

$$\mu_{n+2} - 3\mu_{n+1} + 2\mu_n = 0 \text{ Given that}$$

$$\mu_0 = 0, \mu_1 = 1$$

Solution:-let $Z(\mu_n) = \mu Z$

$$Z(\mu_{n+1}) = Z[\mu Z] - \mu_0$$

$Z(\mu_{n+2}) = Z^2 \mu Z - \mu_0 - \mu Z^1$ now taking Z-Transform on both sides of the given equation we get

$$Z(\mu_{n+2}) - 3Z(\mu_{n+1}) + 2Z(\mu_n) = 0 \quad Z^2 - \mu_0 - \mu Z^1$$

$$[Z^2 - 3Z + 2] \mu Z - \mu_0 = 0 \text{ using the given}$$

conditions it reduces to

$$[Z^2 - 3Z + 2] \mu Z - \mu_0 = 0$$

$$Z^2 - 3Z + 2 = 0$$

$$Z = \frac{3 \pm \sqrt{9 - 8}}{2} = \frac{3 \pm 1}{2}$$

$$= (Z - 1)(Z - 2)$$

$$= Z[Z - 1 - Z - 2]$$

$$\underline{Z} \quad \underline{Z}$$

$$= Z^{-2} - Z^{-1}$$

on taking Inverse Z-Transform on both sides we get

$$Z^{-1} \mu Z = Z^{-1} \left[\frac{Z}{Z-2} - \frac{Z}{Z-1} \right]$$

$$\mu^n = Z^{-1} \left[Z^2 \right] - Z^{-1} \left[Z^{Z-1} \right] \quad 2$$

$$\mu_n = 2n - 1$$

3. Solve the difference equation using Z-Transform

$$y_{n+2} - 4y_{n+1} + 3y_n = 0$$

Given that $y_0 = 2$ and $y_1 = 4$

Solution:- let $Z[y_n] = Y(Z)$

$$Z[y_{n+1}] = Z \left[Y(Z) \right] y_0 \quad Z[y_{n+2}] = Z^2 Y(Z) - y_0 - y_1 Z^{-1}$$

taking Z-Transform of the given equation we get

$$Z(y_{n+2}) - 4Z(y_{n+1}) + 3Z(y_n) = 0$$

$$Z^2 \left[Y(Z) - y_0 - y_1 Z^{-1} \right] - 4 Z Y(Z) - y_0 + 3Y(Z) = 0 \text{ using}$$

$$\text{the given conditions it reduces to}$$

$$Z^2 \left[Y(Z) - 2 - 4Z^{-1} \right] - 4 Z Y(Z) - 2 + 3Y(Z) = 0$$

$$\text{i.e:- } Y(Z)[Z^2 - 4Z + 3] - 2Z^2 - 4Z + 8Z = 0$$

$$Y(Z)[Z^2 - 4Z + 3] = Z(2Z - 4)$$

$$\frac{Y(Z)}{Z} = \frac{2Z - 4}{[Z^2 - 4Z + 3]}$$

$$= \frac{2Z - 4}{(Z - 1)(Z - 3)}$$

$$\frac{Y(Z)}{Z} = \frac{1}{Z - 1} + \frac{1}{Z - 3} \quad (\text{reducing by partial fractions})$$

$$Y(Z) = \frac{Z}{Z - 1} + \frac{Z}{Z - 3}$$

on taking Inverse Z-Transform on both sides we obtain

$$Z^{-1}[Y(Z)] = Z^{-1} \left[\frac{Z}{Z - 1} \right] + Z^{-1} \left[\frac{Z}{Z - 3} \right]$$

$$y_n = 1 + 3^n$$